

THE GERSTENHABER-SCHACK COMPLEX FOR PRESTACKS

DINH VAN HOANG AND WENDY LOWEN

ABSTRACT. The aim of this work is to construct a complex which through its higher structure directly controls deformations of general prestacks, building on the work of Gerstenhaber and Schack for presheaves of algebras. In defining a Gerstenhaber-Schack complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ for an arbitrary prestack \mathcal{A} , we have to introduce a differential with an infinite sequence of components instead of just two as in the presheaf case. If $\tilde{\mathcal{A}}$ denotes the Grothendieck construction of \mathcal{A} , which is a \mathcal{U} -graded category, we explicitly construct inverse quasi-isomorphisms \mathcal{F} and \mathcal{G} between $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ and the Hochschild complex $\mathbf{C}_{\mathcal{U}}(\tilde{\mathcal{A}})$, as well as a concrete homotopy $T : \mathcal{F}\mathcal{G} \rightarrow 1$, which had not been obtained even in the presheaf case. As a consequence, by applying the Homotopy Transfer Theorem, one can transfer the dg Lie structure present on the Hochschild complex in order to obtain an L_∞ -structure on $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$, which controls the higher deformation theory of the prestack \mathcal{A} . This answers the open problem about the higher structure on the Gerstenhaber-Schack complex at once in the general prestack case.

1. INTRODUCTION

Throughout the introduction, let k be a field. In [8], [9], [10] Gerstenhaber and Schack define the Hochschild cohomology of a presheaf \mathcal{A} of k -algebras over a poset \mathcal{U} as an Ext of bimodules $HH^n(\mathcal{A}) = \text{Ext}_{\mathcal{A}\text{-}\mathcal{A}}^n(\mathcal{A}, \mathcal{A})$, in analogy with the case of k -algebras. They construct a complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ which computes this cohomology, obtained as the totalization of a double complex with horizontal Hochschild differential and vertical simplicial differential. From \mathcal{A} , they construct a single k -algebra $\mathcal{A}!$ such that

$$(1.1) \quad HH^n(\mathcal{A}) \cong HH^n(\mathcal{A}!)$$

for the standard Hochschild cohomology of $\mathcal{A}!$ on the right hand side. Further, the authors construct two explicit cochain maps

$$(1.2) \quad \tau : \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}) \longrightarrow \mathbf{C}^\bullet(\mathcal{A}!) \quad \text{and} \quad \hat{\tau} : \mathbf{C}^\bullet(\mathcal{A}!) \longrightarrow \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$$

relating their complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ to the Hochschild complex $\mathbf{C}^\bullet(\mathcal{A}!)$, which they prove to be inverse quasi-isomorphisms. They present two essentially different approaches to (1.1), (1.2) and the relationship between the two:

- (A1) In a first approach [8], [9], (1.1) follows from their (difficult) Special Cohomology Comparison Theorem (SCCT) which compares more general bimodule Ext groups. Both sides of (1.1) are particular cases of such Ext groups, and a universal delta functor argument shows that the isomorphism (1.1) is actually induced by the map τ in (1.2), whence the latter is a quasi-isomorphism.
- (A2) In a second approach [10], in case \mathcal{U} is a finite poset, the authors focus on the compositions $\hat{\tau}\tau$ and $\tau\hat{\tau}$. They prove directly that $\hat{\tau}\tau = 1$, and a comparison of lifts of resolutions implies that $\tau\hat{\tau}$ and the identity are homotopic. Thus, in this case the isomorphism (1.1) follows without invoking the SCCT.

The authors acknowledge the support of the European Union for the ERC grant No 257004-HHNcdMir and of the Research Foundation Flanders (FWO) under Grant No G.0112.13N.

Although deformation theory for presheaves of algebras was thoroughly studied by Gerstenhaber and Schack, the following problems remained:

- (P1) Unlike in the algebra case, first order deformations of \mathcal{A} as a presheaf are classified by the second cohomology group $HH_s^2(\mathcal{A})$ of the sub-complex of $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ containing simple cochains. In general, as we have $HH_s^2(\mathcal{A}) \neq HH^2(\mathcal{A})$, this raises the question as to the precise role of the entire Gerstenhaber-Schack complex in deformation theory.
- (P2) It is a fundamental principle of deformation theory in characteristic 0, due to Deligne, that every deformation problem is governed by a dg Lie algebra or, by extension, an L_∞ -algebra. So it is natural to ask what higher structure is present on the Gerstenhaber-Schack complex.

Concerning (P2), when the poset \mathcal{U} is finite, Gerstenhaber and Schack's argument in (A2) proves the existence of a homotopy $\tau\hat{\tau} \rightarrow 1$, yet it is not constructive. Thus the Homotopy Transfer Theorem (HTT) implies the existence of an L_∞ -structure on $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ transferred from the dg Lie algebra structure on $\mathbf{C}(\mathcal{A})$, yet it does not provide the tools to make this structure concrete. More recently, using operadic methods Y. Frégier, M. Markl and D. Yau constructed an explicit L_∞ -structure on the Gerstenhaber-Schack complex of a morphism of algebras [7], which corresponds to the special case of a presheaf over a single arrow. However, the general problem (P2) has remained open both as regards existence and construction of the higher structure. One of the main results in our paper is the construction of a concrete homotopy map (Theorem 4.13) in the general prestack case which resolves this open problem.

Concerning (P1), in our recent joint work with L. Liu [3], the second cohomology of $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ is shown to classify deformations of \mathcal{A} as a *twisted presheaf*, as is seen from direct inspection of the complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$.

Another way to understand the occurrence of twists is by viewing a presheaf of algebras as a prestack, that is a pseudofunctor taking values in k -linear categories (algebras are considered as one object categories). If \mathcal{A} is a prestack over a small category \mathcal{U} , then \mathcal{A} has an associated \mathcal{U} -graded category $\tilde{\mathcal{A}}$, obtained through a k -linear version of the Grothendieck construction from [1]. If \mathcal{A} is a presheaf over a poset, then $\tilde{\mathcal{A}}$ and $\mathcal{A}!$ are closely related. In [12] it was shown based upon the construction of $\tilde{\mathcal{A}}$ that the appropriate \mathcal{U} -graded Hochschild complex $\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}})$ of $\tilde{\mathcal{A}}$ satisfies

$$(1.3) \quad H^n \mathbf{C}_{\mathcal{U}}(\tilde{\mathcal{A}}) = \text{Ext}_{\tilde{\mathcal{A}}-\tilde{\mathcal{A}}}^n(\tilde{\mathcal{A}}, \tilde{\mathcal{A}})$$

and controls deformations of $\tilde{\mathcal{A}}$ as a \mathcal{U} -graded category ($\text{Def}_{\mathcal{U}}(\tilde{\mathcal{A}})$) and, equivalently, deformations of \mathcal{A} as a prestack ($\text{Def}_{\text{pre}}(\mathcal{A})$). Further, in [14], Lowen and Van den Bergh prove a Cohomology Comparison Theorem (CCT) for prestacks \mathcal{A} . If we define $HH^n(\mathcal{A}) = \text{Ext}_{\mathcal{A}-\mathcal{A}}^n(\mathcal{A}, \mathcal{A})$ and $HH_{\mathcal{U}}^n(\tilde{\mathcal{A}}) = \text{Ext}_{\tilde{\mathcal{A}}-\tilde{\mathcal{A}}}^n(\tilde{\mathcal{A}}, \tilde{\mathcal{A}})$, it follows in particular from the CCT that

$$(1.4) \quad HH^n(\mathcal{A}) \cong HH_{\mathcal{U}}^n(\tilde{\mathcal{A}}),$$

that is, the analogue of (1.1) holds.

All of the above suggests that it is most natural to work at once in the context of arbitrary prestacks \mathcal{A} . In particular, it should be possible to define a Gerstenhaber-Schack complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ which is directly seen to control prestack deformations of \mathcal{A} , and such that we can define a new version of the inverse quasi-isomorphisms (1.2) above in this setup. Realizing this is the main goal of this paper. In summary, we have the following picture of the references in which various relations are studied

for a prestack \mathcal{A} , where $[*]$ stands for the present paper:

$$\begin{array}{ccccc} \text{Ext}_{\mathcal{A}-\mathcal{A}}(\mathcal{A}, \mathcal{A}) & & \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}) & \overset{[*]}{\rightsquigarrow} & \text{Def}_{\text{pre}}(\mathcal{A}) \\ \left. \vphantom{\text{Ext}_{\mathcal{A}-\mathcal{A}}(\mathcal{A}, \mathcal{A})} \right\} [11] & & \left. \vphantom{\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})} \right\} [*] & & \left. \vphantom{\text{Def}_{\text{pre}}(\mathcal{A})} \right\} [10] \\ \text{Ext}_{\tilde{\mathcal{A}}-\tilde{\mathcal{A}}}(\tilde{\mathcal{A}}, \tilde{\mathcal{A}}) & \overset{[10]}{\rightsquigarrow} & \mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}) & \overset{[10]}{\rightsquigarrow} & \text{Def}_{\mathcal{U}}(\tilde{\mathcal{A}}) \end{array}$$

The content of the paper is as follows. After recalling basic terminology on prestacks and map-graded categories in §2, the complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ for a prestack \mathcal{A} on a small category \mathcal{U} is defined in §3. As a graded module, according to (3.12), $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ is the totalization of a double object which is a modification of the one in the presheaf case. Precisely, we put

$$\mathbf{C}^{p,q}(\mathcal{A}) = \prod \text{Hom}_k(\mathcal{A}(U_p)(A_{q-1}, A_q) \otimes \cdots \otimes \mathcal{A}(U_p)(A_0, A_1), \mathcal{A}(U_0)(\sigma^\# A_0, \sigma^* A_q)).$$

Here, the product is taken over all p -simplices

$$(1.5) \quad \sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \cdots \xrightarrow{u_{p-1}} U_{p-1} \xrightarrow{u_p} U_p)$$

in the nerve of \mathcal{U} and all $(q+1)$ -tuples (A_0, \dots, A_q) of objects in $\mathcal{A}(U_p)$. Further, if we denote, for $u : V \rightarrow U$ in \mathcal{U} , the associated restriction functor by $u^* : \mathcal{A}(U) \rightarrow \mathcal{A}(V)$, then we put $\sigma^* = (u_p \dots u_2 u_1)^*$ and $\sigma^\# = u_1^* u_2^* \dots u_p^*$.

The occurrence of the twists significantly complicates the definition of the differential. Precisely, we have to introduce an infinite family of components $(d_j)_{j \geq 0}$ with

$$d_j : \mathbf{C}_{\text{GS}}^{p,q}(\mathcal{A}) \rightarrow \mathbf{C}_{\text{GS}}^{p+j, q+1-j}(\mathcal{A}),$$

and for each n , we define

$$(1.6) \quad d = d_0 + d_1 + \cdots + d_n : \mathbf{C}_{\text{GS}}^{n-1}(\mathcal{A}) \rightarrow \mathbf{C}_{\text{GS}}^n(\mathcal{A}).$$

We have $d_0 = d_{\text{Hoch}}$ for the horizontal Hochschild differential d_{Hoch} and $d_1 = (-1)^n d_{\text{simp}}$ for the vertical simplicial differential d_{simp} . The additional components d_j of d , given in (3.18), are necessary to make the differential square to zero, as is shown in Theorem 3.9. Note that the algebraic structure of the prestack \mathcal{A} naturally corresponds to an element

$$(m, f, c) \in \mathbf{C}^{0,2}(\mathcal{A}) \oplus \mathbf{C}^{1,1}(\mathcal{A}) \oplus \mathbf{C}^{2,0}(\mathcal{A}) = \mathbf{C}_{\text{GS}}^2(\mathcal{A})$$

with m encoding compositions, f encoding restrictions, and c encoding twists. Our definition of the components d_j ensures the following desired result (Theorem 3.20), of which the proof makes use of normalized reduced cochains as defined in §3.4:

Theorem 1.1. *The second cohomology group $H^2 \mathbf{C}_{\text{GS}}(\mathcal{A})$ classifies first order deformations of \mathcal{A} as a prestack.*

The definition of the higher components d_j is combinatorial in nature. It makes essential use of the following ingredients:

- So called *paths* of natural transformations between $\sigma^\#$ and σ^* , each path building up a $(p-1)$ -simplex in the nerve of $\text{Fun}(\mathcal{A}(U_p), \mathcal{A}(U_0))$ by using one twist isomorphism in each step (the precise definition is given in the beginning of §3.3).
- The natural action of shuffle permutations on nerves of categories, as discussed in §3.1.

In §4 we go on to define cochain maps

$$(1.7) \quad \mathcal{F} : \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}) \rightarrow \mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}) \quad \text{and} \quad \mathcal{G} : \mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}) \rightarrow \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$$

between $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$ and the \mathcal{U} -graded Hochschild complex $\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}})$ from [12].

Our main theorem is the following (see Proposition 4.9 and Theorem 4.6):

Theorem 1.2. *The maps \mathcal{F} and \mathcal{G} are inverse quasi-isomorphisms. More precisely*

- (1) $\mathcal{G}\mathcal{F}(\phi) = \phi$ for any normalized reduced cochain ϕ ;
- (2) there is an explicit homotopy $T : \mathcal{F}\mathcal{G} \sim 1$.

In combination with (1.4) and (1.3), we thus obtain

Corollary 1.3. $H^n \mathbf{C}_{\text{GS}}(\mathcal{A}) = \text{Ext}_{\mathcal{A}-\mathcal{A}}^n(\mathcal{A}, \mathcal{A})$.

Note that, in contrast with [9], in our setup we do not give a direct proof of Corollary 1.3, whence the approach (A1) is not available to us.

Although the existence of the maps \mathcal{F} and \mathcal{G} is inspired by the existence of the maps in (1.1), due to our more complicated differential on $\mathbf{C}_{\text{GS}}(\mathcal{A})$, the maps in (1.7) are actually new and the development of the appropriate combinatorial tools in order to prove Theorem 1.2 constitutes the technical heart of the paper.

Our construction of the homotopy $T : \mathcal{F}\mathcal{G} \sim 1$ in part (2) is new even in the presheaf case and has the following important consequence. By the Homotopy Transfer Theorem [11, Theorem 10.3.9], using T we can transfer the dg Lie algebra structure present on $\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}})$ (see [12]) in order to obtain an L_∞ -structure on $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$. This L_∞ -structure determines the higher deformation theory of \mathcal{A} as a prestack, which thus becomes equivalent to the higher deformation theory of the \mathcal{U} -graded category $\tilde{\mathcal{A}}$ described in [12]. A more detailed elaboration of this L_∞ -structure, as well as a comparison with the L_∞ deformation complex described in the literature in an operadic context [7], [5], [15] is work in progress [4].

In future work, we intend to extend the techniques and constructions in this paper in order to shed new light on the difficulties arising in Shresta and Yetter's deformation theories of monoidal categories and pasting diagrams [16] [17] and Elgueta's deformation theory of monoidal bicategories [6]. In particular, in [16], after introducing the components d_0, d_1 of a desired differential on the Yetter complex of monoidal categories, Shresta describes the components d_2, d_3 for cochains of low degrees, and conjectures that there is an infinite family of components constituting a differential on this complex. By applying shuffle products of morphisms and natural transformations as described in Section 3.1, we succeeded in describing the differentials d_2, d_3 on cochains of arbitrary higher degrees, and we found the formula of the differential d_4 . Details will appear in [2].

Acknowledgement. The second author is very grateful to Michel Van den Bergh for many interesting discussions, and in particular for his proposal of map-graded Hochschild cohomology which was originally made in the context of a local-to-global spectral sequence [13]. Both authors are grateful to Jim Stasheff for valuable comments on an earlier version of the paper, and thank both Jim Stasheff and an anonymous referee for suggesting an illuminating interpretation of our paths, currently explained after Example 3.5.

2. PRESTACKS AND MAP-GRADED CATEGORIES

Let k be a commutative ground ring. Except for certain Ext interpretations of cohomologies, notably the ones occurring in (4.1), all our results hold true in this generality.

In this section, we recall the notions of prestacks and map-graded categories, thus fixing terminology and notations. As described explicitly in [12], prestacks and map-graded categories constitute two different incarnations of mathematical data that are equivalent in a suitable sense. A prestack is a pseudofunctor taking values in k -linear categories. We use the same terminology as in [14], [3].

Let \mathcal{U} be a small category.

Definition 2.1. A *prestack* $\mathcal{A} = (\mathcal{A}, m, f, c)$ on \mathcal{U} consists of the following data:

- for every object $U \in \mathcal{U}$, a k -linear category $(\mathcal{A}(U), m^U, 1^U)$ where m^U is the composition of morphisms in $\mathcal{A}(U)$ and 1^U encodes the identity morphisms on $\mathcal{A}(U)$;
- for every morphism $u: V \rightarrow U$ in \mathcal{U} , a k -linear functor $f^u = u^*: \mathcal{A}(U) \rightarrow \mathcal{A}(V)$. For $u = 1_U$, we require that $f^{1_U} = 1_U$.
- for every couple of morphisms $v: W \rightarrow V$, $u: V \rightarrow U$ in \mathcal{U} , a natural isomorphism

$$c^{u,v}: v^*u^* \rightarrow (uv)^*.$$

For $u = 1$ or $v = 1$, we require that $c^{u,v} = 1$. Moreover the natural isomorphisms have to satisfy the following coherence condition for every triple $w: T \rightarrow W$, $v: W \rightarrow V$, $u: V \rightarrow U$:

$$(2.1) \quad c^{u,vw}(c^{v,w} \circ u^*) = c^{uv,w}(w^* \circ c^{u,v}).$$

Remark 2.2. A presheaf of k -linear categories is considered as a prestack in which $c^{u,v} = 1$ for every $v: W \rightarrow V$, $u: V \rightarrow U$.

A prestack being a pseudofunctor, we obviously define a morphism of prestacks to be a pseudonatural transformation.

Definition 2.3. Consider prestacks (\mathcal{A}, m, f, c) and $(\mathcal{A}', m', f', c')$ on \mathcal{U} . A *morphism of prestacks* $(g, \tau): \mathcal{A} \rightarrow \mathcal{A}'$ consists of the following data:

- for each $U \in \mathcal{U}$, a functor $g^U: \mathcal{A}(U) \rightarrow \mathcal{A}'(U)$;
- for each $u: V \rightarrow U$ and $A \in \mathcal{A}(U)$, an element

$$\tau^{u,A} \in \mathcal{A}'(V)(u'^*g^U(A), g^V(u^*A))$$

These data further satisfy the following conditions: for any $v: W \rightarrow V$, $u: V \rightarrow U$ and $a \in \mathcal{A}(U)(A, B)$,

- (1) $m'^V(g^V u^*(a), \tau^u) = m'^V(\tau^u, u'^*g^U(a))$;
- (2) $m'^W(\tau^{uv}, c'^{u,v}) = m'^W(g^W(c^{u,v}), \tau^v, v'^*(\tau^u))$;
- (3) $m'^U(\tau^{1_U}, 1'_U) = g^U(1_U)$.

Let $\text{Mod}(k)$ be the category of k -modules and let $\underline{\text{Mod}}(k)$ be the constant prestack on \mathcal{U} with value $\text{Mod}(k)$. We are mainly interested in modules and bimodules.

Definition 2.4. Let \mathcal{A} be a prestack on \mathcal{U} . An \mathcal{A} -*module* is a morphism of prestacks $M: \mathcal{A}^{\text{op}} \rightarrow \underline{\text{Mod}}(k)$. More precisely, an \mathcal{A} -module consists of the following data:

- for every $U \in \mathcal{U}$, an $\mathcal{A}(U)$ -module $M^U: \mathcal{A}(U)^{\text{op}} \rightarrow \text{Mod}(k)$;
- for every $u: V \rightarrow U$, a morphism of $\mathcal{A}(U)$ -modules $M^u: M^U \rightarrow M^V u^*$; such that the following coherence condition holds for every $u: V \rightarrow U$, $v: W \rightarrow V$: the morphism M^{uv} equals the canonical composition

$$M^U \xrightarrow{M^u} M^V u^* \xrightarrow{M^v u^*} M^W v^* u^* \xrightarrow{M^W(c^{u,v})} M^W (uv)^* .$$

Definition 2.5. Let \mathcal{A}, \mathcal{B} be prestacks on \mathcal{U} . An \mathcal{A} - \mathcal{B} -*bimodule* is a module over $\mathcal{A}^{\text{op}} \otimes \mathcal{B}$. More concretely, an \mathcal{A} - \mathcal{B} -bimodule M consists of abelian groups

$$M^U(B, A)$$

for $U \in \text{Ob}(\mathcal{U})$, $A \in \text{Ob}(\mathcal{A}(U))$, $B \in \text{Ob}(\mathcal{B}(U))$, together with restriction morphisms

$$M^u(B, A): M^U(B, A) \rightarrow M^V(u^*B, u^*A)$$

for $u: V \rightarrow U$ in \mathcal{U} satisfying the natural coherence condition obtained from Definition 2.4.

Next we turn to map-graded categories in the sense of [12], where “map” stands for the maps in the underlying small category \mathcal{U} .

Definition 2.6. A \mathcal{U} -graded k -category $\mathfrak{a} = (\mathfrak{a}, \mu, \text{id})$ consists of the following data:

- for every object $U \in \mathcal{U}$, we have a set of *objects* $\mathfrak{a}(U)$;
- for every morphism $u : V \rightarrow U$ in \mathcal{U} and objects $A \in \mathfrak{a}(V), B \in \mathfrak{a}(U)$, we have a k -module $\mathfrak{a}_u(A, B)$ of *morphisms*.

These data are further equipped with compositions and identity morphisms in the following sense. The composition μ on \mathfrak{a} consists of operations

$$\mu^{u,v,A,B,C} : \mathfrak{a}_u(B, C) \otimes \mathfrak{a}_v(A, B) \longrightarrow \mathfrak{a}_{uv}(A, C)$$

satisfying the associativity condition

$$\mu^{w,uv,A,C,D}(\mu^{u,v,A,B,C} \otimes 1_{\mathfrak{a}_w(C,D)}) = \mu^{wu,v,A,B,D}(1_{\mathfrak{a}_v(A,B)} \otimes \mu^{w,u,B,C,D}).$$

The identity id on \mathfrak{a} consists of elements $\text{id}^A \in \mathfrak{a}_1(A, A)$ satisfying the condition

$$\mu^{u,1,A,A,B}(1_{\mathfrak{a}_u(A,B)} \otimes \text{id}^A) = 1_{\mathfrak{a}_u(A,B)} = \mu^{1,u,A,B,B}(\text{id}^B \otimes 1_{\mathfrak{a}_u(A,B)}).$$

The most natural type of modules to consider over a map-graded category turn out to be a kind of bimodules:

Definition 2.7. Let \mathfrak{a} be a \mathcal{U} -graded k -category. An \mathfrak{a} -bimodule M consists of k -modules

$$M_u(A, B)$$

for $u : V \rightarrow U, A \in \mathfrak{a}(V), B \in \mathfrak{a}(U)$ and compositions

$$\rho : \mathfrak{a}_u(C, D) \otimes M_v(B, C) \otimes \mathfrak{a}_w(A, B) \longrightarrow M_{uvw}(A, D)$$

satisfying the following associativity and identity conditions:

- (1) $\rho(\mu \otimes 1 \otimes \mu) = \rho(1 \otimes \rho \otimes 1)$;
- (2) $\rho(\text{id} \otimes 1 \otimes \text{id}) = 1$.

Let (\mathcal{A}, m, f, c) be prestack on \mathcal{U} . The associated \mathcal{U} -graded category $(\tilde{\mathcal{A}}, \mu, \text{id})$ is defined as a k -linear version of the Grothendieck construction from [1], more precisely:

- for each object $U \in \mathcal{U}$, we put $\tilde{\mathcal{A}}(U) = \text{Ob}(\mathcal{A}(U))$;
- for every morphism $u : V \rightarrow U$ and objects $A \in \tilde{\mathcal{A}}(V), B \in \tilde{\mathcal{A}}(U)$, we put

$$\tilde{\mathcal{A}}_u(A, B) = \mathcal{A}(U)(A, u^*B).$$

The composition operations

$$\mu : \tilde{\mathcal{A}}_u(B, C) \otimes \tilde{\mathcal{A}}_v(A, B) \longrightarrow \tilde{\mathcal{A}}_{uv}(A, C)$$

are defined by setting $\mu(b, a) = m(c^{u,v,C}, v^*b, a)$ for every $a \in \tilde{\mathcal{A}}_v(A, B), b \in \tilde{\mathcal{A}}_u(B, C)$ and the identities are given by $\text{id}^A = 1^{U,A} \in \mathcal{A}(U)(A, A) = \tilde{\mathcal{A}}_{1_U}(A, A)$ for $A \in \mathcal{A}(U)$.

There is a natural functor

$$\widetilde{(-)} : \text{Bimod}(\mathcal{A}) \longrightarrow \text{Bimod}(\tilde{\mathcal{A}}) : M \longmapsto \tilde{M}$$

defined by

$$\tilde{M}_u(A, B) := M^V(A, u^*B)$$

for every $u : V \rightarrow U, A \in \tilde{\mathcal{A}}(V), B \in \tilde{\mathcal{A}}(U)$. In [14], inspired by Gerstenhaber and Schack’s Cohomology Comparison Theorem [9], this functor is shown to induce a fully faithful functor on the level of the derived categories. In particular:

Theorem 2.8. [14, Theorem 1.1] *For any $M, N \in \text{Bimod}(\mathcal{A})$, we have*

$$\text{Ext}_{\mathcal{A}-\mathcal{A}}^n(M, N) \cong \text{Ext}_{\widetilde{\mathcal{A}}-\widetilde{\mathcal{A}}}^n(\widetilde{M}, \widetilde{N})$$

for all n .

3. THE GERSTENHABER-SCHACK COMPLEX FOR PRESTACKS

If \mathcal{A} is a presheaf of k -categories, then in analogy with the case of presheaves of k -algebras treated in [9, §21] and [8], one defines the Gerstenhaber-Schack (GS) complex $(\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M), d_{\text{GS}})$ for an \mathcal{A} -bimodule M as the total complex of a double complex with $d_{\text{GS}} = d_{\text{Hoch}} + d_{\text{simp}}$ for the horizontal Hochschild differential d_{Hoch} and the vertical simplicial differential d_{simp} . The cohomology of this complex is called Gerstenhaber-Schack (GS) cohomology and denoted

$$HH_{\text{GS}}^n(\mathcal{A}, M) = H^n \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M).$$

We denote $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}) = \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, \mathcal{A})$ and $HH_{\text{GS}}^n(\mathcal{A}) = H^n \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$.

In analogy with [3, §2] one sees that the second cohomology group $HH_{\text{GS}}^2(\mathcal{A})$ naturally classifies the first order deformations of \mathcal{A} as a *prestack*. Even though many prestacks of interest are in fact presheaves - for instance (restricted) structure sheaves of schemes as treated in [3] - the fact that prestacks turn up naturally as deformations suggests that it is really *prestacks* of which one should understand Gerstenhaber-Schack cohomology and deformations in the first place.

Our main aim in this section is to define a Gerstenhaber-Schack (GS) complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M)$ for an arbitrary prestack \mathcal{A} . Contrary to what one may at first expect, the change from the presheaf case to the prestack case is a major one. Indeed, if \mathcal{A} is non-trivially twisted ($c^{u,v} \neq 1$), with the natural definitions of d_{Hoch} and d_{simp} we now in general have $d_{\text{simp}}^2 \neq 0$ so we do not obtain a double complex. Instead, we construct a more complicated differential on the total double object $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M)$ by adding more components to the formula. After introducing the double object $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M)$ in §3.2 as a slight modification of the object associated to a presheaf, in §3.3 we introduce the infinite family of components $(d_j)_{j \geq 0}$ in (3.18), with

$$d_j : \mathbf{C}_{\text{GS}}^{p,q}(\mathcal{A}, M) \longrightarrow \mathbf{C}_{\text{GS}}^{p+j, q+1-j}(\mathcal{A}, M),$$

and we define the total differential

$$(3.1) \quad d = d_0 + d_1 + \cdots + d_n : \mathbf{C}_{\text{GS}}^{n-1}(\mathcal{A}, M) \longrightarrow \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M).$$

We have $d_0 = d_{\text{Hoch}}$ and $d_1 = (-1)^n d_{\text{simp}}$. The new differential d is shown to square to zero in Theorem 3.9. The definition of the higher components d_j is combinatorial in nature. It makes essential use of certain *paths* of natural transformations introduced in §3.3 and of the natural action of shuffle permutations on nerves of categories, as discussed in §3.1.

In order to properly relate the GS cohomology to deformation theory, we have to turn to the complex of normalized reduced cochains, which is introduced in §3.4 as a subcomplex of the GS complex and shown to be quasi-isomorphic to the latter in Propositions 3.13, 3.17. Finally, in §3.5 generalizing [3, Thm 2.21], in Theorem 3.20 we prove that HH_{GS}^2 classifies first order deformations of \mathcal{A} as a prestack.

3.1. Shuffle products. In this section, we discuss the natural action of shuffle permutations on nerves of categories. Let S_n be the symmetric group of permutations of $\{1, \dots, n\}$. For $n_i \geq 0$ with $\sum_{i=1}^k n_i = n$, a permutation $\beta \in S_n$ is an $(n_i)_i$ -*shuffle* if the following holds: for $1 \leq i \leq k$ and $\sum_{j=1}^{i-1} n_j + 1 \leq x \leq y \leq \sum_{j=1}^i n_j$

we have $\beta(x) \leq \beta(y)$. The permutation is a *conditioned* $(n_i)_i$ -*shuffle* if moreover we have

$$\beta\left(\sum_{i=1}^{l-1} n_i + 1\right) \leq \beta\left(\sum_{i=1}^l n_i + 1\right)$$

for all $1 \leq l \leq k-1$. Let $S_{(n_i)_i} \subseteq S_n$ be the subset of all $(n_i)_i$ -shuffles and $\bar{S}_{(n_i)_i} \subseteq S_{(n_i)_i}$ the subset of conditioned $(n_i)_i$ -shuffles. For any set X , S_n obviously has an action of X^n . For $\beta \in S_n$ and $(x_1, \dots, x_n) \in X^n$, we define

$$\beta^{(0)}(x_1, \dots, x_n) = (x_{\beta(1)}, \dots, x_{\beta(n)}).$$

When working with $(n_i)_i$ -shuffles, we will often consider different sets X_i for $1 \leq i \leq k$ and elements $x^i = (x_1^i, \dots, x_{n_i}^i) \in (X_i)^{n_i}$ for $1 \leq i \leq k$. Thus, for a permutation β , we obtain the *formal shuffle* by β of $(x^i)_i$:

$$(3.2) \quad \beta^{(0)}((x^i)_i) = \beta^{(0)}(x_1^1, \dots, x_{n_1}^1, \dots, x_1^k, \dots, x_{n_k}^k) \in \left(\prod_{i=1}^k X_i\right)^n.$$

For instance, for $k=2$, $\beta \in S_{m,n}$, $x = (x_1, \dots, x_m) \in X^m$ and $y = (y_1, \dots, y_n) \in Y^n$, we denote the formal shuffle by β of (x, y) by:

$$x *_{\beta}^{(0)} y = \beta^{(0)}(x, y) = \beta^{(0)}(x_1, \dots, x_m, y_1, \dots, y_n).$$

In the remainder of this section, we discuss the action of shuffle permutations on nerves of categories. Consider categories \mathcal{A}_i for $1 \leq i \leq k$. We now refine action (3.2) to obtain a *shuffle action*

$$(3.3) \quad S_{(n_i)_i} \times \prod_{i=1}^k \mathcal{N}_{n_i}(\mathcal{A}_i) \longrightarrow \mathcal{N}_n\left(\prod_{i=1}^k \mathcal{A}_i\right).$$

Consider $\beta \in S_{(n_i)_i}$ and

$$a^i = (A_0^i \xrightarrow{a_{n_i}^i} A_1^i \xrightarrow{a_{n_i-1}^i} \dots \xrightarrow{a_2^i} A_{n_i-1}^i \xrightarrow{a_1^i} A_{n_i}^i) \in \mathcal{N}_{n_i}(\mathcal{A}_i).$$

Note that it may occur that $n_i = 0$ and $a^i = A_0^i \in \mathcal{N}_0(\mathcal{A}_i) = \text{Ob}(\mathcal{A}_i)$. For the associated elements $\underline{a}^i = (a_1^i, a_2^i, \dots, a_{n_i-1}^i, a_{n_i}^i) \in \text{Mor}(\mathcal{A}_i)^{n_i}$, we obtain the formal shuffle $\underline{b} = \beta^{(0)}((\underline{a}^i)_i) = (\underline{b}_1, \dots, \underline{b}_n)$. We now inductively associate to \underline{b} an element

$$b = \beta((a^i)_i) \in \mathcal{N}_n\left(\prod_{i=1}^k \mathcal{A}_i\right)$$

with source $\prod_{i=1}^k A_0^i$ and target $\prod_{i=1}^k A_{n_i}^i$. Then b is called the *shuffle product* by β of $(a^i)_i$, and \underline{b} is called the *formal sequence* of b .

Since β is a shuffle permutation, we have $\underline{b}_1 = a_1^j : A_{n_j-1}^j \longrightarrow A_{n_j}^j$ for some $1 \leq j \leq k$. We put $B_n = \prod_{i=1}^k A_{n_i}^i$, $B_{n-1} = A_{n_1}^1 \times \dots \times A_{n_j-1}^j \times \dots \times A_{n_k}^k$ and

$$b_1 = (1_{A_{n_1}^1}, \dots, a_1^j, \dots, 1_{A_{n_k}^k}) : B_{n-1} \longrightarrow B_n.$$

Now suppose

$$\hat{b}_l = (B_{n-l} \xrightarrow{b_l} B_{n-l+1} \xrightarrow{b_{l-1}} \dots \xrightarrow{b_2} B_{n-1} \xrightarrow{b_1} B_n) \in \mathcal{N}_l\left(\prod_{i=1}^k \mathcal{A}_i\right)$$

has been defined with $B_{n-l} = \prod_{i=1}^k B_{n-l}^i$ and $B_{n-l}^i = A_{n_i-\alpha_i}^i$ where $\alpha_i = \max\{t \mid a_t^i \in \{\underline{b}_1, \dots, \underline{b}_l\}\}$. It then follows that $\underline{b}_{l+1} = a_{\alpha_j+1}^j$ for some $1 \leq j \leq k$ and we put $B_{n-l-1} = A_{n_1-\alpha_1}^1 \times \dots \times A_{n_j-\alpha_j-1}^j \times \dots \times A_{n_k-\alpha_k}^k$ and

$$(3.4) \quad b_{l+1} = (1_{A_{\alpha_1}^1}, \dots, a_{\alpha_j+1}^j, \dots, 1_{A_{\alpha_k}^k}) : B_{n-l-1} \longrightarrow B_{n-l}.$$

We thus arrive at the element

$$b = \beta((a^i)_i) = \hat{b}_n = (b_1, \dots, b_n) \in \mathcal{N}_n\left(\prod_{i=1}^k \mathcal{A}_i\right)$$

which concludes the definition of (3.3).

Remark 3.1. Suppose \mathcal{A} is a category and $\phi : \prod_{i=1}^k \mathcal{A}_i \longrightarrow \mathcal{A}$ is a functor. We naturally obtain an induced map $\mathcal{N}_n(\prod_{i=1}^k \mathcal{A}_i) \longrightarrow \mathcal{N}_n(\mathcal{A})$ which upon composition with (3.3) gives rise to a ϕ -shuffle action

$$(3.5) \quad S_{(n_i)_i} \times \prod_{i=1}^k \mathcal{N}_{n_i}(\mathcal{A}_i) \longrightarrow \mathcal{N}_n(\mathcal{A}) : (\beta, (a^i)_i) \longmapsto \beta^{(\phi)}((a^i)_i).$$

Obviously, taking $\phi = 1_{\prod_{i=1}^k \mathcal{A}_i}$, we recover the shuffle action (3.3). If ϕ is understood from the context, it will be omitted from the notation.

Example 3.2. Let \mathfrak{a} and \mathfrak{b} be small categories and put $\mathcal{A}_1 = \text{Fun}(\mathfrak{a}, \mathfrak{b})$, $\mathcal{A}_2 = \mathfrak{a}$, $\mathcal{A} = \mathfrak{b}$ and

$$\phi : \text{Fun}(\mathfrak{a}, \mathfrak{b}) \times \mathfrak{a} \longrightarrow \mathfrak{b} : (F, A) \longmapsto F(A).$$

Consider $a = (a_1 : A_1 \longrightarrow A_0) \in \mathcal{N}_1(\mathfrak{a})$ and

$$\epsilon = (T_0 \xrightarrow{\epsilon_2} T_1 \xrightarrow{\epsilon_1} T_2) \in \mathcal{N}_2(\text{Fun}(\mathfrak{a}, \mathfrak{b})).$$

The three elements in $S_{2,1}$ correspond to the following three formal shuffles of $\underline{\epsilon}$ and \underline{a} : $(a, \epsilon_1, \epsilon_2)$, $(\epsilon_1, a, \epsilon_2)$ and $(\epsilon_1, \epsilon_2, a)$. The three corresponding shuffles in $\mathcal{N}_3(\text{Fun}(\mathfrak{a}, \mathfrak{b}) \times \mathfrak{a})$ according to (3.3) are given by:

$$\begin{aligned} T_0 \times A_0 &\xrightarrow{\epsilon_2 \times 1_{A_0}} T_1 \times A_0 \xrightarrow{\epsilon_1 \times 1_{A_0}} T_2 \times A_0 \xrightarrow{1_{T_2} \times a} T_2 \times A_1; \\ T_0 \times A_0 &\xrightarrow{\epsilon_2 \times 1_{A_0}} T_1 \times A_0 \xrightarrow{1_{T_1} \times a} T_1 \times A_1 \xrightarrow{\epsilon_1 \times 1_{A_1}} T_2 \times A_1; \\ T_0 \times A_0 &\xrightarrow{1_{T_0} \times a} T_0 \times A_1 \xrightarrow{\epsilon_2 \times 1_{A_1}} T_1 \times A_1 \xrightarrow{\epsilon_1 \times 1_{A_1}} T_2 \times A_1. \end{aligned}$$

The three corresponding ϕ -shuffles in $\mathcal{N}_3(\mathfrak{b})$ according to (3.7) are given by:

$$\begin{aligned} T_0(A_0) &\xrightarrow{\epsilon_2(A_0)} T_1(A_0) \xrightarrow{\epsilon_1(A_0)} T_2(A_0) \xrightarrow{T_2(a)} T_2(A_1); \\ T_0(A_0) &\xrightarrow{\epsilon_2(A_0)} T_1(A_0) \xrightarrow{T_1(a)} T_1(A_1) \xrightarrow{\epsilon_1(A_1)} T_2(A_1); \\ T_0(A_0) &\xrightarrow{T_0(a)} T_0(A_1) \xrightarrow{\epsilon_2(A_1)} T_1(A_1) \xrightarrow{\epsilon_1(A_1)} T_2(A_1). \end{aligned}$$

Remark 3.3. Consider small categories $\mathfrak{b}_0, \dots, \mathfrak{b}_k$ and put $\mathcal{A}_i = \text{Fun}(\mathfrak{b}_{k-i}, \mathfrak{b}_{k-i+1})$. Applying the natural composition of functors to each element b_{l+1} in (3.4), we obtain

$$(3.6) \quad b'_{l+1} = A_{\alpha_1}^1 \circ \dots \circ a_{\alpha_j+1}^j \circ \dots \circ A_{\alpha_k}^k : B'_{n-l-1} \longrightarrow B'_{n-l}$$

where $B'_{n-l-1} = A_{n_1-\alpha_1}^1 \circ \dots \circ A_{n_j-\alpha_j-1}^j \circ \dots \circ A_{n_k-\alpha_k}^k$. Concatenating these morphisms, we obtain the simplex

$$\hat{b}'_n = (b'_1, \dots, b'_n) \in \mathcal{N}_n(\text{Fun}(\mathfrak{b}_0, \mathfrak{b}_k)).$$

Example 3.4. Consider

$$\epsilon = (T_0 \xrightarrow{\epsilon_2} T_1 \xrightarrow{\epsilon_1} T_2) \in \mathcal{N}_2(\text{Fun}(\mathfrak{b}_0, \mathfrak{b}_1))$$

and

$$\xi = (S_0 \xrightarrow{\xi} S_1) \in \mathcal{N}_1(\mathbf{Fun}(\mathbf{b}_1, \mathbf{b}_2)).$$

The shuffle products of ξ and ϵ with respect to composition of functors corresponding to the formal sequences $(\xi\epsilon_1, \epsilon_2), (\epsilon_1, \xi, \epsilon_2), (\epsilon_1, \epsilon_2, \xi)$ are

$$\begin{aligned} S_0T_0 &\xrightarrow{S_0\epsilon_2} S_0T_1 \xrightarrow{S_0\epsilon_1} S_0T_2 \xrightarrow{\xi T_2} S_1T_2; \\ S_0T_0 &\xrightarrow{S_0\epsilon_2} S_0T_1 \xrightarrow{\xi T_1} S_1T_1 \xrightarrow{S_1\epsilon_1} S_1T_2; \\ S_0T_0 &\xrightarrow{\xi T_0} S_1T_0 \xrightarrow{S_1\epsilon_2} S_1T_1 \xrightarrow{S_1\epsilon_1} S_1T_2. \end{aligned}$$

Now suppose $\beta \in \bar{S}_{(n_i)_i}$ is a conditioned shuffle. In this case it is possible to adapt the inductive procedure we just described in order to arrive at the datum, for $(a^i)_i$ as before, of a sequence

$$(3.7) \quad (\hat{c}_1, \dots, \hat{c}_k) \in \prod_{l=1}^k \mathcal{N}_{\gamma_l} \left(\prod_{i=1}^l \mathcal{A}_i \right)$$

where the numbers γ_l are determined by β and satisfy $\sum_{l=1}^k \gamma_l = n$. We put $\phi = 1$ and suppress it in the notations (the adaptation to arbitrary ϕ is easily made and will be used in the paper). Since β is a conditioned shuffle, there are uniquely determined numbers γ_l such that $\underline{b}_1 = a_1^1, \underline{b}_{\gamma_1+1} = a_1^2, \dots, \underline{b}_{\sum_{i=1}^l \gamma_i+1} = a_1^{l+1}, \dots, \underline{b}_{\sum_{i=1}^{k-1} \gamma_i+1} = a_1^k$ and $\gamma_k = n - \sum_{i=1}^{k-1} \gamma_i$. For $1 \leq l \leq k$ we now have that for every $\sum_{i=1}^{l-1} \gamma_i + 1 \leq \rho \leq \sum_{i=1}^l \gamma_i$ there exists $1 \leq j \leq l$ and t with $\underline{b}_\rho = a_t^j$. Moreover, for fixed j , there exists

$$a^{j,l} = (A_{n_j-t+1-m_j^l}^j \xrightarrow{a_{t+m_j^l-1}^j} \dots \xrightarrow{a_t^j} A_{n_j-t+1}^j) \in \mathcal{N}_{m_j^l}(\mathcal{A}_j)$$

such that the morphisms a_s^j occurring in $a^{j,l}$ coincide precisely with the elements occurring as \underline{b}_ρ for $\sum_{i=1}^{l-1} \gamma_i + 1 \leq \rho \leq \sum_{i=1}^l \gamma_i$. Here we make the convention that if no a_s^j occurs as such \underline{b}_ρ , we put $a^{j,l} \in \mathcal{N}_0(\mathcal{A}_j)$ equal to the domain of $a^{j,l-1}$, or equal to $a^{j,l-1}$ in case $a^{j,l-1} \in \mathcal{N}_0(\mathcal{A}_j)$. We have $\sum_{j=1}^l m_j^l = \gamma_l$. As a consequence, there is a unique $\beta_l \in S_{(m_j^l)_j}$ such that

$$\beta_l^{(0)}((a^{j,l})_j) = (\underline{b}_\rho)_{\sum_{i=1}^{l-1} \gamma_i+1 \leq \rho \leq \sum_{i=1}^l \gamma_i}.$$

In (3.7) we now put $\hat{c}_l = \beta_l((a^{j,l})_j) \in \mathcal{N}_{\gamma_l}(\prod_{i=1}^l \mathcal{A}_i)$.

3.2. The Gerstenhaber-Schack complex. Let \mathcal{U} be a small category, \mathcal{A} a prestack on \mathcal{U} and M a bimodule over \mathcal{A} . Let $\mathcal{N}(\mathcal{U})$ denote the simplicial nerve of the small category \mathcal{U} . Our standard notation for a p -simplex $\sigma \in \mathcal{N}(\mathcal{U})_p$ is

$$(3.8) \quad \sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \dots \xrightarrow{u_{p-1}} U_{p-1} \xrightarrow{u_p} U_p).$$

If confusion can arise, we write $U_i = U_i^\sigma$ and $u_i = u_i^\sigma$ instead. We also write $\sigma = (u_1, \dots, u_p)$ for short.

For $\sigma \in \mathcal{N}_p(\mathcal{U})$, we obtain a functor

$$\sigma^* = (u_p \dots u_2 u_1)^* : \mathcal{A}(U_p) \longrightarrow \mathcal{A}(U_0)$$

and a functor

$$\sigma^\sharp = u_1^* u_2^* \dots u_p^* : \mathcal{A}(U_p) \longrightarrow \mathcal{A}(U_0).$$

For each $1 \leq k \leq p-1$, denote by $L_k(\sigma)$ and $R_k(\sigma)$ the following simplices

$$\begin{aligned} L_k(\sigma) &= (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \dots \xrightarrow{u_{p-1}} U_{k-1} \xrightarrow{u_k} U_k) \\ R_k(\sigma) &= (U_k \xrightarrow{u_{k+1}} U_{k+2} \xrightarrow{u_{k+2}} \dots \xrightarrow{u_{p-1}} U_{p-1} \xrightarrow{u_p} U_p) \end{aligned}$$

We consider the following natural isomorphisms:

$$(3.9) \quad c^{\sigma,k} = c^{u_k \cdots u_1, u_p \cdots u_{k+1}} : (L_k \sigma)^* (R_k \sigma)^* \longrightarrow \sigma^*$$

$$(3.10) \quad \epsilon^{\sigma,k} = u_1^* \cdots u_{k-1}^* c^{u_k, u_{k+1}} u_{k+2}^* \cdots u_p^* : \sigma^\sharp \longrightarrow u_1^* \cdots (u_{k+1} u_k)^* \cdots u_p^*$$

For $A \in \text{Ob}(\mathcal{A}(U_p))$, we write $c^{\sigma,k,A} = c^{\sigma,k}(A)$ and $\epsilon^{\sigma,k,A} = \epsilon^{\sigma,k}(A)$.

For the category $\mathcal{A}(U)$, $U \in \mathcal{U}$, we use the following standard notation for a q -simplex $a \in \mathcal{N}(\mathcal{A}(U))_q$:

$$(3.11) \quad a = (A_0 \xrightarrow{a_q} A_1 \xrightarrow{a_{q-1}} \dots \xrightarrow{a_2} A_{q-1} \xrightarrow{a_1} A_q).$$

We also write $a = (a_1, \dots, a_q)$ for short.

Let

$$\mathbf{C}^{\sigma,A}(\mathcal{A}, M) = \text{Hom}_k(\mathcal{A}(U_p)(A_{q-1}, A_q) \otimes \cdots \otimes \mathcal{A}(U_p)(A_0, A_1), M^{U_0}(\sigma^\sharp A_0, \sigma^* A_q)).$$

and put

$$\begin{aligned} \mathbf{C}^{\sigma,q}(\mathcal{A}, M) &= \prod_{A \in \mathcal{A}(U_p)^{q+1}} \mathbf{C}^{\sigma,A}(\mathcal{A}, M), \\ \mathbf{C}^{p,q}(\mathcal{A}, M) &= \prod_{\sigma \in \mathcal{N}_p(\mathcal{U})} \mathbf{C}^{\sigma,q}(\mathcal{A}, M). \end{aligned}$$

Then we obtain the double object

$$(3.12) \quad \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M) = \prod_{p+q=n} \mathbf{C}^{p,q}(\mathcal{A}, M)$$

The usual Hochschild differential defines vertical maps

$$d_{\text{Hoch}} : \mathbf{C}^{p,q-1}(\mathcal{A}) \longrightarrow \mathbf{C}^{p,q}(\mathcal{A}).$$

Precisely, given $(\phi^\sigma)_\sigma \in \mathbf{C}^{p,q}(\mathcal{A}, M)$, for each p -simplex σ and for $(a_1, \dots, a_q) \in \mathcal{A}(U_p)(A_{q-1}, A_q) \otimes \cdots \otimes \mathcal{A}(U_p)(A_0, A_1)$, then we have

$$(d_{\text{Hoch}} \phi)^\sigma(a_1, \dots, a_q) = \sum_{i=0}^q (-1)^i (d_{\text{Hoch}}^i \phi)^\sigma(a_1, \dots, a_q)$$

where

$$(d_{\text{Hoch}}^i \phi)^\sigma(a_1, \dots, a_q) = \begin{cases} \sigma^*(a_1) \phi^\sigma(a_2, \dots, a_q) & \text{if } i = 0 \\ \phi^\sigma(a_1, \dots, a_i a_{i+1}, \dots, a_q) & \text{if } 1 \leq i \leq q-1 \\ \phi^\sigma(a_{q-1}, \dots, a_1) \sigma^\sharp(a_q) & \text{if } i = q. \end{cases}$$

We also write $\phi^\sigma(d_{\text{Hoch}}^i(a_q, \dots, a_1))$ instead of $(d_{\text{Hoch}}^i(\phi))^\sigma(a_q, \dots, a_1)$.

As a part of the simplicial structure of $\mathcal{N}(\mathcal{U})$, we have maps

$$\partial_i : \mathcal{N}_{p+1}(\mathcal{U}) \longrightarrow \mathcal{N}_p(\mathcal{U}) : \sigma \longmapsto \partial_i \sigma$$

for $i = 0, 1, \dots, p+1$. For $\sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \dots \xrightarrow{u_p} U_p \xrightarrow{u_{p+1}} U_{p+1})$, we have

$$\partial_{p+1} \sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \dots \xrightarrow{u_{p-1}} U_{p-1} \xrightarrow{u_p} U_p)$$

$$\partial_0 \sigma = (U_1 \xrightarrow{u_2} U_2 \xrightarrow{u_3} \dots \xrightarrow{u_p} U_p \xrightarrow{u_{p+1}} U_{p+1})$$

and

$$\partial_i \sigma = (U_0 \xrightarrow{u_1} \dots \xrightarrow{u_{i+1} u_i} U_{i+1} \xrightarrow{u_{p+1}} \dots \xrightarrow{u_{p+1}} U_{p+1})$$

for $i = 1, \dots, p$. Each ∂_i gives rise to a map

$$d_{\text{simp}}^i : \mathbf{C}^{p-1, q}(\mathcal{A}, M) \longrightarrow \mathbf{C}^{p, q}(\mathcal{A}, M)$$

given by

$$(d_{\text{simp}}^i(\phi))^\sigma(a_1, \dots, a_q) := \begin{cases} c^{\sigma, 1, A_q} M^{u_1} \phi^{\partial_0 \sigma}(a_1, \dots, a_q) & \text{if } i = 0 \\ \phi^{\partial_i \sigma}(a_1, \dots, a_q) \epsilon^{\sigma, i, A_0} & \text{if } 1 \leq i \leq p \\ c^{\sigma, p-1, A_q} \phi^{\partial_p \sigma}(u_p^* a_1, \dots, u_p^* a_q) & \text{if } i = p. \end{cases}$$

Hence we obtain the horizontal maps

$$d_{\text{simp}} = \sum_{i=0}^p (-1)^i d_{\text{simp}}^i : \mathbf{C}^{p-1, q}(\mathcal{A}, M) \longrightarrow \mathbf{C}^{p, q}(\mathcal{A}, M).$$

We define the maps

$$d_{\text{GS}} = d_{\text{Hoch}} + (-1)^n d_{\text{simp}} : \mathbf{C}^{n-1}(\mathcal{A}, M) \longrightarrow \mathbf{C}^n(\mathcal{A}, M).$$

Now if $c^{u, v} = 1$ for all $u : V \longrightarrow U, v : W \longrightarrow V$, then \mathcal{A} is a presheaf of k -linear categories. It is easy to check that $d_{\text{Hoch}}^2 = d_{\text{simp}}^2 = d_{\text{Hoch}} d_{\text{simp}} - d_{\text{simp}} d_{\text{Hoch}} = 0$, so $d_{\text{GS}}^2 = 0$. In analogy with [9], if k is a field one shows that $(\mathbf{C}^\bullet(\mathcal{A}, M), d_{\text{GS}})$ computes Ext groups of bimodules:

$$HH_{\text{GS}}^n(\mathcal{A}, M) = H^n(\mathbf{C}^\bullet(\mathcal{A}, M), d_{\text{GS}}) = \text{Ext}_{\mathcal{A}-\mathcal{A}}^n(\mathcal{A}, M).$$

Moreover, by analogous computations as in [3, §2.21], it is seen that the second cohomology group $HH_{\text{GS}}^2(\mathcal{A})$ naturally controls the first order deformations of the presheaf \mathcal{A} as a prestack.

3.3. The new differential. When \mathcal{A} is a prestack with non-trivial twists $c^{u, v}$, then for d_{GS} defined as in the previous section, we have $d_{\text{GS}}^2 \neq 0$ because $d_{\text{simp}}^2 \neq 0$. To fix this problem we add new components to d_{GS} to obtain the new differential

$$(3.13) \quad d = d_0 + d_1 + \dots + d_n : \mathbf{C}_{\text{GS}}^{n-1}(\mathcal{A}, M) \longrightarrow \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M)$$

where $d_0 = d_{\text{Hoch}}, d_1 = (-1)^n d_{\text{simp}}$ as above. The cohomology with respect to the new differential is denoted

$$HH_{\text{GS}}^n(\mathcal{A}, M) = H^n \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M).$$

Let \mathcal{A} be a prestack. Consider a simplex $\sigma = (u_1, \dots, u_n) \in \mathcal{N}_n(\mathcal{U})$ with $n \geq 2$. For every $u : V \longrightarrow U, v : W \longrightarrow V$ we have the natural isomorphism $c^{u, v} : v^* u^* \longrightarrow (uv)^*$. From these isomorphisms we inductively construct a set

$$(3.14) \quad \mathcal{P}(u_1, \dots, u_n) \subseteq \mathcal{N}_{n-1}(\text{Fun}(\mathcal{A}(U_n), \mathcal{A}(U_0)))$$

of simplices r with source $u_1^* u_2^* \dots u_n^*$ and target $(u_n u_{n-1} \dots u_1)^*$. Our standard notation for a simplex r of natural transformations is

$$r = (T_0 \xrightarrow{r_{n-1}} T_1 \xrightarrow{r_{n-2}} T_2 \longrightarrow \dots \xrightarrow{r_1} T_{n-1})$$

which is abbreviated to $r = (r_1, \dots, r_{n-1})$. Elements of $\mathcal{P}(u_1, \dots, u_n)$ are called *paths* from $u_1^* u_2^* \dots u_n^*$ to $(u_n u_{n-1} \dots u_1)^*$. Further, we define a *sign* map

$$\text{sign} : \mathcal{P}(u_1, \dots, u_n) \longrightarrow \{1, -1\} : r \longmapsto \text{sign}(r).$$

We start with $n = 2$. Consider $c^{u_1, u_2} : u_1^* u_2^* \longrightarrow (u_2 u_1)^*$. We put $\mathcal{P}(u_1, u_2) := \{(c^{u_1, u_2})\}$ and we set $\text{sign}(c^{u_1, u_2}) = -1$.

For $n > 2$, given $\sigma = (u_1, \dots, u_n)$, for each $i = 1, \dots, n-1$, consider the natural isomorphism $\epsilon^{\sigma, i} = u_1^* \cdots c^{u_i, u_{i+1}} \cdots u_n^*$ as defined in (3.10) and put

$$\text{sign}(\epsilon_i) = (-1)^i.$$

For each path $r = (r_1, \dots, r_{n-2}) \in \mathcal{P}(u_1, \dots, u_{i-1}, u_{i+1}u_i, u_{i+2}, \dots, u_n)$, the simplex $(r_1, \dots, r_{n-2}, \epsilon_i)$ is called a *path* from $u_1^*u_2^* \cdots u_n^*$ to $(u_n u_{n-1} \cdots u_1)^*$ and $\mathcal{P}(u_1, \dots, u_n)$ is defined to be the set of all such paths. Thus,

$$\mathcal{P}(u_1, \dots, u_n) = \{(r_1, \dots, r_{n-2}, \epsilon^{\sigma, i}) : 1 \leq i \leq n-1 \text{ and } r \in \mathcal{P}(\partial_i \sigma)\}.$$

For a path $r = (r_1, \dots, r_{n-1})$, we define

$$(-1)^r \equiv \text{sign}(r) = \prod_{i=1}^{n-1} \text{sign}(r_i).$$

For a permutation $\beta \in S_n$, we similarly denote $(-1)^\beta \equiv \text{sign}(\beta)$ for the standard sign of permutations and denote $(-1)^{r+\beta} = (-1)^r (-1)^\beta$.

There are $(n-1)!$ paths in $\mathcal{P}(u_1, \dots, u_n)$, for each path $r = (r_1, r_2, \dots, r_{n-1})$ denote the isomorphism $\|r\| = r_1 r_2 \cdots r_{n-1}$.

Example 3.5. Given $\sigma = (u_1, u_2, u_3)$, there are two paths from $u_1^*u_2^*u_3^*$ to $(u_3u_2u_1)^*$:

$$\mathcal{P} = \{r = (c^{u_2u_1, u_3}, c^{u_1, u_2}u_3^*), s = (c^{u_1, u_3}u_2, u_1^*c^{u_2, u_3})\}$$

and $\text{sign}(r) = 1$, $\text{sign}(s) = -1$.

The set of paths $\mathcal{P}(u_1, \dots, u_n)$ can be visualised in the following way. Let $[2] = \{0, 1\}$ be the poset with $0 < 1$, and consider the $(n-1)$ -dimensional ordered cube $[2]^{n-1}$. Every element $a = (a_i)_i \in [2]^{n-1}$ corresponds bijectively to a partition of u_1, \dots, u_n into a formal expression with parantheses

$$(3.15) \quad a(u_1, \dots, u_n) = (u_1, \dots, u_{i_1})(u_{i_1+1}, \dots, u_{i_2}) \cdots (u_{i_{k+1}}, \dots, u_n)$$

for $a_{i_1} = a_{i_2} = \cdots = a_{i_k} = 1$ and all other a_j equal to zero. Hence, we can define a function $F : [2]^{n-1} \rightarrow \text{Fun}(\mathcal{A}(U_n), \mathcal{A}(U_0))$ given by

$$F(a) = (u_1, \dots, u_{i_1})^*(u_{i_1+1}, \dots, u_{i_2})^* \cdots (u_{i_{k+1}}, \dots, u_n)^*$$

for $a(u_1, \dots, u_n)$ as in (3.15).

Example 3.6. For $\sigma = (u_1, u_2, u_3)$, the vertices $(0, 0), (1, 0), (0, 1)$ and $(1, 1)$ of the cube $[2]^2$ correspond to the functors $(u_3u_2u_1)^*, (u_1)^*(u_3u_2)^*, (u_2u_1)^*u_3^*$ and $u_1^*u_2^*u_3^*$ respectively.

For every two adjacent vertices in the cube $[2]^{n-1}$, there is a unique natural transformation between the corresponding functors under F that is induced from the twists $c^{u, v}$. Hence, we can visualise our paths as corresponding to composition series, or, equivalently, non-degenerate $(n-1)$ -simplices in the poset $[2]^{n-1}$.

The following lemma, which can alternatively be deduced from the universal property of cartesian liftings, shows that the function F on objects can actually be extended to a functor.

Lemma 3.7. *Assume given an n -simplex $\sigma = (u_1, \dots, u_n)$. Let $r = (r_1, r_2, \dots, r_{n-1})$ and $s = (s_1, s_2, \dots, s_{n-1})$ be two arbitrary paths in $\mathcal{P}(u_1, \dots, u_n)$. Then $\|r\| = \|s\|$.*

Proof. By the coherence condition (2.1) our lemma is true for $n = 3$. For $n > 3$, we assume that $r_{n-1} = \epsilon^{\sigma, i}$ and $s_{n-1} = \epsilon^{\sigma, j}$ for some $i \leq j$. If $i = j$ then $r_{n-1} = s_{n-1}$, by induction hypothesis we have $\|r\| = \|s\|$. If $i < j$, it is sufficient to prove that $\|r\| = \|t\|$ for some path $t = (t_1, \dots, t_{n-1})$ in which $t_{n-1} = \epsilon^{\sigma, i+1}$. Thus, let $h = (h_1, \dots, h_{n-2})$ be a path in $\mathcal{P}(u_1, \dots, u_{i+1}u_i, \dots, u_n)$ such that $h_{n-2} = u_1^* \cdots u_{i-1}^* c^{(u_{i+1}u_i, u_{i+2})} u_{i+3}^* \cdots u_n^*$, by the induction hypothesis

$$h_1 \cdots h_{n-2} = r_1 \cdots r_{n-2}.$$

Let $t_{n-2} = u_1^* \cdots u_{i-1}^* c^{(u_i, u_{i+2} u_{i+1})} u_{i+3}^* \cdots u_n^*$, again by (2.1) we have the commutative diagram

$$\begin{array}{ccc} u_1^* \cdots u_i^* u_{i+1}^* u_{i+2}^* \cdots u_n^* & \xrightarrow{t_{n-1}} & u_1^* \cdots u_i^* (u_{i+2} u_{i+1})^* u_{i+3}^* \cdots u_n^* \\ \downarrow r_{n-1} & & \downarrow t_{n-2} \\ u_1^* \cdots u_{i-1}^* (u_{i+1} u_i)^* u_{i+2}^* \cdots u_n^* & \xrightarrow{h_{n-2}} & u_1^* \cdots u_{i-1}^* (u_{i+2} u_{i+1} u_i)^* u_{i+3}^* \cdots u_n^* \end{array}$$

Choose $t = (h_1, \dots, h_{n-3}, t_{n-2}, t_{n-1})$, then $\|t\| = \|(h, r_{n-1})\| = \|r\|$. \square

Given a simplex $\sigma = (u_1, \dots, u_n)$, let $r = (r_1, \dots, r_{n-1})$ be a path in $\mathcal{P}(\sigma)$. For each $1 \leq k \leq n-2$, assume that $r_{k+1} = \epsilon^{\gamma, i}$ for some simplex $\gamma = (v_1, \dots, v_{k+2})$ and $1 \leq i \leq k+1$. Then $r_k = \epsilon^{\partial_i \gamma, j}$ for some $1 \leq j \leq k$. We put

$$\begin{cases} r'_{k+1} = \epsilon^{\gamma, j} \text{ and } r'_k = \epsilon^{\partial_j \gamma, i-1} \text{ if } i > j; \\ r'_{k+1} = \epsilon^{\gamma, j+1} \text{ and } r'_k = \epsilon^{\partial_j \gamma, i} \text{ if } i \leq j. \end{cases}$$

Denote by $\text{flip}(r, k)$ the path $(r_1, \dots, r_{k-1}, r'_k, r'_{k+1}, r_{k+2}, \dots, r_{n-1})$ in $\mathcal{P}(\sigma)$. It is easy to see that $\text{flip}(\text{flip}(r, k), k) = r$ and

$$(3.16) \quad \text{sign}(\text{flip}(r, k)) = -\text{sign}(r).$$

Due to Lemma 3.7, we have

$$(3.17) \quad r'_k r'_{k+1} = r_k r_{k+1}.$$

In the next lemma, which is easy to show, the shuffle product of natural transformations is taken with respect to the composition of functors as in Example (3.4).

Lemma 3.8. *Assume given an n -simplex $\sigma = (u_1, \dots, u_n)$. Then,*

- (1) *Consider two paths $r = (r_1, \dots, r_{n-k-1}) \in \mathcal{P}(R_k(\sigma))$, $s = (s_1, \dots, s_{k-1}) \in \mathcal{P}(L_k(\sigma))$. For each $\beta \in S_{n-k-1, k-1}$, the simplex $\omega = (c^{\sigma, k}, \beta(r, s))$ is a path in $\mathcal{P}(\sigma)$. Moreover*

$$(-1)^\omega = (-1)^{n-k} (-1)^\beta (-1)^r (-1)^s.$$

- (2) *Consider a path $\omega = (\omega_1, \dots, \omega_{n-1})$ in $\mathcal{P}(\sigma)$ in which $\omega_1 = c^{\sigma, k}$. There exist unique paths $r = (r_1, \dots, r_{n-k-1}) \in \mathcal{P}(R_k(\sigma))$, $s = (s_1, \dots, s_{k-1}) \in \mathcal{P}(L_k(\sigma))$ and $\beta \in S_{n-k-1, k-1}$ such that $\omega = (c^{\sigma, k}, \beta(r, s))$.*

Now we are able to define the components $d_j (j \geq 2)$ of the differential d from (3.13) in formula (3.18) below with $d_j : \mathbf{C}_{\text{GS}}^{p, q}(\mathcal{A}, M) \rightarrow \mathbf{C}_{\text{GS}}^{p+j, q+1-j}(\mathcal{A}, M)$.

$$\begin{array}{ccccc} & & & & \mathbf{C}^{p, q+1} \\ & & & & \uparrow \\ & & & & d_0 \\ & & & & \mathbf{C}^{p, q} \\ & \xrightarrow{d_1} & \mathbf{C}^{p+1, q} & & \\ & \searrow^{d_2} & & \searrow^{d_{q+1}} & \\ & & \mathbf{C}^{p+2, q-1} & & \\ & & & & \dots \\ & & & & \mathbf{C}^{p+q+1, 0} \end{array}$$

Consider $\phi \in \mathbf{C}_{\text{GS}}^{p,q}(\mathcal{A}, M)$. Let $\sigma = (u_1, \dots, u_{p+j})$ be a $(p+j)$ -simplex as in (3.8). Given $A_0, \dots, A_t \in \text{Ob}(\mathcal{A}(U_{p+j}))$ where $t = q+1-j$, let $a = (a_1, \dots, a_t)$ where $a_i \in \mathcal{A}(U_p)(A_{t-i}, A_{t-i+1})$ as in (3.11). We define

$$(3.18) \quad (d_j(\phi))^\sigma(a_1, \dots, a_t) = \sum_{\substack{r \in \mathcal{P}(R_j(\sigma)) \\ \beta \in S_{t,j-1}}} (-1)^r (-1)^\beta (-1)^t c^{\sigma,p,A_t} \phi^{L_p(\sigma)}(\beta(a, r))$$

where $\beta(a, r)$ is the shuffle product by β of $a = (a_1, \dots, a_t)$ and $r = (r_1, \dots, r_{j-1})$, with respect to the evaluation of functors (see Remark 3.1 and Example 3.2).

Theorem 3.9. $d \circ d = 0$.

Proof. For $N \geq 2$, for each cochain $\phi \in \mathbf{C}_{\text{GS}}^{p,q+N-2}(\mathcal{A}, M)$, we show the component of $d(d(\phi))$ which lies in $\mathbf{C}_{\text{GS}}^{p+N,q}(\mathcal{A}, M)$ is zero. Given a simplex $\sigma = (u_1, \dots, u_{p+N}) \in \mathcal{N}_{p+N}(\mathcal{U})$ and objects $A_0, A_1, \dots, A_q \in \mathcal{A}(U_{p+N})$. Let $a = (a_1, \dots, a_q)$ where $a_i \in \mathcal{A}(U_{p+N})(A_{q-i}, A_{q-i+1})$ as in (3.11). We need to show that

$$(d(d\phi))^\sigma(a) = \sum_{i=0}^N (d_{N-i}(d_i\phi))^\sigma(a) = 0.$$

This equation is equivalent to

$$(3.19) \quad (d_{\text{Hoch}} d_N \phi + d_{N-1} d_1 \phi + d_1 d_{N-1} \phi + \sum_{i=2}^{N-2} d_{N-i} d_i \phi)^\sigma(a) = -(d_N d_{\text{Hoch}} \phi)^\sigma(a).$$

By definition we have

$$-(d_N d_{\text{Hoch}} \phi)^\sigma(a) = \sum_{i=0}^{q+N-1} \sum_{r \in \mathcal{P}(R_p(\sigma)), \beta \in S_{q,N-1}} T(q, r, \beta, i)$$

where

$$T(a, r, \beta, i) = -(-1)^{q+i} (-1)^r (-1)^\beta c^{\sigma,p,A_q} (d_{\text{Hoch}}^i \phi)^{L_p(\sigma)}(\beta(a, r)).$$

We prove the equation (3.19) in the following steps:

- (1) For each term T_1 occurring in the expression of $d_{\text{Hoch}} d_N \phi$, there is a unique term $T(a, r, \beta, i)$ in $-(d_N d_{\text{Hoch}} \phi)$ such that $T_1 = T(a, r, \beta, i)$.
- (2) For $j = 2, \dots, (N-2)$, for each term T_2 occurring in $d_{N-j} d_j \phi$, there is a unique term $T(a, r, \beta', j')$ in $-(d_N d_{\text{Hoch}} \phi)$ such that $T_2 = T(a, r, \beta', j')$.
- (3) After cancellation, for each term T_3 in $d_{N-1} d_1 + d_1 d_{N-1}$, there is a unique term $T(a, r, \beta, i)$ in $-(d_N d_{\text{Hoch}} \phi)$ such that $T_3 = T(a, r, \beta, i)$.
- (4) After the cancellation with the terms T_1, T_2, T_3 as in step 1,2,3, denote X the remaining terms in $-(d_N d_{\text{Hoch}} \phi)$, then we show that $X = 0$.

Step 1. We have

$$d_{\text{Hoch}}(d_N \phi)^\sigma(a) = \sum_{i=0}^q \sum_{r \in \mathcal{P}(R_p(\sigma)), \beta \in S_{q-1,N-1}} T_1(d_{\text{Hoch}}^i(a), \beta, r, j)$$

where

$$T_1(d_{\text{Hoch}}^j(a), r, \beta, j) = (-1)^j (-1)^{q-1} (-1)^r (-1)^\beta c^{\sigma,p,A_q} \phi^{L_p(\sigma)}(\beta(d_{\text{Hoch}}^j(a), r)).$$

• Consider $j = 1, \dots, q-1$. For each path $r \in \mathcal{P}(R_p(\sigma))$, each $\beta \in S_{q-1,N-1}$, we write the formal sequence $\beta^{(0)}(d_{\text{Hoch}}^j(a), r) = (\beta_1, \dots, \beta_k, a_j a_{j+1}, \beta_{k+2} \dots \beta_{q+N-2})$ for some k . There is a unique shuffle permutation $\beta' \in S_{q,N-1}$ such that

$$\beta'^{(0)}(a, r) = (\underline{\beta}_1, \dots, \underline{\beta}_k, a_j, a_{j+1}, \underline{\beta}_{k+2}, \dots, \underline{\beta}_{q+N-2}).$$

Straightforward computations show that $T_1(d_{\text{Hoch}}^j(a), r, \beta, j) = T(a, r, \beta', k+1)$.

- Consider $j = 0$ or $j = q$. For $j = 0$, we have

$$T_1(d_{\text{Hoch}}^0(a), \beta, r, 0) = (-1)^{q-1}(-1)^r(-1)^\beta \sigma^*(a_1) c^{\sigma, p, A_{q-1}} \phi^{L_p}(\beta(a_2, \dots, a_q; r)).$$

Upon writing the formal sequence $\beta^{(0)}(a_2, \dots, a_q; r) = (\underline{\beta}_1, \dots, \underline{\beta}_{N+q-2})$, there is a unique $\beta' \in S_{q, N-1}$ such that $\beta^{(0)}(a, r) = (a_1, \underline{\beta}_1, \dots, \underline{\beta}_{q+N-2})$, and thus

$$T(a, r, \beta', 0) = T_1(d_{\text{Hoch}}^0(a), \beta, r, 0).$$

For $j = q$, we have

$$T_1(d_{\text{Hoch}}^q(a), \beta, r, 0) = -(-1)^r(-1)^\beta c^{\sigma, p, A_q} \phi^{L_p}(\beta(a_2, \dots, a_q; r)) \sigma^\#(a_q).$$

Assume that $\beta^{(0)}(a_1, \dots, a_{q-1}; r) = (\underline{\beta}_1, \dots, \underline{\beta}_{N+q-2})$, there is a unique $\beta' \in S_{q, N-1}$ such that $\beta^{(0)}(a, r) = (\underline{\beta}_1, \dots, \underline{\beta}_{q+N-2}, a_q)$, so we get that $T(a, r, \beta', q+N) = T_1(d_{\text{Hoch}}^q(a), \beta, r, q)$.

Step 2. We write

$$\sigma = (u_1, \dots, u_p, \dots, u_{p+N-j}, \dots, u_{p+N}).$$

Let $\Delta = (u_1, \dots, u_p, \dots, u_{p+N-j}) = L_{p+N-j}(\sigma)$. By definition, we have

$$(d_j(d_{N-j}\phi))^\sigma(a) = \sum_{\substack{r \in \mathcal{P}(R_{p+N-j}(\sigma)) \\ \beta \in S_{q, j-1}}} \sum_{\substack{s \in \mathcal{P}(R(\Delta, p)) \\ \gamma \in S_{q+j-1, N-j-1}}} T_2(a, r, \beta, s, \gamma)$$

where

$$T_2(a, r, \beta, s, \gamma) = (-1)^{j-1}(-1)^{r+s+\beta+\gamma} c^{\sigma, p+N-j, A_q} c^{\Delta, p, (R_{p+N-j}(\sigma))^* A_q} \phi^{L_p(\Delta)}(\gamma(\beta(a, r), s)).$$

The shuffle product is associative, hence $\gamma(\beta(a, r), s) = \beta(a, \gamma(r, s))$. Let $c_0 = c^{\sigma, p+1}$, by Lemma 3.8, we have $\omega = (c_0, \gamma(r, s))$ is a path in $\mathcal{P}(R_p(\sigma))$. There is a unique $\beta' \in S_{q, N-1}$ such that $\beta'(a, \omega) = (c_0(A_q), \beta(a, \gamma(r, s)))$. This implies that $T(a, \omega, \beta', 0) = T_2(a, r, \beta, s, \gamma)$.

Step 3. By definition we have

$$(d_{N-1}(d_1\phi))^\sigma(a) = \sum_{r \in \mathcal{P}(R_{p+1}(\sigma)), \beta \in S_{q, N-2}} (B(a, r, \beta) + \sum_{i=1}^p C(a, r, \beta, i) + D(a, r, \beta))$$

where

$$\begin{aligned} B(a, r, \beta) &= (-1)^{p+N-1}(-1)^{r+\beta} c^{\sigma, p+1, A_q} c^{L_{p+1}(\sigma), 1, (R_{p+1}(\sigma))^* A_q} M^{u_1}(\phi^{\partial_0(L_{p+1}(\sigma))}(\beta(a, r))); \\ C(a, r, \beta, i) &= (-1)^{p+N+i-1}(-1)^{r+\beta} c^{\sigma, p+1, A_q} \phi^{\partial_i L_{p+1}(\sigma)}(\beta(a, r)) \epsilon^{L_{p+1}(\sigma), i, (R_{p+1}(\sigma))^\# A_0}; \\ D(a, r, \beta) &= (-1)^N(-1)^{r+\beta} c^{\sigma, p+1, A_q} c^{L_{p+1}(\sigma), p, (R_{p+1}(\sigma))^* A_q} \phi^{\partial_{p+1} L_{p+1}(\sigma)}(u_{p+1}^*(\beta(a, r))). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} (d_1(d_{N-1}\phi))^\sigma(a) &= \sum_{\substack{r \in \mathcal{P}(R_{p+1}(\partial_0\sigma)) \\ \beta \in S_{q, N-2}}} B'(a, r, \beta) + \sum_{i=1}^p \sum_{\substack{r \in \mathcal{P}(R_{p+1}(\partial_i\sigma)) \\ \beta \in S_{q, N-2}}} C'(a, r, \beta, i) \\ &+ \sum_{i=p+1}^{p+N-1} \sum_{\substack{r \in \mathcal{P}(R_p(\partial_i\sigma)) \\ \beta \in S_{q, N-2}}} C''(a, r, \beta, i) + \sum_{\substack{r \in \mathcal{P}(R_p \partial_{p+N}\sigma) \\ \beta \in S_{q, N-2}}} D'(a, r, \beta) \end{aligned}$$

where

$$\begin{aligned} B'(a, r, \beta) &= (-1)^{p+N} (-1)^{r+\beta} c^{\sigma, 1, A_q} u_1^* (c^{\partial_0 \sigma, p+1, A_q}) M^{u_1} \left(\phi^{L_{p+1}(\partial_0 \sigma)}(\beta(a, r)) \right); \\ C'(a, r, \beta, i) &= (-1)^{p+N+i} (-1)^{r+\beta} c^{\partial_i \sigma, p+1, A_q} \phi^{L_{p+1}(\partial_i \sigma)}(\beta(a, r)) \epsilon^{\sigma, i, A_0}; \\ C''(a, r, \beta, i) &= (-1)^{p+N+i} (-1)^{r+\beta} c^{\partial_i \sigma, p, A_q} \phi^{L_p(\partial_i \sigma)}(\beta(a, r)) \epsilon^{\sigma, i, A_0}; \\ D'(a, r, \beta) &= (-1)^{r+\beta} c^{\sigma, p+N-1, A_q} c^{\partial_{p+N} \sigma, p, A_q} \phi^{L_p(\partial_{p+N} \sigma)}(\beta(u_{p+N}^*(a), r)). \end{aligned}$$

By computation, we get

$$\sum_{r \in \mathcal{P}(R_{p+1}(\partial_0 \sigma)), \beta \in S_{q, N-2}} B'(a, r, \beta) + \sum_{r \in \mathcal{P}(R_{p+1}(\sigma)), \beta \in S_{q, N-2}} B(a, r, \beta) = 0$$

and

$$\sum_{i=1}^p \sum_{r \in \mathcal{P}(R_{p+1}(\sigma)), \beta \in S_{q, N-2}} C(a, r, \beta, i) + \sum_{i=1}^p \sum_{r \in \mathcal{P}(R_{p+1}(\partial_i \sigma)), \beta \in S_{q, N-2}} C'(a, r, \beta, i) = 0.$$

So we obtain

$$\begin{aligned} (d_{N-1}(d_1 \phi))^\sigma(a) + (d_1(d_{N-1} \phi))^\sigma(a) &= \\ \sum_{\substack{r \in \mathcal{P}(R_{p+1}(\sigma)) \\ \beta \in S_{q, N-2}}} D(a, r, \beta) + \sum_{i=p+1}^{p+N-1} \sum_{\substack{r \in \mathcal{P}(R_p(\partial_i \sigma)) \\ \beta \in S_{q, N-2}}} C''(a, r, \beta, i) + \sum_{\substack{r \in \mathcal{P}(R_p \partial_{p+N} \sigma) \\ \beta \in S_{q, N-2}}} D'(a, r, \beta). \end{aligned}$$

We complete step 3 by showing that every term at the right hand side of this equation is matched with a unique term in $-(d_N d_{\text{Hoch}} \phi)^\sigma(a)$. First, consider the term $D(a, r, \beta)$ for $r = (r_1, \dots, r_{N-2}) \in \mathcal{P}(R_{p+1}(\sigma))$ and $\beta \in S_{q, N-2}$. Let $c_0 = c^{R_p(\sigma), 1}$, denote $u_{p+1}^* r = (u_{p+1}^* r_1, \dots, u_{p+1}^* r_{N-2})$ then $s = (c_0, u_{p+1}^* r)$ is a path in $\mathcal{P}(R_p \sigma)$ and there is a unique $\beta' \in S_{q, N-1}$ such that $\beta'(a, s) = (c_0(A_q), u_{p+1}^* \beta(a, r))$. This implies that $D(a, r, \beta) = T(a, s, \beta', 0)$.

Consider the term $C''(a, r, \beta, i)$ for $r \in \mathcal{P}(R_p \partial_i \sigma)$, $\beta \in S_{q, N-2}$ and $p+1 \leq i \leq p+N-1$. Then $s = (r, \epsilon^{\sigma, i})$ is a path in $\mathcal{P}(R_p \sigma)$, there is a unique $\beta' \in S_{q, N-1}$ such that $\beta'(a, s) = (\beta(a, r), \epsilon^{\sigma, i, A_0})$. Thus, we find that $C''(a, r, \beta, i) = T(a, s, \beta', q+N-1)$.

Consider the term $D'(a, r, \beta)$ where $r = (r_1, \dots, r_{N-2}) \in \mathcal{P}(R_p \partial_{p+N} \sigma)$ and $\beta \in S_{q, N-2}$. Let $c_0 = c^{R_p \sigma, p+N-1}$, denote $ru_{p+N}^* = (r_1 u_{p+N}^*, \dots, r_{N-2} u_{p+N}^*)$, then $s = (c_0, ru_{p+N}^*)$ is a path in $\mathcal{P}(R_p \sigma)$. There is a unique $\beta' \in S_{q, N-1}$ such that $\beta'(a, s) = (c_0(A_q), \beta(u_{p+N}^*(a), r))$. Hence, we obtain that $T(a, s, \beta', 0) = D'(a, r, \beta)$.

Step 4. For $\beta \in S_{q, N-1}$, $r \in \mathcal{P}(R_p \sigma)$, we write $\beta^{(0)}(a, r) = (\underline{\beta}_1, \dots, \underline{\beta}_{q+N-1})$. For each $k = 1, \dots, (q+N-2)$, denote by $S_{q, N-1}^k$ the set of all $(q, N-1)$ -shuffle permutations β such that $(\underline{\beta}_k, \underline{\beta}_{k+1}) \neq (a_i, a_{i+1})$, $\forall i = 1, \dots, q-1$. After steps 1, 2, 3, now it is seen that

$$-(d_N d_{\text{Hoch}} \phi)^\sigma(a) = (d_{\text{Hoch}} d_N \phi + d_{N-1} d_1 \phi + d_1 d_{N-1} \phi + \sum_{i=2}^{N-2} d_{N-i} d_i \phi)^\sigma(a) + X$$

where

$$X = \sum_{k=1}^{q+N-2} \sum_{r \in \mathcal{P}(R_p(\sigma)), \beta \in S_{q, N-1}^k} T(q, r, \beta, k).$$

Recall that

$$T(a, r, \beta, k) = (-1)^{q+1+k} (-1)^{r+\beta} c^{\sigma, p, A_q} (d_{\text{Hoch}}^k \phi)^{L_p(\sigma)}(\beta(a, r)).$$

Let $\beta \in S_{q, N-1}^k$, $r = (r_1, \dots, r_{N-1}) \in \mathcal{P}(R_p \sigma)$. In the expression

$$\beta^{(0)}(a, r) = (\underline{\beta}_1, \dots, \underline{\beta}_k, \underline{\beta}_{k+1}, \dots, \underline{\beta}_{q+N-1})$$

if $(\underline{\beta}_k, \underline{\beta}_{k+1}) = (a_i, r_j)$ or $(\underline{\beta}_k, \underline{\beta}_{k+1}) = (r_j, a_i)$ for some (i, j) , then take $\beta' = (k, k+1) \circ \beta$, then

$$T(a, r, \beta, k) + T(a, r, \beta', k) = 0.$$

Otherwise, $(\underline{\beta}_k, \underline{\beta}_{k+1}) = (r_i, r_{i+1})$ for some i . Then, by equations (3.17) and (3.16), we get

$$T(a, r, \beta, k) + T(a, \text{flip}(r, k), \beta, k) = 0.$$

Hence $X = 0$, this completes our proof. \square

3.4. Normalized reduced cochains. In this section, in analogy with [3, §2.4], we study the subcomplex $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M) \subseteq \mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ of normalized reduced cochains. Let $\sigma = (u_1, \dots, u_p)$ be a p -simplex as in (3.8). The simplex σ is said to be *right k -degenerate* if $u_i = 1_{U_i}$ for some $p-k+1 \leq i \leq p$ and σ is said to be *degenerate* if it is right k -degenerate for $k = p$. For $A = (A_1, \dots, A_q) \in \mathcal{A}(U_p)$ and $a = (a_1, \dots, a_q)$ as in (3.11), a is said to be *normal* if $a_i = 1$ for some i .

Given a cochain $\phi = (\phi^\sigma)_\sigma \in \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M)$, ϕ^σ is said to be *normalized* if $\phi^\sigma(a) = 0$ as soon as a is normal, and ϕ is said to be *normalized* if ϕ^σ is normalized for every simplex σ . The normalized cochains form a subcomplex $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ of $\mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$. The cochain ϕ is said to be *right k -reduced* if $\phi^\sigma = 0$ for every right k -degenerate simplex σ and ϕ is said to be *reduced* if $\phi^\sigma = 0$ for every degenerate simplex σ . The normalized reduced cochains further form a subcomplex $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ of $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$.

Inspired by [3, §2.4], [8, §7], we first prove that the inclusion $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M) \hookrightarrow \mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ is a quasi-isomorphism. It is more subtle to prove that $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M) \hookrightarrow \bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ is also a quasi-isomorphism. Due to the higher components of our new differential, the spectral sequence argument does not apply as in [3, §2.4]. As a single filtration is not sufficient, we use a double filtration instead.

Remark 3.10. If \mathcal{A} is a presheaf of k -linear categories, then the new differential d on $\mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ does not reduce to d_{GS} from §3.2. However, on the quasi-isomorphic subcomplex $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M) \subseteq \mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ of normalized reduced cochains, d and d_{GS} do coincide in this case.

Lemma 3.11. *Consider a cochain complex (D^{\bullet}, δ) with a subcomplex $(D'^{\bullet}, \delta) \subseteq (D^{\bullet}, \delta)$. Assume that for all n , for every cochain $f \in D^n$ with $\delta(f) \in D'^{n+1}$, there exists $h \in D'^{n-1}$ such that $f - \delta(h) \in D'^n$. Then the inclusion $(D'^{\bullet}, \delta) \hookrightarrow (D^{\bullet}, \delta)$ is a quasi-isomorphism.*

Proof. The condition spelled out is readily seen to be equivalent to the quotient complex D^{\bullet}/D'^{\bullet} being acyclic. \square

It is seen that for each simplex σ , $\mathbf{C}_{\text{GS}}^{\sigma, \bullet}(\mathcal{A}, M)$ is a cochain complex with the differential d_{Hoch} . By similar computations as in [8, §7] we obtain

Lemma 3.12. *Let $f \in \mathbf{C}_{\text{GS}}^{\sigma, n}(\mathcal{A}, M)$ be a cochain. If $d_{\text{Hoch}}(f)$ is normalized, then there exists $h \in \mathbf{C}_{\text{GS}}^{\sigma, n-1}(\mathcal{A}, M)$ such that $f - d_{\text{Hoch}}(h)$ is normalized.*

Equip $\mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ with a filtration

$$\dots \subseteq F^p \mathbf{C}^n \subseteq F^{p-1} \mathbf{C}^n \subseteq \dots \subseteq F^0 \mathbf{C}^n \subseteq F^{-1} \mathbf{C}^n = \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M)$$

by setting

$$F^j \mathbf{C}^n = \{ \phi = (\phi^\sigma)_\sigma \in \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M) \mid \phi^\sigma \text{ is normalized if } |\sigma| \leq j \}.$$

Since $d(F^j \mathbf{C}^p) \subseteq F^j \mathbf{C}^{p+1}$, $F^j \mathbf{C}^{\bullet}$ is a complex. There is a sequence of complexes

$$(3.20) \quad \dots \hookrightarrow F^j \mathbf{C}^{\bullet} \hookrightarrow F^{j-1} \mathbf{C}^{\bullet} \hookrightarrow \dots \hookrightarrow F^0 \mathbf{C}^{\bullet}.$$

Proposition 3.13. *The following inclusions are quasi-isomorphisms:*

- (1) $l: F^j \mathbf{C}^{\bullet} \hookrightarrow F^{j-1} \mathbf{C}^{\bullet}$;

$$(2) \quad \bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M) \hookrightarrow \mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M).$$

Proof. It suffices to prove that (1) is a quasi-isomorphism. By Lemma 3.11 it is sufficient to prove that for every cochain $\phi \in F^{j-1}\mathbf{C}^n$, if $d(\phi) \in F^j\mathbf{C}^{n+1}$ then there exists a cochain $\psi \in F^{j-1}\mathbf{C}^{n-1}$ such that $\phi - d(\psi) \in F^j\mathbf{C}^n$. Writing $\phi = (\phi_{p,q})_{p+q=n}$, we assume that $d(\phi) \in F^j\mathbf{C}^{n+1}$. Let σ be a j -simplex and let $a = (a_1, \dots, a_{n+1-j})$ be normal, then $(d(\phi))^\sigma(a) = 0$. By definition, we have

$$(d\phi)^\sigma(a) = \sum_{i=0}^j (d_i \phi_{j-i, n-j+i})^\sigma(a).$$

Note that $(d_i \phi_{j-i, n-j+i})^\sigma(a) = 0$ for $i > 0$ as $\phi \in F^{j-1}\mathbf{C}^n$. Hence we get

$$(d_{\text{Hoch}} \phi_{j, n-j})^\sigma(a) = 0.$$

By Lemma 3.12, there exists $h^\sigma \in \mathbf{C}^{\sigma, n-j-1}$ such that $\phi_{j, n-j}^\sigma - d_{\text{Hoch}}(h^\sigma)$ is normalized. We define $\psi^\sigma = h^\sigma$ if $|\sigma| = j$ and $\psi^\sigma = 0$ otherwise. Thus $\psi \in F^{j-1}\mathbf{C}^{n-1}$ and it is easy to see that $\phi - d(\psi) \in F^j\mathbf{C}^n$. \square

Now equip $\bar{\mathbf{C}}_{\text{GS}}^{\bullet}(\mathcal{A}, M)$ with a filtration

$$\dots \subseteq F^{lp}\bar{\mathbf{C}}^n \subseteq F^{l(p-1)}\bar{\mathbf{C}}^n \subseteq \dots \subseteq F^{l0}\bar{\mathbf{C}}^n = \bar{\mathbf{C}}_{\text{GS}}^n(\mathcal{A}, M)$$

by setting, for each $k \geq 1$,

$$F'^k \bar{\mathbf{C}}^n = \{\phi = (\phi^\sigma)_\sigma \in \bar{\mathbf{C}}_{\text{GS}}^n(\mathcal{A}, M) \mid \phi^\sigma(a) = 0 \forall a, \text{ if } \sigma \text{ is right } k\text{-degenerate}\}.$$

By straightforward computations, we obtain the following lemma.

Lemma 3.14. $d(F'^k \bar{\mathbf{C}}^n) \subseteq F'^k \bar{\mathbf{C}}^{n+1}$.

By Lemma 3.14 we obtain a sequence of complexes

$$(3.21) \quad \dots \hookrightarrow F'^k \bar{\mathbf{C}}^{\bullet} \hookrightarrow F'^{k-1} \bar{\mathbf{C}}^{\bullet} \hookrightarrow \dots \hookrightarrow F'^0 \bar{\mathbf{C}}^{\bullet}.$$

Next, for each $k \geq 0$, we equip $F'^k \bar{\mathbf{C}}$ with a further filtration

$$F'^{k+1} \bar{\mathbf{C}}^n = G^{n+1} F'^k \bar{\mathbf{C}}^n \subseteq \bar{\mathbf{C}}^n \dots \subseteq G^{l+1} F'^k \bar{\mathbf{C}}^n \subseteq G^l F'^k \bar{\mathbf{C}}^n \subseteq \dots \subseteq G^0 F'^k \bar{\mathbf{C}}^n = F'^k \bar{\mathbf{C}}^n$$

by setting

$$G^l F'^k \bar{\mathbf{C}}^n = \{\phi \in F'^k \bar{\mathbf{C}}^n \mid \phi^\sigma = 0 \text{ for } |\sigma| \geq n - l + 1 \text{ and } \sigma \text{ is right } (k+1)\text{-degenerate}\}.$$

By analogous computations as in Lemma 3.14, we get

$$d(G^l F'^k \bar{\mathbf{C}}^n) \subseteq G^l F'^k \bar{\mathbf{C}}^{n+1}.$$

Thus, for each k , we obtain a sequence of complexes

$$(3.22) \quad \dots \hookrightarrow G^{l+1} F'^k \bar{\mathbf{C}}^{\bullet} \hookrightarrow G^l F'^k \bar{\mathbf{C}}^{\bullet} \hookrightarrow \dots \hookrightarrow F'^k \bar{\mathbf{C}}^{\bullet}.$$

Lemma 3.15. *Let ϕ be a right k -reduced cochain in $\bar{\mathbf{C}}_{\text{GS}}^{p,q}(\mathcal{A}, M)$. If $d_{\text{simp}}\phi$ is a right $(k+1)$ -reduced cochain in $\bar{\mathbf{C}}_{\text{GS}}^{p+1,q}(\mathcal{A}, M)$, then there exists a right k -reduced cochain $\psi \in \bar{\mathbf{C}}_{\text{GS}}^{p-1,q}(\mathcal{A}, M)$ such that $\phi - d_{\text{simp}}\psi$ is a right $(k+1)$ -reduced cochain.*

Proposition 3.16. *The following inclusions are quasi-isomorphism:*

- (1) $G^{l+1} F'^k \bar{\mathbf{C}}^{\bullet} \hookrightarrow G^l F'^k \bar{\mathbf{C}}^{\bullet}$;
- (2) $F'^{k+1} \bar{\mathbf{C}}^{\bullet} \hookrightarrow F'^k \bar{\mathbf{C}}^{\bullet}$;
- (3) $\bar{\mathbf{C}}'_{\text{GS}}(\mathcal{A}, M) \hookrightarrow \bar{\mathbf{C}}_{\text{GS}}(\mathcal{A}, M)$.

Proof. For each n , the filtrations (3.21) and (3.22) are stationary, so we only need to prove (1). By Lemma 3.11, it is sufficient to prove that for any cochain $\phi = (\phi_{p,q}) \in G^l F'^k \bar{\mathbf{C}}^n$ which satisfies $d\phi \in G^{l+1} F'^k \bar{\mathbf{C}}^{n+1}$, there exists a cochain $\psi \in G^l F'^k \bar{\mathbf{C}}^{n-1}$ such that $\phi - d\psi \in G^{l+1} F'^k \bar{\mathbf{C}}^n$.

Set $p = n - l + 1$ and let σ be $(k + 1)$ -right degenerate p -simplex. By definition, we have $\phi_{p,n-p}^\sigma = 0$. Assume that $(d\phi)^\sigma = 0$. This implies $(d_{\text{simp}}\phi_{p-1,n-p+1})^\sigma = 0$. Apply Lemma 3.15, there exists $h \in F'^k \bar{\mathbf{C}}^{p-2,n-p+1}$ such that

$$(\phi_{p-1,n-p+1} - d_1(h))^{\sigma'} = 0$$

for every $(k + 1)$ -right degenerate $(p - 1)$ -simplex σ' .

We define $\psi^\sigma = h^\sigma$ if $|\sigma| = p - 2$ and $\psi^\sigma = 0$ elsewhere. It is seen that $\psi \in G^l F'^k \bar{\mathbf{C}}^{n-1}$ and $\phi - d\psi \in G^{l+1} F'^k \bar{\mathbf{C}}^n$ as desired. \square

Combining Propositions 3.13 and 3.16, we now obtain the following isomorphisms.

Proposition 3.17. *Let M be an \mathcal{A} -bimodule. Then*

$$H^n \bar{\mathbf{C}}'_{\text{GS}}(\mathcal{A}, M) \simeq H^n \bar{\mathbf{C}}_{\text{GS}}(\mathcal{A}, M) \simeq H^n \mathbf{C}_{\text{GS}}(\mathcal{A}, M).$$

Remark 3.18. If \mathcal{A} is a presheaf of k -linear categories, then the new differential d does not reduce to the old d from §3.2. However, on the quasi-isomorphic subcomplex $\bar{\mathbf{C}}'_{\text{GS}}(\mathcal{A}, M) \subseteq \mathbf{C}_{\text{GS}}(\mathcal{A}, M)$ of normalized reduced cochains defined in §3.4, they do coincide in this case.

3.5. First-order deformations of prestacks. In this section, generalizing [3, Thm 2.21], we prove that HH_{GS}^2 classifies first order deformations of prestacks.

Definition 3.19. (see Def 3.24 in [12]) Let (\mathcal{A}, m, f, c) be a prestack over \mathcal{U} .

- (1) A *first order deformation* of \mathcal{A} is given by a prestack

$$(\bar{\mathcal{A}}, \bar{m}, \bar{f}, \bar{c}) = (\mathcal{A}[\epsilon], m + m_1\epsilon, f + f_1\epsilon, c + c_1\epsilon)$$

of $k[\epsilon]$ -categories where $(m_1, f_1, c_1) \in \mathbf{C}^{0,2}(\mathcal{A}) \oplus \mathbf{C}^{1,1}(\mathcal{A}) \oplus \mathbf{C}^{2,0}(\mathcal{A})$.

- (2) For two deformations $(\bar{\mathcal{A}}, \bar{m}, \bar{f}, \bar{c})$ and $(\bar{\mathcal{A}}', \bar{m}', \bar{f}', \bar{c}')$ an *equivalence of deformations* is given by an isomorphism of the form $(g, \tau) = (1 + g_1\epsilon, 1 + \tau_1\epsilon)$ where $(g_1, \tau_1) \in \mathbf{C}^{0,1}(\mathcal{A}) \oplus \mathbf{C}^{1,0}(\mathcal{A})$.

Theorem 3.20. *Let $\mathcal{A} = (\mathcal{A}, m, f, c)$ be a prestack with GS complex $(\mathbf{C}_{\text{GS}}(\mathcal{A}), d)$. Then the second cohomology $HH_{\text{GS}}^2(\mathcal{A})$ classifies the first order deformations of \mathcal{A} . More precisely:*

- (1) For (m_1, f_1, c_1) in $\mathbf{C}^{0,2}(\mathcal{A}) \oplus \mathbf{C}^{1,1}(\mathcal{A}) \oplus \mathbf{C}^{2,0}(\mathcal{A})$, we have that $(\mathcal{A}[\epsilon], \bar{m} = m + m_1\epsilon, \bar{f} = f + f_1\epsilon, \bar{c} = c + c_1\epsilon)$ is a first order deformation of \mathcal{A} if and only if $(m_1, f_1, c_1) \in \mathbf{C}'_{\text{GS}}(\mathcal{A})$ and $d(m_1, f_1, c_1) = 0$.
- (2) For (m_1, f_1, c_1) and (m'_1, f'_1, c'_1) in $Z^2 \bar{\mathbf{C}}'_{\text{GS}}(\mathcal{A})$, and $(g_1, -\tau_1) \in \mathbf{C}^{0,1}(\mathcal{A}) \oplus \mathbf{C}^{1,0}(\mathcal{A})$, we have that $(g, \tau) = (1 + g_1\epsilon, 1 + \tau_1\epsilon)$ is an isomorphism between the corresponding deformed prestacks $\bar{\mathcal{A}}$ and $\bar{\mathcal{A}}'$ if and only if $(g_1, -\tau_1) \in \bar{\mathbf{C}}^1_{\text{GS}}(\mathcal{A})$ and $d(g_1, -\tau_1) = (m_1, f_1, c_1) - (m'_1, f'_1, c'_1)$. We have an isomorphism of sets

$$(3.23) \quad H^2 \bar{\mathbf{C}}'_{\text{GS}}(\mathcal{A}) \longrightarrow \text{Def}(\mathcal{A}).$$

Hence, the second cohomology group $HH^2(\mathcal{A})_{\text{GS}} \cong H^2 \bar{\mathbf{C}}'_{\text{GS}}(\mathcal{A})$ classifies first order deformations of \mathcal{A} up to equivalence.

Proof. (1) For each $U \in \mathcal{U}$, the composition \bar{m}^U of $\mathcal{A}(U)$ is associative if and only if

$$(d_{\text{Hoch}} m_1)^U = 0.$$

For each $a \in \mathcal{A}(U)(A, B)$, the unity condition $\bar{m}^U(1_B, a) = a = \bar{m}^U(a, 1_A)$ holds if and only if $m_1^U(1_B, a) = m_1^U(a, 1_A) = 0$.

For each 1-simplex $\sigma = (V \xrightarrow{v} U)$ and $(a, b) \in \mathcal{A}(U)(A, B) \times \mathcal{A}(U)(B, C)$. The condition $\bar{m}^V(\bar{f}(b), \bar{f}(a)) = \bar{f}(\bar{m}^U(b, a))$ holds if and only if

$$(d_{\text{Hoch}}f_1)^\sigma(b, a) - (d_{\text{simp}}m_1)^\sigma(b, a) = 0.$$

The condition $\bar{f}^\sigma(1_A) = 1_{\bar{f}^\sigma(A)}$ is equivalent to $f_1^\sigma(1_A) = 0$. The condition $\bar{f}^{1_U} = 1_U$ holds if and only if $f_1^{1_U} = 0$.

For each 2-simplex $\sigma = (W \xrightarrow{v} V \xrightarrow{u} U)$ and $a \in \mathcal{A}(U)(A, B)$, the condition $\bar{m}(\bar{c}^{u,v,B}, \bar{f}^v \bar{f}^u(a)) = \bar{m}(\bar{f}^{uv}(a), \bar{c}^{u,v,A})$ holds if and only if

$$(d_{\text{Hoch}}c_1)^\sigma(a) - (d_{\text{simp}}f_1)^\sigma(a) + (d_2m_1)^\sigma(a) = 0.$$

The condition that $\bar{c}^\sigma = 1$ when σ is degenerated holds if and only if $c_1^\sigma = 0$ if σ is degenerated.

For each 3-simplex $\sigma = (T \xrightarrow{w} W \xrightarrow{v} V \xrightarrow{u} U)$, the compatibility of \bar{c} holds if and only if

$$-(d_{\text{simp}}c_1)^\sigma(A) + (d_2f_1)^\sigma(A) + (d_3m_1)^\sigma(A) = 0.$$

Recall that

$$d(m_1, f_1, c_1) = (d_{\text{Hoch}}m_1, d_{\text{Hoch}}f_1 - d_{\text{simp}}m_1, d_{\text{Hoch}}c_1 - d_{\text{simp}}f_1 + d_2m_1, \\ - d_{\text{simp}}c_1 + d_2f_1 + d_2m_1).$$

These facts yield that (m_1, f_1, c_1) gives rise to a deformation of the prestack \mathcal{A} if and only if it is a normalized reduced cocycle.

(2) For each $U \in \mathcal{U}$, we have that g^U is a functor if and only if $g_1^U(1) = 0$ and

$$d_{\text{Hoch}}(g_1) = m_1 - m'_1.$$

For each 1-simplex $\sigma = (V \xrightarrow{u} U)$ and $a \in \mathcal{A}(U)(A, B)$, the condition $m'^V(g^V u^*(a), \tau^u) = m'^V(\tau^u, u'^* g^U(a))$ holds if and only if

$$(d_{\text{Hoch}}g_1)^\sigma(a) + (d_{\text{simp}}(-\tau_1))^\sigma(a) = f_1^\sigma(a) - f_1'^\sigma(a).$$

The condition $m'^U(\tau^{1_U}, 1'_U) = g^U(1_U)$ holds if and only if $\tau_1^{1_U} = 0$.

For each 2-simplex $\sigma = (W \xrightarrow{v} V \xrightarrow{u} U)$ and $A \in \mathcal{A}(U)$, the condition $m'^W(\tau^{uv}, c'^{u,v}) = m'^W(g^W(c^{u,v}), \tau^v, v'^*(\tau^u))$ holds if and only if

$$(d_{\text{simp}}(-\tau_1))^\sigma(A) + (d_2g_1)^\sigma(A) = c_1^\sigma(A) - c_1'^\sigma(A).$$

Hence $(g, \tau) = (1 + g_1\epsilon, 1 + \tau_1\epsilon)$ is an isomorphism between \mathcal{A} and \mathcal{A}' if and only if $(g_1, -\tau_1)$ is a normalized reduced cochain and

$$d(g_1, -\tau_1) = (d_{\text{Hoch}}g_1, d_{\text{Hoch}}(-\tau_1) + d_{\text{simp}}g_1, d_{\text{simp}}(-\tau_1) + d_2g_1) \\ = (m_1, f_1, c_1) - (m'_1, f'_1, c'_1).$$

□

4. COMPARISON OF COMPLEXES

Let \mathcal{U} be a small category, \mathcal{A} a prestack on \mathcal{U} , and M an \mathcal{A} -bimodule. In this section, we define cochain maps

$$\mathcal{F} : \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M) \longrightarrow \mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M}) \quad \text{and} \quad \mathcal{G} : \mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M}) \longrightarrow \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M)$$

between the GS complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M)$ and the Hochschild complex $\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M})$ as defined in [12]. We prove that \mathcal{F} and \mathcal{G} are inverse quasi-isomorphisms. In combination with [12, Prop. 3.13] and the Cohomology Comparison Theorem [14, Thm. 1.1] it follows that - as in the case of presheaves - if k is a field then the cohomology

of the complex $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M)$ computes bimodule Ext groups. More precisely, in this case we obtain

$$(4.1) \quad HH_{\text{GS}}^n(\mathcal{A}, M) \cong H^n(\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M})) \cong \text{Ext}_{\tilde{\mathcal{A}}-\tilde{\mathcal{A}}}^n(\tilde{\mathcal{A}}, \tilde{M}) \cong \text{Ext}_{\tilde{\mathcal{A}}-\mathcal{A}}^n(\mathcal{A}, M).$$

Due to the presence of the infinite family of components of the differential on $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$, the definition of \mathcal{F} and \mathcal{G} requires new combinatorial constructions. A key element is the use of partitions combined with appropriate shuffle products. Consider an n -simplex $\sigma = (u_1, \dots, u_n)$, objects $A_i \in \text{Ob}(\tilde{\mathcal{A}}(U_i))$, and morphisms $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n)$ where

$$\tilde{a}_i \in \tilde{\mathcal{A}}_{u_{n+1-i}}(A_{n-i}, A_{n+1-i}) = \mathcal{A}(U_{n-i})(A_{n-i}, u_{n+1-i}^* A_{n+1-i})$$

as follows:

$$A_0 \xrightarrow{\tilde{a}_n} A_1 \xrightarrow{\tilde{a}_{n-1}} \cdots \xrightarrow{\tilde{a}_2} A_{n-1} \xrightarrow{\tilde{a}_1} A_n$$

$$U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \cdots \xrightarrow{u_{n-1}} U_{n-1} \xrightarrow{u_n} U_n.$$

To each partition $\bar{m} = (m_1, \dots, m_k)$ of n , by induction on k , we associate the set $\text{Seq}(\sigma, \tilde{a}, \bar{m})$ of special sequences of morphisms obtained from shuffle products of \tilde{a} and certain paths corresponding to \bar{m} (4.3). The sets $\text{Seq}(\sigma, \tilde{a}, \bar{m})$ are crucial in defining the cochain map \mathcal{F} (4.4). Further, we define the sets $\text{Seqq}(\sigma, \bar{m})$ containing *conditioned shuffle product* of certain paths corresponding to \bar{m} (4.9), these sets are essentially used in defining the cochain map \mathcal{G} (4.10).

The proof that \mathcal{F} and \mathcal{G} are inverse quasi-isomorphisms has two parts. The fact that $\mathcal{G}\mathcal{F}(\phi) = \phi$ for any normalized reduced cochain ϕ can be proved by direct computation (Proposition 4.9). The hard part is Theorem 4.6, which relies on the construction of a homotopy $T : \mathcal{F}\mathcal{G} \rightarrow 1$. By induction, we define the family $(\Omega_n)_{n \geq 1}$ in (4.12) which is essentially used in defining the homotopy T . This homotopy is new even in the presheaf case.

Theorem 4.6 has an important consequence, as by the Homotopy Transfer Theorem [11, Theorem 10.3.9], we can transfer the dg Lie algebra structure present on $\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}})$ (see [12]) in order to obtain an L_∞ -structure on $\mathbf{C}_{\text{GS}}^\bullet(\mathcal{A})$. This L_∞ -structure determines the higher deformation theory of \mathcal{A} as a prestack, which thus becomes equivalent to the higher deformation theory of the \mathcal{U} -graded category $\tilde{\mathcal{A}}$ described in [12]. A more detailed elaboration of this L_∞ -structure, as well as a comparison with the L_∞ deformation complex described in the literature in an operadic context [7], [5], [15] will appear in [4].

4.1. The cochain map \mathcal{F} . Following [12] the Hochschild complex $(\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M}), \delta)$ of the \mathcal{U} -graded category $\tilde{\mathcal{A}}$ is defined as

$$\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M}) = \prod_{\substack{u_1, \dots, u_n \\ A_0, A_1, \dots, A_n}} = \text{Hom}_k\left(\bigotimes_{i=1}^n \tilde{\mathcal{A}}_{u_{n+1-i}}(A_{n-i}, A_{n+1-i}), \tilde{\mathcal{A}}_{u_n \cdots u_1}(A_0, A_n)\right)$$

where δ is the usual Hochschild differential.

In order to define the cochain map

$$\mathcal{F} : \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M) \rightarrow \mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M})$$

we need to introduce the following notations. For each $n \in \mathbb{N}$ denote the set of all partitions of n as

$$\text{Part}(n) = \{\bar{m} = (m_k, \dots, m_1) \mid m_k + \cdots + m_1 = n, k \geq 1, m_i \geq 1\}$$

We define $(-1)^{\bar{m}} = (-1)^{n-k}$ for $\bar{m} = (m_k, \dots, m_1)$.

Let $\sigma = (u_1, \dots, u_n)$ be a n -simplex as in (3.8), denote $\|\sigma\| = u_n \cdots u_1$. For $i \leq k$ denote by $\sigma[m_i]$ the m_i -simplex $(u_{m_k+\dots+m_{i+1}+1}, \dots, u_{m_k+\dots+m_i})$. For example, we have $\sigma[m_k] = (u_1, \dots, u_{m_k})$ and $\sigma[m_{k-1}] = (u_{m_k+1}, \dots, u_{m_k+m_{k-1}})$. Put $c^{\sigma, \bar{m}} = \|r\|$ for an arbitrary $r \in \mathcal{P}(\|\sigma[m_k]\|, \dots, \|\sigma[m_1]\|)$.

Given $A_i \in \text{Ob}(\tilde{\mathcal{A}}(U_i))$, consider $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n)$ where

$$\tilde{a}_i \in \tilde{\mathcal{A}}_{u_{n+1-i}}(A_{n-i}, A_{n+1-i}) = \mathcal{A}(U_{n-i})(A_{n-i}, u_{n+1-i}^* A_{n+1-i})$$

as follows:

$$(4.2) \quad A_0 \xrightarrow{\tilde{a}_n} A_1 \xrightarrow{\tilde{a}_{n-1}} \dots \xrightarrow{\tilde{a}_2} A_{n-1} \xrightarrow{\tilde{a}_1} A_n$$

$$U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \dots \xrightarrow{u_{n-1}} U_{n-1} \xrightarrow{u_n} U_n.$$

For each $i = 1, \dots, n$, denote

$$\begin{aligned} \tilde{a}_i &= u_1^* \cdots u_{n-i}^* \tilde{a}_i \in \mathcal{A}(U_0)(u_1^* \cdots u_{n-i}^* A_{n-i}, u_1^* \cdots u_{n+1-i}^* A_{n+1-i}); \\ \tilde{a}_{i, \dots, n} &= \tilde{a}_i \circ \dots \circ \tilde{a}_n \in \mathcal{A}(U_0)(A_0, u_1^* \cdots u_{n+1-i}^* A_{n+1-i}). \end{aligned}$$

Given a partition $\bar{m} = (m_k, \dots, m_1) \in \text{Part}(n)$, denote

$$\tilde{a}[m_i] = \tilde{a}_{m_{i-1}+\dots+m_1+1} \circ \dots \circ \tilde{a}_{m_i+\dots+m_1},$$

thus $\tilde{a}[m_1] = \tilde{a}_1 \circ \dots \circ \tilde{a}_{m_1}$ and $\tilde{a}[m_k] = \tilde{a}_{n-m_k+1, \dots, n}$.

For $r = (r_1, \dots, r_{n-1}) \in \mathcal{P}(\sigma)$, we obtain the following n -simplex in $\mathcal{A}(U_0)$:

$$(r(A_n), \tilde{a}_{1, \dots, n}) \equiv (r_1(A_n), \dots, r_{n-1}(A_n), \tilde{a}_{1, \dots, n}).$$

Now for each partition $\bar{m} = (m_k, \dots, m_1)$ of n we define by induction a set

$$(4.3) \quad \text{Seq}(\sigma, \bar{m}) \equiv \text{Seq}(\sigma, \tilde{a}, \bar{m}) \subseteq \mathcal{N}_n(\mathcal{A}(U_0))(A_0, \sigma[m_k]^* \cdots \sigma[m_1]^* A_n)$$

along with a sign map

$$\text{Seq}(\sigma, \bar{m}) \longrightarrow \{1, -1\} : \xi \longmapsto \text{sign}(\xi) \equiv (-1)^\xi.$$

Simultaneously, for each sequence $\xi \in \text{Seq}(\sigma, \bar{m})$ we define the *formal sequence* $\underline{\xi}$ of ξ , then denote the set of all these formal sequences

$$\underline{\text{Seq}}(\sigma, \bar{m}) = \{\underline{\xi} \mid \xi \in \text{Seq}(\sigma, \bar{m})\}.$$

- For $k = 1$, $\bar{m} = (m_1)$ where $m_1 = n$, we define

$$\text{Seq}(\sigma, \bar{m}) = \{(r(A_n), \tilde{a}_{1, \dots, n}) \mid r \in \mathcal{P}(\sigma)\}.$$

For each element $\xi = (r(A_n), \tilde{a}_{1, \dots, n}) \in \text{Seq}(\sigma, \bar{m})$ we define

$$\text{sign}(\xi) = (-1)^r.$$

The formal sequence of ξ is defined to be

$$\underline{\xi} = (r, \tilde{a}_{1, \dots, n}).$$

- For $k \geq 2$, $R_{m_k}\sigma$ is an $(n - m_k)$ -simplex. Let $\xi = (\xi_1, \dots, \xi_{n-m_k}) \in \text{Seq}(R_{m_k}\sigma, (m_{k-1}, \dots, m_1)) \subseteq \mathcal{N}_{n-m_k}(\mathcal{A}(U_{m_k}))(A_{m_k}, \sigma[m_{k-1}]^* \cdots \sigma[m_1]^* A_n)$. Let $\underline{\xi} = (\underline{\xi}_1, \dots, \underline{\xi}_{n-m_k})$ be the formal sequence of ξ .

- Case $m_k = 1$. Let $u_1^*\xi = (u_1^*\xi_1, \dots, u_1^*\xi_{n-m_k})$, then we obtain the concatenation

$$(u_1^*\xi, \tilde{a}_n) \in \mathcal{N}_n(\mathcal{A}(U_0))(A_0, \sigma[m_k]^* \cdots \sigma[m_1]^* A_n).$$

We define

$$\text{Seq}(\sigma, \bar{m}) = \{(u_1^*\xi, \tilde{a}_n) \mid \xi \in \text{Seq}(R_{m_k}\sigma, (m_{k-1}, \dots, m_1))\}.$$

For each element $\xi' = (u_1^* \xi, \tilde{a}_n) \in \text{Seq}(\sigma, \bar{m})$, we define

$$\text{sign}(\xi') = \text{sign}(\xi).$$

Now we define the formal sequence of ξ' to be

$$\underline{\xi}' = (\underline{\xi}, \tilde{a}_n).$$

- (ii) Case $m_k \geq 2$. For $s \in \mathcal{P}(L_{m_k} \sigma)$ and $\beta \in S_{n-m_k, m_k-1}$, we obtain the shuffle

$$\xi *_{\beta} s \in \mathcal{N}_{n-1}(\mathcal{A}(U_0))(\sigma[m_k]^{\sharp} A_{m_k}, \sigma[m_k]^* \cdots \sigma[m_1]^* A_n)$$

taken with respect to evaluation of functors. Concatenation with $\tilde{a}_{n+1-m_k, \dots, n} \in \mathcal{A}(U_0)(A_0, \sigma[m_k]^{\sharp} A_{m_k})$ yields an n -simplex

$$(\xi *_{\beta} s, \tilde{a}_{n+1-m_k, \dots, n}) \in \mathcal{N}_n(\mathcal{A}(U_0))(A_0, \sigma[m_k]^* \cdots \sigma[m_1]^* A_n).$$

Put $m' = (m_{k-1}, \dots, m_1)$. We define

$$\text{Seq}(\sigma, \bar{m}) = \{(\xi *_{\beta} r, \tilde{a}_{n+1-m_k, \dots, n}) \mid \xi \in \text{Seq}(R_{m_k} \sigma, m'), r \in \mathcal{P}(R_{m_k} \sigma), \beta \in S_{n-m_k, m_k-1}\}.$$

For each element $\xi' = (\xi *_{\beta} r, \tilde{a}_{n+1-m_k, \dots, n}) \in \text{Seq}(\sigma, \bar{m})$ we define

$$\text{sign}(\xi') = (-1)^r (-1)^{\beta} \text{sign}(\xi).$$

Let $\beta(\underline{\xi}, r)$ be the formal shuffle product of $\underline{\xi}$ and r . The formal sequence of ξ' is defined to be

$$\underline{\xi}' = (\beta^{(0)}(\underline{\xi}, r), \tilde{a}_{n+1-m_k, \dots, n}).$$

Example 4.1. Consider a partition $m = (m_3, m_2, m_1)$ of n where $m_i \geq 2$. Each element $\xi \in \text{Seq}(\sigma, \bar{m})$ is of the form

$$\xi = \left(((r_1, \tilde{a}[m_1]) *_{\beta_1} r_2, \tilde{a}[m_2]) *_{\beta_2} r_3, \tilde{a}[m_3] \right)$$

where $r_1 \in \mathcal{P}(\sigma[m_1])$, $r_2 \in \mathcal{P}(\sigma[m_2])$, $r_3 \in \mathcal{P}(\sigma[m_3])$ and $\beta_1 \in S_{m_1, m_2-1}$, $\beta_2 \in S_{m_1+m_2, m_3-1}$.

Now we are able to define the maps $F_p : \mathbf{C}_{\text{GS}}^{p, n-p}(\mathcal{A}, M) \rightarrow \mathbf{C}^n(\tilde{\mathcal{A}}, \tilde{M})$. Let $\sigma = (u_1, \dots, u_n)$ be an n -simplex and $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n)$ as in (4.2). For each cochain $\phi = (\phi_{p,q}) \in \mathbf{C}_{\text{GS}}^n(\mathcal{A}, M)$, we define

$$(4.4) \quad \mathcal{F}_p \phi_{p, n-p}(\tilde{a}) = \sum_{\bar{m} \in \text{Part}(n-p)} \sum_{\xi \in \text{Seq}(R_p \sigma, \bar{m})} (-1)^{\bar{m} + \xi} \mathcal{F}_p^{\sigma, \bar{m}, A_n} \phi_{p, n-p}^{L_p \sigma}(\xi) \tilde{a}_{n+1-p, \dots, n}$$

where $\mathcal{F}_p^{\sigma, \bar{m}, A_n} = c^{\sigma, p, A_n}(L_p \sigma)^* c^{R_p \sigma, \bar{m}, A_n}$. The map \mathcal{F} is as follows

$$\mathcal{F}(\phi) = \sum_{p+q=n} \mathcal{F}_p(\phi_{p,q}).$$

Proposition 4.2. *The map \mathcal{F} commutes with differentials. More precisely, let $p+q=n-1$, for $\phi \in \mathbf{C}_{\text{GS}}^{p,q}(\mathcal{A}, M)$, then $F(d\phi) = \delta(F\phi)$.*

Proof. Let $\sigma = (u_1, \dots, u_n)$ be a n -simplex as in (3.8), and let $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n)$ as in (4.2). First, we prove that $\mathcal{F}(d\phi) = \delta(\mathcal{F}\phi)$ for the case $\phi \in \mathbf{C}_{\text{GS}}^{0, n-1}(\mathcal{A}, M)$.

The equation

$$(4.5) \quad \sum_{i=0}^n (-1)^i \mathcal{F} d_{\text{Hoch}}^i \phi + (-1)^n \mathcal{F}(d_{\text{simp}}^0 - d_{\text{simp}}^1) \phi + \sum_{i=2}^n \mathcal{F} d_i \phi = \sum_{i=0}^n \delta_n \mathcal{F} \phi$$

holds true if the following equations hold true:

$$(i) \quad -(-1)^n \mathcal{F}d_{\text{Hoch}}^n \phi = (-1)^{n+1} \mathcal{F}d_{\text{simp}}^1 \phi + \sum_{i=2}^n \mathcal{F}d_i \phi;$$

$$(ii) \quad \mathcal{F}d_{\text{simp}}^0 \phi = \delta_n \mathcal{F} \phi;$$

$$(iii) \quad \mathcal{F}d_{\text{Hoch}}^0 \phi + \sum_{i=1}^{n-1} (-1)^i \mathcal{F}d_{\text{Hoch}}^i \phi = \sum_{i=0}^{n-1} (-1)^i \delta_i \mathcal{F} \phi.$$

Step 1. We prove the equation (i). Note that $L_0 \sigma = (U_0)$ is a 0-simplex, by definition, we have

$$(-1)^{n+1} (\mathcal{F}d_{\text{Hoch}}^n \phi)^\sigma(\tilde{a}) = \sum_{\bar{m} \in \text{Part}(n)} \sum_{\xi \in \text{Seq}(\sigma, \bar{m})} T(\bar{m}, \xi)$$

where $T(\bar{m}, \xi) = (-1)^{n+1} (-1)^{\bar{m}+\xi} c^{\sigma, \bar{m}, A_n} (d_{\text{Hoch}}^n \phi)^{U_0}(\xi)$. On the right hand side, we have

$$(-1)^{n+1} (\mathcal{F}d_{\text{simp}}^1 \phi)^\sigma(\tilde{a}) = \sum_{\substack{\bar{m}' \in \text{Part}(n-1) \\ \xi' \in \text{Seq}(R_1 \sigma, \bar{m}')}} (-1)^{n+1} (-1)^{\bar{m}'+\xi'} \mathcal{F}_1^{\sigma, \bar{m}', A_n} (d_{\text{simp}}^1 \phi)^{L_1 \sigma}(\xi') \tilde{a}_n$$

where $(d_{\text{simp}}^1 \phi)^{L_1 \sigma}(\xi') \tilde{a}_n = \phi^{U_0}(u_1^* \xi') \tilde{a}_n$.

For each $\bar{m}' = (m'_k, \dots, m'_1) \in \text{Part}(n-1)$ and $\xi' \in \text{Seq}(R_1 \sigma, \bar{m}')$, let $\bar{m} = (1, m'_k, \dots, m'_1) \in \text{Part}(n)$. Then by definition, there exists a unique element $\xi \in \text{Seq}(\sigma, \bar{m})$ such that $\xi = (u_1^* \xi', \tilde{a}_n)$. Hence, we get

$$T(\xi, \bar{m}) = (-1)^{n+1} (-1)^{\bar{m}'+\xi'} \mathcal{F}_1^{\sigma, \bar{m}', A_n} (d_{\text{simp}}^1 \phi)^{L_1 \sigma}(\xi') \tilde{a}_n.$$

So all the terms occurring in $(-1)^{n+1} (\mathcal{F}d_{\text{simp}}^1 \phi)^\sigma(\tilde{a})$ are canceled.

For $2 \leq i \leq n$, we have

$$(\mathcal{F}d_i \phi)^\sigma(\tilde{a}) = \sum_{\substack{\bar{m}' \in \text{Part}(n-i) \\ \xi' \in \text{Seq}(R_i \sigma, \bar{m}')}} (-1)^{\bar{m}'+\xi'} c^{\sigma, i, A_n} (L_i \sigma)^* c^{R_i \sigma, \bar{m}', A_n} (d_i \phi)^{L_i \sigma}(\xi') \tilde{a}_{n+1-i, \dots, n}$$

where $(d_i \phi)^{L_i \sigma}(\xi') = \sum_{r \in \mathcal{P}(L_i \sigma), \beta \in S_{n-i, i-1}} (-1)^{n-i} (-1)^{r+\beta} \phi^{U_0}(\xi' *_\beta r)$. For each $\bar{m}' \in \text{Part}(n-i)$, $r \in \mathcal{P}(L_i \sigma)$ and $\beta \in S_{n-i, i-1}$, there exists a unique element $\xi \in \text{Seq}(\sigma, \bar{m})$, where $\bar{m} = (n-i, \bar{m}') \in \text{Part}(n)$, such that $\xi = (\xi' *_\beta r, \tilde{a}_{n+1-k, \dots, n})$. We

get $T(\xi, \bar{m}) = (-1)^{\bar{m}'+\xi'} (-1)^{n-i} (-1)^{r+\beta} c^{\sigma, i, A_n} (L_i \sigma)^* c^{R_i \sigma, \bar{m}', A_n} \phi^{U_0}(\xi' *_\beta r) \tilde{a}_{n+1-k, \dots, n}$.

So every term occurring in $(\mathcal{F}d_i \phi)^\sigma(\tilde{a})$ is cancelled.

Step 2. The equation (ii) is obvious. We prove the equation (iii). For $i = 1, \dots, (n-1)$, we have

$$(\mathcal{F}d_{\text{Hoch}}^i \phi)^\sigma(\tilde{a}) = \sum_{\bar{m} \in \text{Part}(n), \xi \in \text{Seq}(\sigma, \bar{m})} (-1)^{\bar{m}+\xi} c^{\sigma, \bar{m}, A_n} (d_{\text{Hoch}}^i \phi)^{U_0}(\xi).$$

Let $\bar{m} = (m_k, \dots, m_1)$. Assume $\xi = (\xi_1, \dots, \xi_n) \in \text{Seq}(\sigma, \bar{m})$, we have

$$(d_{\text{Hoch}}^i \phi)^{U_0} = \phi^{U_0}(\xi_1, \dots, \xi_i \xi_{i+1}, \dots, \xi_n).$$

Let $\underline{\xi} = (\underline{\xi}_1, \dots, \underline{\xi}_n) \in \text{Seq}(\sigma, \bar{m})$ be the formal sequence of ξ . Then $(\underline{\xi}_i, \underline{\xi}_{i+1})$ can only be one of the following cases

$$(\underline{\xi}_i, \underline{\xi}_{i+1}) = \begin{cases} (r_j, s_l) \text{ or } (s_l, r_j) & \text{for } r \in \mathcal{P}(\sigma[m_t]), s \in \mathcal{P}(\sigma[m_{t+1}]); \\ (\tilde{a}[m_t], r_j) \text{ or } (r_j, \tilde{a}[m_t]) & \text{for } r \in \mathcal{P}(\sigma[m_{t+1}]); \\ (\tilde{a}[m_t], \tilde{a}[m_{t+1}]) & \text{for some } t; \\ (r_{m_t-1}, \tilde{a}[m_t]) & \text{for } r \in \mathcal{P}(\sigma[m_t]). \end{cases}$$

Case 1. Assume that $(\underline{\xi}_i, \underline{\xi}_{i+1}) = (r_j, s_l)$ or (s_l, r_j) , for some $r = (r_1, \dots, r_{m_t-1}) \in \mathcal{P}(\sigma[m_t])$ and $s = (s_1, \dots, s_{m_{t+1}-1}) \in \mathcal{P}(\sigma[m_{t+1}])$. There exists a unique element

$\xi' \in \text{Seq}(\sigma, \bar{m})$ such that its formal sequence satisfies $\underline{\xi}' = (\underline{\xi}_1, \dots, \underline{\xi}_{i-1}, \underline{\xi}_{i+1}, \underline{\xi}_i, \underline{\xi}_{i+2}, \dots, \underline{\xi}_n)$. Hence

$$(-1)^{\bar{m}+\xi} c^{\sigma, \bar{m}, A_n} (d_{\text{Hoch}}^i \phi)^{U_0}(\xi) + (-1)^{\bar{m}+\xi'} c^{\sigma, \bar{m}, A_n} (d_{\text{Hoch}}^i \phi)^{U_0}(\xi') = 0.$$

The same argument applies for the cases $(\underline{\xi}_i, \underline{\xi}_{i+1}) = (\tilde{a}[m_t], r_j)$ or $(r_j, \tilde{a}[m_t])$.

Case 2. Assume $(\underline{\xi}_i, \underline{\xi}_{i+1}) = (\tilde{a}[m_t], \tilde{a}[m_{t+1}])$ for some $1 \leq t \leq k-1$. Without loss of generality we assume that $\underline{\xi} = (\underline{\xi}_1, \dots, \underline{\xi}_j, r, s, \tilde{a}[m_t], \tilde{a}[m_{t+1}], \underline{\xi}_{j+m_t+m_{t+1}+1}, \dots, \underline{\xi}_n)$ for some paths $r \in \mathcal{P}(\sigma[m_t])$, $s \in \mathcal{P}(\sigma[m_{t+1}])$. Denote $\gamma = \sigma[m_t] \sqcup \sigma[m_{t+1}]$ the concatenation of the simplices $\sigma[m_{t+1}]$ and $\sigma[m_t]$, then $(c^{\gamma, m_{t+1}}, r, s)$ is a path in $\mathcal{P}(\gamma)$. We have

$$(\mathcal{F}d_{\text{Hoch}}^0 \phi)^\sigma(\tilde{a}) = \sum_{\bar{m}' \in \text{Part}(n)} \sum_{\xi' \in \text{Seq}(\sigma, \bar{m}')} (-1)^{\bar{m}'+\xi'} c^{\sigma, \bar{m}', A_n} (d_{\text{Hoch}}^0 \phi)^{U_0}(\xi').$$

Consider the partition $\bar{m}' = (m_k, \dots, m_{t+1} + m_t, \dots, m_1)$, there exists a unique element $\xi' \in \text{Seq}(\sigma, \bar{m}')$ such that its formal sequence satisfies

$$\begin{aligned} \underline{\xi}' &= (c^{\gamma, m_{t+1}}, \underline{\xi}_1, \dots, \underline{\xi}_j, r, s, \tilde{a}[m_t], \tilde{a}[m_{t+1}], \underline{\xi}_{j+m_t+m_{t+1}+1}, \dots, \underline{\xi}_n) \\ &= (c^{\gamma, m_{t+1}}, \underline{\xi}). \end{aligned}$$

We obtain $(-1)^{\bar{m}'+\xi'} c^{\sigma, \bar{m}', A_n} (d_{\text{Hoch}}^0 \phi)^{U_0}(\xi') + (-1)^{\bar{m}+\xi} c^{\sigma, \bar{m}, A_n} (d_{\text{Hoch}}^0 \phi)^{U_0}(\xi) = 0$.

Case 3. Assume that $(\underline{\xi}_i, \underline{\xi}_{i+1}) = (r_{m_t-1}, \tilde{a}[m_t])$ for some $r = (r_1, \dots, r_{m_t-1}) \in \mathcal{P}(\sigma[m_t])$. We have $r_{m_t-1} = \epsilon^{\sigma[m_t], j}$ for some $1 \leq j \leq m_t - 1$. Let $j' = n + 1 - (m_k + \dots + m_{t+1} + j) = m_1 + \dots + m_t + 1 - j$. In the right hand side of equation (iii), we have

$$\begin{aligned} (-1)^{j'} (\delta_{j'} \mathcal{F} \phi)^\sigma(\tilde{a}) &= (-1)^{j'} (\mathcal{F} \phi)^{\partial_{n-j'} \sigma}(\partial_{j'} \tilde{a}) \\ &= \sum_{\bar{m}' \in \text{Part}(n-1)} \sum_{\xi' \in \text{Seq}(\partial_{n-j'} \sigma, \bar{m}')} (-1)^{j'} (-1)^{\bar{m}'+\xi'} c^{\partial_{n-j'} \sigma, \bar{m}', A_n} \phi^{U_0}(\xi'). \end{aligned}$$

Choose $\bar{m}' = (m_k, \dots, m_t - 1, \dots, m_1) \in \text{Part}(n-1)$. There exists a unique element $\xi' \in \text{Seq}(\partial_{n-j'} \sigma, \bar{m}')$ such that

$$(-1)^{j'} (-1)^{\bar{m}'+\xi'} c^{\partial_{n-j'} \sigma, \bar{m}', A_n} \phi^{U_0}(\xi') = (-1)^{\bar{m}+\xi} c^{\sigma, \bar{m}, A_n} (d_{\text{Hoch}}^i \phi)^{U_0}(\xi).$$

After considering all cases 1,2,3 as above, we find that all the terms occurring in $\sum_{i=1}^{n-1} (\mathcal{F}d_{\text{Hoch}}^i \phi)^\sigma(\tilde{a})$ and $\sum_{i=1}^{n-1} (\delta_i \mathcal{F} \phi)^\sigma(\tilde{a})$ are canceled. The remaining terms in $(\mathcal{F}d_{\text{Hoch}}^0 \phi)^\sigma(\tilde{a})$ are only

$$\sum_{\bar{m}' = (m'_k, \dots, m'_2, 1) \in \text{Part}(n)} \sum_{\xi' \in \text{Seq}(\sigma, \bar{m}')} (-1)^{\bar{m}'+\xi'} c^{\sigma, \bar{m}', A_n} (d_{\text{Hoch}}^0 \phi)^{U_0}(\xi')$$

which are in turn canceled by all the terms in $(\delta_0 \mathcal{F} \phi)^\sigma(\tilde{a})$. We conclude that the equation (iii) holds.

In the general case, we consider $\phi \in \mathbf{C}_{\text{GS}}^{p, n-1-p}(\mathcal{A}, M)$ for $p > 0$. Applying the same arguments as above, we can prove the following equations hold true:

$$(i') \quad -(-1)^{n-p} \mathcal{F}d_{\text{Hoch}}^{n-p} \phi = (-1)^{n+1-p} \mathcal{F}d_{\text{simp}}^{p+1} \phi + \sum_{i=2}^{n-p} \mathcal{F}d_i \phi;$$

$$(ii') \quad \mathcal{F}d_{\text{simp}}^i \phi = \delta_{n-i} \mathcal{F} \phi \text{ for } i = 0, \dots, p;$$

$$(iii'') \quad \mathcal{F}d_{\text{Hoch}}^0 \phi + \sum_{i=1}^{n-p-1} (-1)^i \mathcal{F}d_{\text{Hoch}}^i \phi = \sum_{i=0}^{n-p-1} (-1)^i \delta_i \mathcal{F} \phi.$$

These equations yield

$$\sum_{i=0}^n (-1)^i \mathcal{F} d_{\text{Hoch}}^i \phi + (-1)^n \mathcal{F} \left(\sum_{i=0}^{p+1} d_{\text{simp}}^i \right) \phi + \sum_{i=2}^n \mathcal{F} d_i \phi = \sum_{i=0}^n \delta_n \mathcal{F} \phi,$$

which means $\mathcal{F}(d\phi) = \delta(\mathcal{F}\phi)$. \square

4.2. The cochain map \mathcal{G} . In this section we define the cochain map

$$\mathcal{G} : \mathbf{C}_{\mathcal{U}}^{\bullet}(\tilde{\mathcal{A}}, \tilde{M}) \longrightarrow \mathbf{C}_{\text{GS}}^{\bullet}(\mathcal{A}, M).$$

Consider a p -simplex $\sigma = (u_1, \dots, u_p) \in \mathcal{N}_p(\mathcal{U})$ as follows

$$(4.6) \quad \sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} \dots \xrightarrow{u_{p-1}} U_{p-1} \xrightarrow{u_p} U_p)$$

and a q -simplex $a = (a_1, \dots, a_q) \in \mathcal{N}(\mathcal{A}(U_p))_q$ as follows

$$(4.7) \quad a = (A_0 \xrightarrow{a_q} A_1 \xrightarrow{a_{q-1}} \dots \xrightarrow{a_2} A_{q-1} \xrightarrow{a_1} A_q).$$

Using conditioned shuffles, we will describe several ways to build a $(p+q)$ -simplex in $\mathcal{N}_{p+q}(\tilde{\mathcal{A}})$ from these data. Let $\bar{m} = (m_k, \dots, m_1)$ be a partition of p with $m_i \geq 1$ for all i and let $\beta \in \bar{S}_{\bar{m}}$ be a conditioned \bar{m} -shuffle as defined in §3.1. For $1 \leq i \leq k$, let $r^i = (r_1^i, \dots, r_{m_i-1}^i) \in \mathcal{P}(\sigma[m_i])$ be a path and consider the associated m_i -simplex

$$(4.8) \quad \bar{r}^i = (1_{\sigma[m_i]^*}, r_1^i, \dots, r_{m_i-1}^i) \in \mathcal{N}_{m_i}(\mathcal{C}_i).$$

where

$$\mathcal{C}_i = \text{Fun}(\mathcal{A}(U_{p-m_1 \dots - m_{i-1}}), \mathcal{A}(U_{p-m_1 \dots - m_i})).$$

First, consider the formal shuffle by β of the associated tuples $(\bar{r}^i)_i$ as described in (3.2). Assume that

$$\beta^{(0)}((\bar{r}^i)_i) = \underline{s} = (\underline{s}_1, \dots, \underline{s}_p).$$

Since β is a conditioned shuffle, there are uniquely determined numbers $\gamma_l \geq 1$, $1 \leq l \leq k$ such that $\underline{s}_1 = 1_{\sigma[m_1]^*}$, $\underline{s}_{\gamma_1+1} = 1_{\sigma[m_2]^*}$, \dots , $\underline{s}_{\sum_{i=1}^l \gamma_i+1} = 1_{\sigma[l+1]^*}$, \dots , $\underline{s}_{\sum_{i=1}^{k-1} \gamma_i+1} = 1_{\sigma[m_k]^*}$ and $\gamma_k = p - \sum_{i=1}^{k-1} \gamma_i$. Following the pattern explained at the end of §3.1, we obtain the sequence

$$(\hat{c}^1, \dots, \hat{c}^k) \in \prod_{l=1}^k \mathcal{N}_{\gamma_l} \left(\prod_{i=1}^l \mathcal{C}_i \right).$$

Using the composition of functors as in Remark 3.3, we obtain the following sequence which we define as the shuffle product of $(\bar{r}^i)_i$ by β

$$\beta((\bar{r}^i)_i) := (\bar{c}^1, \dots, \bar{c}^k) \in \prod_{l=1}^k \mathcal{N}_{\gamma_l}(\mathcal{D}_l)$$

where

$$\mathcal{D}_l = \text{Fun}(\mathcal{A}(U_p), \mathcal{A}(U_{p-m_1 \dots - m_l})).$$

We denote by $\text{Seqq}(\sigma, \bar{m})$ the set of all such conditioned shuffle products. Thus

$$(4.9) \quad \text{Seqq}(\sigma, \bar{m}) = \{ \beta((\bar{r}^i)_i) \mid \beta \in \bar{S}_{\bar{m}}, \bar{r}^i = (1_{\sigma[m_i]^*}, r^i), r^i \in \mathcal{P}(\sigma[m_i]) \}.$$

For each $\zeta = \beta((\bar{r}^i)_i) \in \text{Seqq}(\sigma, \bar{m})$, we denote the formal sequence $\beta^{(0)}((\bar{r}^i)_i)$ of ζ by $\underline{\zeta}$, and denote the set of all such formal sequence as

$$\underline{\text{Seqq}}(\sigma, \bar{m}) = \{ \underline{\zeta} \mid \zeta \in \text{Seqq}(\sigma, \bar{m}) \}.$$

We define

$$\text{sign}(\zeta) = \text{sign}(\beta((\bar{r}^i)_i)) = (-1)^\beta \prod_{i=1}^k (-1)^{r^i}$$

and equip this shuffle product with a certain underlying simplex denoted by $\text{simp}(\beta((\bar{r}^i)_i))$. Writing

$$\bar{c}^l = (\bar{c}_1^l, \dots, \bar{c}_{\gamma_l}^l)$$

we define

$$\text{simp}(\bar{c}_1^l) = (U_{p-m_1 \dots - m_l} \xrightarrow{\|\sigma[m_l]\|} U_{p-m_1 \dots - m_{l-1}});$$

$$\text{simp}(\bar{c}_j^l) = (U_{p-m_1 \dots - m_l} \xrightarrow{1} U_{p-m_1 \dots - m_l}), \quad j > 1.$$

The simplex $\text{simp}(\bar{c}^l)$ is obtained by concatenation of $(\text{simp}(\bar{c}_j^l))_j$, the simplex $\text{simp}(\beta((\bar{r}^i)_i)) \equiv \text{simp}(\bar{c}^1, \dots, \bar{c}^k)$ is obtained by concatenation of the simplices $(\text{simp}(\bar{c}^l))_l$.

Example 4.3. Consider the simplex $\sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} U_2 \xrightarrow{u_3} U_3 \xrightarrow{u_4} U_4)$ and the partition $\bar{m} = (m_2, m_1) = (2, 2)$. There are three conditioned formal shuffles $(1_{\sigma[m_1]}^*, c^{u_3, u_4}, 1_{\sigma[m_2]}^*, c^{u_1, u_2}); (1_{\sigma[m_1]}^*, 1_{\sigma[m_2]}^*, c^{u_3, u_4}, c^{u_1, u_2}); (1_{\sigma[m_1]}^*, 1_{\sigma[m_2]}^*, c^{u_1, u_2}, c^{u_3, u_4})$. The set $\text{Seq}(\sigma, (m_2, m_1))$ consists of following sequences:

$$\begin{pmatrix} \bullet & \xrightarrow{c^{u_1, u_2} u_3^* u_4^*} & \bullet & \xrightarrow{1_{\sigma[m_2]}^* u_3^* u_4^*} & \bullet & \xrightarrow{c^{u_3, u_4}} & \bullet & \xrightarrow{1_{\sigma[m_1]}^*} & \bullet \\ U_0 & \xrightarrow{1} & U_0 & \xrightarrow{u_2 u_1} & U_2 & \xrightarrow{1} & U_2 & \xrightarrow{u_4 u_3} & U_4 \end{pmatrix}$$

$$\begin{pmatrix} \bullet & \xrightarrow{c^{u_1, u_2} u_3^* u_4^*} & \bullet & \xrightarrow{(u_2 u_1)^* c^{u_3, u_4}} & \bullet & \xrightarrow{1_{\sigma[m_2]}^* (u_4 u_3)^*} & \bullet & \xrightarrow{1_{\sigma[m_1]}^*} & \bullet \\ U_0 & \xrightarrow{1} & U_0 & \xrightarrow{1} & U_0 & \xrightarrow{u_2 u_1} & U_2 & \xrightarrow{u_4 u_3} & U_4 \end{pmatrix}$$

$$\begin{pmatrix} \bullet & \xrightarrow{u_1^* u_2^* c^{u_3, u_4}} & \bullet & \xrightarrow{c^{u_1, u_2} (u_4 u_3)^*} & \bullet & \xrightarrow{1_{\sigma[m_2]}^* (u_4 u_3)^*} & \bullet & \xrightarrow{1_{\sigma[m_1]}^*} & \bullet \\ U_0 & \xrightarrow{1} & U_0 & \xrightarrow{1} & U_0 & \xrightarrow{u_2 u_1} & U_2 & \xrightarrow{u_4 u_3} & U_4 \end{pmatrix}.$$

Next consider a shuffle permutation $\omega \in S_{p,q}$. We are now to define the shuffle product of a and $(\bar{c}^1, \dots, \bar{c}^k)$ by ω to be the element

$$(\hat{b}^0, \hat{b}^1, \dots, \hat{b}^k) \equiv a *_{\omega} (\bar{c}^1, \dots, \bar{c}^k) \in \mathcal{N}_{p+q}(\tilde{\mathcal{A}}).$$

The formal shuffle product $\omega^{(0)}(a, \beta^{(0)}((\bar{r}^i)_i))$ is called the formal sequence of $(\hat{b}^0, \hat{b}^1, \dots, \hat{b}^k)$. First consider the formal shuffle

$$\omega^{(0)}(a; (\bar{c}^1, \dots, \bar{c}^k)) = (b_1, \dots, b_{p+q}).$$

Since ω is shuffle, there are unique numbers t_1, \dots, t_{k+1} such that $b_{t_1+1} = \bar{c}_1^1$, $b_{t_1+t_2+1} = \bar{c}_1^2, \dots, b_{\sum_{i=1}^k t_i+1} = \bar{c}_1^k$ and $t_{k+1} = p+q - \sum_{i=1}^k t_i$. Following the procedure at the end of section 3.1, for $0 \leq l \leq k$ consider

$$a^l = (a_1^l, \dots, a_{j_l}^l) = \{b_{\sum_{i=1}^l t_i+1}, \dots, b_{\sum_{i=1}^{l+1} t_i}\} \cap \{a_1, \dots, a_q\}.$$

Obviously $a^0 = (a_1, \dots, a_{t_1})$. There is unique shuffle $\omega_l \in S_{j_l, \gamma_l}$ such that the formal shuffle product of a^l and \bar{c}^l by ω is exactly

$$(b_{\sum_{i=1}^l t_i+1}, \dots, b_{\sum_{i=1}^{l+1} t_i}).$$

Now we put $\hat{b}^0 = a^0$. For $l = 1 \dots k$, take the shuffle product $\omega_l(a^l, \bar{c}^l)$ with respect to evaluation of functors as in Example 3.2, and put

$$\hat{b}^l = (\hat{b}_1^l, \dots, \hat{b}_{j_l+\gamma_l}^l) = \omega_l(a^l, \bar{c}^l).$$

Now we associate the underlying simplex to $(\hat{b}^0, \hat{b}^1, \dots, \hat{b}^k)$ to show that

$$(\hat{b}^0, \hat{b}^1, \dots, \hat{b}^k) \in \mathcal{N}_{p+q}(\tilde{\mathcal{A}})(\sigma^\# A_0, A_q).$$

We have $\hat{b}_1^l = \sigma[m_l]^* T_{l-1}(A_{\alpha_l})$ for a certain $T_{l-1} \in \mathcal{D}_{l-1}$ and a certain $A_{\alpha_l} \in \{A_0, \dots, A_q\}$. Thus it can be regarded as an element of $\mathcal{N}_1(\tilde{\mathcal{A}})$ as follows:

$$\sigma[m_l]^* T_{l-1} A_{\alpha_l} \xrightarrow{1} T_{l-1} A_{\alpha_l}$$

$$U_{p-m_1 \dots - m_l} \xrightarrow{\|\sigma[m_l]\|} U_{p-m_1 \dots - m_{l-1}}.$$

We consider $\hat{b}_j^l = \mathcal{A}_{U_{p-m_1 \dots - m_l}}(B, B')$ as an element of $\mathcal{N}_1(\tilde{\mathcal{A}})$ as follows:

$$B \xrightarrow{\hat{b}_j^l} B'$$

$$U_{p-m_1 \dots - m_l} \xrightarrow{1} U_{p-m_1 \dots - m_l}.$$

Put

$$\text{simp}(\hat{b}_1^l) = (U_{p-m_1 \dots - m_l} \xrightarrow{\|\sigma[m_l]\|} U_{p-m_1 \dots - m_{l-1}}), \quad l \geq 1;$$

$$\text{simp}(\hat{b}_j^l) = (U_{p-m_1 \dots - m_l} \xrightarrow{1} U_{p-m_1 \dots - m_l}), \quad j > 1;$$

$$\text{simp}(\hat{b}_j^0) = (U_p \xrightarrow{1} U_p), \quad j \geq 0.$$

By concatenation of all these 1-simplices we obtain the simplex $\text{simp}(\hat{b}^0, \hat{b}^1, \dots, \hat{b}^k)$ of $(\hat{b}^0, \hat{b}^1, \dots, \hat{b}^k)$.

Example 4.4. Let $\sigma = (U_0 \xrightarrow{u_1} U_1 \xrightarrow{u_2} U_2)$ and $a \in \mathcal{A}(U_2)(A_0, A_1)$. Let $\bar{m} = (2)$, then $\text{Seqq}(\sigma, (2))$ consist only of the sequence $(1_{(u_2 u_1)^*}, c^{u_1, u_2})$:

$$\left(\begin{array}{ccc} \bullet & \xrightarrow{c^{u_1, u_2}} & \bullet \\ U_0 & \xrightarrow{1} & U_0 \end{array} \xrightarrow{1_{(u_2 u_1)^*}} \begin{array}{ccc} \bullet & & \bullet \\ U_0 & \xrightarrow{u_2 u_1} & U_2 \end{array} \right).$$

The following are shuffle products of a and $(1_{u_2 u_1}, c^{u_1, u_2})$:

$$\begin{aligned} & \left(\begin{array}{ccc} u_1^* u_2^* A_0 & \xrightarrow{c^{u_1, u_2, A_0}} & (u_2 u_1)^* A_0 \\ U_0 & \xrightarrow{1} & U_0 \end{array} \xrightarrow{1_{(u_2 u_1)^*(A_0)}} \begin{array}{ccc} A_0 & \xrightarrow{a} & A_1 \\ U_2 & \xrightarrow{1} & U_2 \end{array} \right) \\ & \left(\begin{array}{ccc} u_1^* u_2^* A_0 & \xrightarrow{c^{u_1, u_2, A_0}} & (u_2 u_1)^* A_0 \\ U_0 & \xrightarrow{1} & U_0 \end{array} \xrightarrow{(u_2 u_1)^* a} \begin{array}{ccc} (u_2 u_1)^* A_1 & \xrightarrow{1_{(u_2 u_1)^*(A_1)}} & A_1 \\ U_0 & \xrightarrow{u_2 u_1} & U_2 \end{array} \right) \\ & \left(\begin{array}{ccc} u_1^* u_2^* A_0 & \xrightarrow{u_1^* u_2^* a} & u_1^* u_2^* A_1 \\ U_0 & \xrightarrow{1} & U_0 \end{array} \xrightarrow{c^{u_1, u_2, A_1}} \begin{array}{ccc} (u_2 u_1)^* A_1 & \xrightarrow{1_{(u_2 u_1)^*(A_1)}} & A_1 \\ U_0 & \xrightarrow{u_2 u_1} & U_2 \end{array} \right). \end{aligned}$$

For each cochain $\psi \in \mathbf{C}_{\mathcal{U}}^{p+q}(\tilde{\mathcal{A}}, \tilde{M})$, we now define

$$(4.10) \quad \mathcal{G}(\psi)^\sigma(a) = \sum_{\substack{\bar{m} \in \text{Part}(n) \\ \zeta \in \text{Seqq}(\sigma, \bar{m})}} \sum_{\beta \in S_{q,p}} (-1)^\beta (-1)^\zeta \psi^{\text{simp}(a^* \zeta)}(a^* \zeta).$$

Proposition 4.5. *The map \mathcal{G} commutes with differentials. Precisely, for $\psi \in \mathbf{C}_{\mathcal{U}}^{n-1}(\tilde{\mathcal{A}}, \tilde{M})$, we have $d\mathcal{G}(\psi) = \mathcal{G}\delta(\psi)$.*

Proof. Assume that $p + q = n$. Let $\sigma = (u_1, \dots, u_p)$ be a p -simplex as in (4.6) and $a = (a_1, \dots, a_q)$ as in (4.7). We prove $(dG(\psi))^\sigma(a) = (G\delta(\psi))^\sigma(a)$.

Put

$$\begin{aligned} \text{LHS} &= (d_0\mathcal{G}\psi)^\sigma(a) + (-1)^n(d_{\text{simp}}\mathcal{G}\psi)^\sigma(a) + (d_2\mathcal{G}\psi)^\sigma(a) + \dots + (d_p\mathcal{G}\psi)^\sigma(a); \\ \text{RHS} &= (\mathcal{G}\delta_0\psi)^\sigma(a) - (\mathcal{G}\delta_1\psi)^\sigma(a) + \dots + (-1)^n(\mathcal{G}\delta_n\psi)^\sigma(a). \end{aligned}$$

We have

$$(-1)^i(\mathcal{G}\delta_i\psi)^\sigma(a) = \sum_{\substack{\bar{m} \in \text{Part}(p), \beta \in S_{q,p} \\ \zeta \in \text{Seqq}(\sigma, \bar{m})}} (-1)^i(-1)^\beta(-1)^\zeta(\delta_i\psi)^{\text{simp}(a*_\beta\zeta)}(a*_\beta\zeta).$$

Denote

$$T(i, \bar{m}, \beta, \zeta) = (-1)^i(-1)^\beta(-1)^\zeta(\delta_i\psi)^{\text{simp}(a*_\beta\zeta)}(a*_\beta\zeta).$$

To prove that LHS = RHS, we show that each term T appearing in the expansion of RHS is either matched with a unique term in the expansion of LHS or canceled out with a term $-T$ in RHS. Simultaneously, this process also shows that every term in LHS is cancelled out.

Take a partition $\bar{m} = (m_k, \dots, m_1) \in \text{Part}(p)$. Fix $\beta \in S_{q,p}$ and $\zeta \in \text{Seqq}(\sigma, \bar{m})$. By definition, there are a unique $\gamma \in \tilde{S}_{\bar{m}}$, $r^{m_i} = (r_1^{m_i}, \dots, r_{m_i-1}^{m_i}) \in \mathcal{P}(\sigma[m_i])$, $i = 1, \dots, k$; such that ζ is the shuffle product

$$(4.11) \quad \zeta = \gamma((\bar{r}^{m_i})_{i=1, \dots, k})$$

where $\bar{r}^{m_i} = (1_{\sigma[m_i]}, r^{m_i})$ as in (4.8).

We denote the shuffle product

$$a*_\beta\zeta = \alpha = (\alpha_1, \dots, \alpha_n)$$

and its formal sequence

$$\underline{\alpha} = (\underline{\alpha}_1, \dots, \underline{\alpha}_n).$$

Step 1. We consider the term $T(0, \bar{m}, \beta, \zeta)$ in RHS. We have

$$(\delta_0\psi)^{\text{simp}(a*_\beta\zeta)}(a*_\beta\zeta) = \mu(\alpha_1, \psi^{\text{simp}(\partial_0\alpha)}(\alpha_2, \dots, \alpha_n))$$

where μ is the composition in the map-graded category $\tilde{\mathcal{A}}$. There are only three cases $\underline{\alpha}_1 = a_1$, $\underline{\alpha}_1 = 1_{u_p}$ or $\underline{\alpha}_1 = 1_{\sigma[m_1]^*}$ where $m_1 \geq 2$.

- Consider the case $\underline{\alpha}_1 = a_1$. We have $\text{simp}(\alpha_1) = (U_p \xrightarrow{1} U_p)$. In the LHS, we consider

$$(d_{\text{Hoch}}^0\mathcal{G}\psi)^\sigma(a) = \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta' \in S_{q-1,p} \\ \zeta' \in \text{Seqq}(\sigma, \bar{m}')}} (-1)^{\beta'}(-1)^{\zeta'}\sigma^*(a_1)\psi^{\text{simp}(\partial_0 a*_\beta'\zeta')}(\partial_0 a*_\beta'\zeta').$$

Choose $\bar{m}' = \bar{m}$ and $\zeta' = \zeta \in \text{Seqq}(\sigma, \bar{m}')$. Then there exists a unique $\beta' \in S_{q-1,p}$ such that $(a_2, \dots, a_q)*_{\beta'}\zeta' = (\alpha_2, \dots, \alpha_n)$. Hence

$$T(0, \bar{m}, \beta, \zeta) = (-1)^{\beta'+\zeta'}\sigma^*(a_1)\psi^{\text{simp}(\partial_0 a*_\beta'\zeta')}(\partial_0 a*_\beta'\zeta').$$

- Consider the case $\underline{\alpha}_1 = 1_{u_p}$. We have $m_1 = 1$, $\text{simp}(\alpha_1) = (U_{p-1} \xrightarrow{u_p} U_p)$, and $(\delta_0\psi)^{\text{simp}(a*_\beta\zeta)}(a*_\beta\zeta) = c^{\sigma, p-1, A_q}\psi^{\text{simp}(\partial_0\alpha)}(\alpha_2, \dots, \alpha_n)$. In the LHS, we have

$$(-1)^{n+p}(d_{\text{simp}}^p\mathcal{G}\psi)^\sigma(a) = (-1)^{n+p}c^{\sigma, p-1, A_q}(\mathcal{G}\psi)^{\partial_p\sigma}(u_p^*a)$$

$$= \sum_{\substack{\bar{m}' \in \text{Part}(p-1), \beta' \in S_{q,p-1} \\ \zeta' \in \text{Seqq}(\sigma, \bar{m}')}} (-1)^{n+p+\beta'+\zeta'} c^{\sigma,p-1,A_q} \psi^{\text{simp}(u_p^* a_{\beta'}^* \zeta')} (u_p^* a_{\beta'}^* \zeta').$$

Choose $\bar{m}' = (m_k, \dots, m_2) \in \text{Part}(p-1)$. There exist unique $\zeta' \in \text{Seqq}(\partial_p \sigma, \bar{m}')$ and $\beta' \in S_{q,p-1}$ such that $u_p^* a_{\beta'}^* \zeta' = (\alpha_2, \dots, \alpha_n)$. This implies

$$T(0, \bar{m}, \beta, \zeta) = (-1)^{n+p+\beta'+\zeta'} c^{\sigma,p-1,A_q} \psi^{\text{simp}(u_p^* a_{\beta'}^* \zeta')} (u_p^* a_{\beta'}^* \zeta').$$

- Consider the case $\underline{\alpha}_1 = 1_{\sigma[m_1]^*}$. We have $\text{simp}(\alpha_1) = (U_{p-m_1} \xrightarrow{u_p \cdots u_{p-m_1+1}} U_p)$, and $(\delta_0 \psi)^{\text{simp}(a_{\beta}^* \zeta)} (a_{\beta}^* \zeta) = c^{\sigma,p-m_1,A_q} \psi^{\text{simp}(\partial_0 \alpha)} (\alpha_2, \dots, \alpha_n)$. We have, in RHS, the terms

$$\begin{aligned} (d_{m_1} \mathcal{G}\psi)^{\sigma}(a) &= \sum_{r \in \mathcal{P}(\sigma[m_1]), \beta' \in S_{q,m_1-1}} (-1)^q (-1)^r (-1)^{\beta'} c^{\sigma,p-m_1,A_q} (\mathcal{G}\psi)^{L_{p-m_1} \sigma} (\beta'(a, r)) \\ &= \sum_{\substack{r \in \mathcal{P}(\sigma[m_1]), \beta' \in S_{q,m_1-1}, \beta'' \in S_{q+m_1-1,p-m_1} \\ \bar{m}' \in \text{Part}(p-m_1), \zeta' \in \text{Seqq}(L_{p-m_1} \sigma, \bar{m}')}} (-1)^{q+r+\beta'+\beta''+\zeta'} \times \\ &\quad c^{\sigma,p-m_1,A_q} \psi^{\text{simp}(\beta'(a,r)_{\beta''}^* \zeta')} (\beta'(a, r)_{\beta''}^* \zeta'). \end{aligned}$$

Let $\bar{m}' = (m_k, \dots, m_2) \in \text{Part}(p-m_1)$. We consider the element ζ' in $\text{Seqq}(L_{p-m_1} \sigma, \bar{m}')$ of the form $\zeta' = \gamma_1(\bar{r}^{m_2}, \dots, \bar{r}^{m_k})$ where $\gamma_1 \in \bar{S}_{\bar{m}'}$. Choose $r = r^{m_1}$, there exist unique $\gamma_1 \in \bar{S}_{\bar{m}'}$, $\beta' \in S_{q,m_1-1}$, $\beta'' \in S_{q+m_1-1,p-m_1} S$ such that $\beta'(a, r)_{\beta''}^* \zeta' = (\alpha_2, \dots, \alpha_n)$. Therefore

$$T(0, \bar{m}, \beta, \zeta) = (-1)^{q+r+\beta'+\beta''+\zeta'} c^{\sigma,p-m_1,A_q} \psi^{\text{simp}(\beta'(a,r)_{\beta''}^* \zeta')} (\beta'(a, r)_{\beta''}^* \zeta').$$

Step 2. We consider the term $T(n, \bar{m}, \beta, \zeta)$ in RHS. We have

$$(-1)^n (\delta_n \psi)^{\text{simp}(a_{\beta}^* \zeta)} (a_{\beta}^* \zeta) = (-1)^n \mu(\psi^{\text{simp}(\partial_0 \alpha)} (\alpha_1, \dots, \alpha_{n-1}), \alpha_n).$$

There are only three cases: $\underline{\alpha}_n = a_n$, $\underline{\alpha}_n = 1_{u_1^*}$ or $\underline{\alpha}_n = r_{m_i-1}^{m_i}$ where $r^{m_i} = (r_1^{m_i}, \dots, r_{m_i-1}^{m_i}) \in \mathcal{P}(\sigma[m_i])$.

- Consider the case $\underline{\alpha}_n = a_q$. Then $\text{simp}(\alpha_n) = (U_0 \xrightarrow{1} U_0)$. In LHS, we have

$$(-1)^q (d_{\text{Hoch}}^q \mathcal{G}\psi)^{\sigma}(a) = \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta' \in S_{q-1,p} \\ \zeta' \in \text{Seqq}(\sigma, \bar{m}')}} (-1)^{q+\beta'+\zeta'} \psi^{\text{simp}(\partial_q a_{\beta'}^* \zeta')} (\partial_q a_{\beta'}^* \zeta') \sigma^{\#}(a_q).$$

Choose $\bar{m}' = \bar{m} \in \text{Part}(p)$ and $\zeta' = \zeta \in \text{Seqq}(\sigma, \bar{m}')$. There exists a unique $\beta' \in S_{q-1,p}$ such that $(a_1, \dots, a_{q-1})_{\beta'}^* \zeta' = (\alpha_1, \dots, \alpha_{n-1})$. This implies

$$T(n, \bar{m}, \beta, \zeta) = (-1)^{q+\beta'+\zeta'} \psi^{\text{simp}(\partial_q a_{\beta'}^* \zeta')} (\partial_q a_{\beta'}^* \zeta') \sigma^{\#}(a_q).$$

- Consider the case $\underline{\alpha}_n = (1_{u_1^*})$. Then $m_k = 1$ and $\text{simp}(\alpha_n) = (U_0 \xrightarrow{1} U_1)$, so $\underline{\zeta} = (\underline{\eta}, 1_{u_1^*})$ for some $\underline{\eta} \in \text{Seqq}(\partial_0 \sigma, (m_{k-1}, \dots, m_1))$. We have

$$(\delta_n \psi)^{\text{simp}(a_{\beta}^* \zeta)} (a_{\beta}^* \zeta) = c^{\sigma,1,A_q} u_1^* \psi^{\text{simp}(\partial_n \alpha)} (\alpha_1, \dots, \alpha_{n-1}).$$

In LHS we have

$$(-1)^n (d_{\text{simp}}^0 \mathcal{G}\psi)^{\sigma}(a) = (-1)^n c^{\sigma,1,A_q} M^{u_1} (\mathcal{G}\psi)^{\partial_0 \sigma}(a)$$

$$= \sum_{\substack{\bar{m}' \in \text{Part}(p-1), \beta' \in S_{q,p-1} \\ \zeta' \in \text{Seqq}(\partial_0 \sigma, \bar{m}')}} (-1)^{n+\beta'+\zeta'} c^{\sigma,1,A_q} M^{u_1} \psi^{\text{simp}(a *_{\beta'} \zeta')} (a *_{\beta'} \zeta').$$

Take $\bar{m}' = (m_{k-1}, \dots, m_1) \in \text{Part}(p-1)$ and $\zeta' = \eta$, there exists a unique $\beta' \in S_{q,p-1}$ such that $(a_1, \dots, a_q) *_{\beta'} \zeta' = (\alpha_1, \dots, \alpha_{n-1})$. By computation,

$$\text{we obtain } T(n, \bar{m}, \beta, \zeta) = (-1)^{n+\beta'+\zeta'} c^{\sigma,1,A_q} M^{u_1} \psi^{\text{simp}(a *_{\beta'} \zeta')} (a *_{\beta'} \zeta').$$

- Consider the case $\underline{\alpha}_n = r_{m_{i-1}}^{m_i}$. Then $\alpha_n = \epsilon^{\sigma, j_0}(A_0)$ for some j_0 , $\text{simp}(\alpha_n) = (U_0 \xrightarrow{1} U_0)$. We have $(\delta_n \psi)^{\text{simp}(a *_{\beta} \zeta)} (a *_{\beta} \zeta) = \psi^{\text{simp}(\partial_n \alpha)} (\alpha_1, \dots, \alpha_{n-1}) \epsilon^{\sigma, j_0}(A_0)$. In LHS we have

$$(-1)^{n+j_0} (d_{\text{simp}}^{j_0} \mathcal{G}\psi)^{\sigma}(a) = \sum_{\substack{\bar{m}' \in \text{Part}(p-1), \beta' \in S_{q,p-1} \\ \zeta' \in \text{Seqq}(\partial_{j_0} \sigma, \bar{m}')}} (-1)^{n+j_0+\beta'+\zeta'} \psi^{\text{simp}(a *_{\beta'} \zeta')} (a *_{\beta'} \zeta') \epsilon^{\sigma, j_0}(A_0).$$

Take $\bar{m}' = (m_k, \dots, m_{i+1}, m_i - 1, m_{i-1}, \dots, m_1) \in \text{Part}(p-1)$. There exist unique $\zeta' \in \text{Seqq}(\partial_{j_0} \sigma, \bar{m}')$ and $\beta' \in S_{q,p-1}$ such that $(a_1, \dots, a_q) *_{\beta'} \zeta' = (\alpha_1, \dots, \alpha_{n-1})$. We get $T(n, \bar{m}, \beta, \zeta) = (-1)^{n+j_0+\beta'+\zeta'} \psi^{\text{simp}(a *_{\beta'} \zeta')} (a *_{\beta'} \zeta') \epsilon^{\sigma, j_0}(A_0)$.

Step 3. Considering the term $T(i, \bar{m}, \beta, \zeta)$ in RHS for $i = 1..(n-1)$, we have

$$(\delta_i \psi)^{\text{simp}(a *_{\beta} \zeta)} (a *_{\beta} \zeta) = \psi^{\text{simp}(\partial_i \alpha)} (\alpha_1, \dots, \alpha_{i-1}, \alpha_i \alpha_{i+1}, \alpha_{i+2}, \dots, \alpha_n).$$

Denote

$$\Gamma = \{1_{u_i^*}, 1_{\sigma[m_j]^*}, r_l^{m_t} \mid r^{m_t} = (r_1^{m_t}, \dots, r_{m_t-1}^{m_t}) \in \mathcal{P}(\sigma[m_t]), i, j, t, l \geq 1\}.$$

We consider the following case

- (i) Assume that $\{\underline{\alpha}_i, \underline{\alpha}_{i+1}\} \cap \{\tilde{a}_1, \dots, \tilde{a}_n\} \neq \emptyset$ then:

- (a) If $(\underline{\alpha}_i, \underline{\alpha}_{i+1}) = (a_j, a_{j+1})$ for some j , we look at $d_{\text{Hoch}}^j \mathcal{G}\psi$ in LHS,

$$(-1)^j (d_{\text{Hoch}}^j \mathcal{G}\psi)^{\sigma}(a) = \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta' \in S_{q-1,p} \\ \zeta' \in \text{Seqq}(\sigma, \bar{m}')}} (-1)^{j+\beta'+\zeta'} \psi^{\text{simp}(\partial_j a *_{\beta'} \zeta')} (\partial_j a *_{\beta'} \zeta').$$

Choose $\bar{m}' = \bar{m}$ and $\zeta' = \zeta$. There exists a unique $\beta' \in S_{q-1,p}$ such that $\partial_j a *_{\beta'} \zeta' = \partial_i \alpha$. Hence, $T(i, \bar{m}, \beta, \zeta) = (-1)^{j+\beta'+\zeta'} \psi^{\text{simp}(\partial_j a *_{\beta'} \zeta')} (\partial_j a *_{\beta'} \zeta')$.

- (b) If $\{\underline{\alpha}_i, \underline{\alpha}_{i+1}\} = \{a_j, b\}$ for some $b \in \Gamma$, there exists a unique $\beta' \in S_{q,p}$ such that $\beta'^{(0)}(a, \zeta) = (\underline{\alpha}_1, \dots, \underline{\alpha}_{i-1}, \underline{\alpha}_{i+1}, \underline{\alpha}_i, \underline{\alpha}_{i+2}, \dots, \underline{\alpha}_n)$. This implies $T(i, \bar{m}, \beta, \zeta) + T(i, \bar{m}, \beta', \zeta) = 0$.
- (ii) Assume that $\{\underline{\alpha}_i, \underline{\alpha}_{i+1}\} \subseteq \Gamma$, then :
- (a) If $\{\underline{\alpha}_i, \underline{\alpha}_{i+1}\} = \{r_j^{m_t}, 1_{u_l^*}\}$, then we repeat the argument in (i').
- (b) If $\{\underline{\alpha}_i, \underline{\alpha}_{i+1}\} = \{r_j^{m_t}, r_l^{m_s}\}$, then if $s \neq t$ we repeat the argument in (i'). Else $s = t$ so $(\underline{\alpha}_i, \underline{\alpha}_{i+1}) = (r_j^{m_t}, r_l^{m_t})$, then $l = j + 1$. In the formula (4.11) of ζ , we keep $\bar{r}^{m_{t'}}$ when $t' \neq t$ and replace $\bar{r}^{m_t} = (1_{\sigma[m_t]^*}, r^{m_t})$ by $(1_{\sigma[m_t]^*}, \text{flip}(r^{m_t}, j))$ to obtain the new element $\eta \in \text{Seqq}(\sigma, \bar{m})$. Then $(-1)^\eta = -(-1)^\zeta$ and by (3.17) we get $\partial_i(a *_{\beta} \eta) = \partial_i \alpha$. This implies $T(i, \bar{m}, \beta, \eta) + T(i, \bar{m}, \beta, \zeta) = 0$.

- (c) If $\{\underline{\alpha}_i, \underline{\alpha}_{i+1}\} = \{1_{\sigma[m_j]}, r_v^{m_t}\}$ where $r^{m_j} = (r_1^{m_j}, \dots, r_{m_j-1}^{m_j}) \in \mathcal{P}(\sigma[m_j])$, then if $j \neq t$ we again repeat the argument in (iⁿ). If $j = t$ then $(\underline{\alpha}_i, \underline{\alpha}_{i+1}) = (1_{\sigma[m_j]}, r_1^{m_j})$. Assume that $r_1^{m_j} = c^{\sigma[m_j], l}$ for some l . Let $\Delta = m_j - l, \Delta' = l$. We decompose $\sigma[m_j] = \sigma[\Delta'] \sqcup \sigma[\Delta]$ as concatenation of $\sigma[\Delta']$ and $\sigma[\Delta]$. By Lemma (3.8) there exist paths $r^\Delta \in \mathcal{P}(\sigma[\Delta]), r^{\Delta'} \in \mathcal{P}(\sigma[\Delta'])$ and $\beta_0 \in S_{\Delta-1, \Delta'-1}$ such that $(r_1^{m_j}, r^\Delta \underset{\beta_0}{*} r^{\Delta'}) = r^{m_j}$. Choose the new partition $\bar{m}' = (m_k, \dots, m_{j+1}, \Delta', \Delta, m_{j-1}, \dots, m_1) \in \text{Part}(p)$. There exists a unique conditioned shuffle permutation $\gamma \in \bar{S}_{\bar{m}}$ such that
- $$\begin{aligned} & \gamma^{(0)}(\bar{r}^{m_1}, \dots, \bar{r}^{m_{i-1}}, \bar{r}^\Delta, \bar{r}^{\Delta'}, \bar{r}^{m_{i+1}}, \dots, \bar{r}^{m_k}) \\ &= (\underline{\alpha}_1, \dots, \underline{\alpha}_{i-1}, 1_{\sigma[\Delta]}, 1_{\sigma[\Delta']}, \underline{\alpha}_{i+1}, \dots, \underline{\alpha}_n). \end{aligned}$$
- Let $\eta = \gamma^{(0)}(\bar{r}^{m_1}, \dots, \bar{r}^{m_{i-1}}, \bar{r}^\Delta, \bar{r}^{\Delta'}, \bar{r}^{m_{i+1}}, \dots, \bar{r}^{m_k}) \in \text{Seqq}(\sigma, \bar{m}')$. Since $\partial_i(a * \eta) = \partial_i \alpha$, we get $T(i, \bar{m}, \beta, \zeta) + T(i, \bar{m}', \beta, \eta) = 0$.

□

4.3. \mathcal{F} and \mathcal{G} are quasi-inverse. In this section we construct homotopy maps

$$\{T_{n+1} : \mathbf{C}_{\mathcal{U}}^{n+1}(\tilde{\mathcal{A}}, \tilde{M}) \longrightarrow \mathbf{C}_{\mathcal{U}}^n(\tilde{\mathcal{A}}, \tilde{M})\}$$

to show that $\mathcal{F}\mathcal{G} \sim 1$, then we prove directly that $\mathcal{G}\mathcal{F}(\phi) = \phi$ for any normalized reduced cochain ϕ . Hence we conclude that both \mathcal{F} and \mathcal{G} are quasi-isomorphisms, in particular, we have

$$HH_{\text{GS}}^n(\mathcal{A}, M) = H^n \mathbf{C}_{\text{GS}}^\bullet(\mathcal{A}, M) \cong H^n(\mathbf{C}_{\mathcal{U}}^\bullet(\tilde{\mathcal{A}}, \tilde{M})) = HH_{\mathcal{U}}^n(\tilde{\mathcal{A}}, \tilde{M}).$$

For each n -simplex $\sigma = (u_1, \dots, u_n)$ as in (3.8), let $A = (A_i)_{i=0}^n$ where $A_i \in \tilde{\mathcal{A}}(U_i)$. Denote

$$\tilde{\mathcal{A}}_{\sigma, A} = \tilde{\mathcal{A}}_{u_n}(A_{n-1}, A_n) \otimes \cdots \otimes \tilde{\mathcal{A}}_{u_1}(A_0, A_1).$$

Let

$$\Lambda = \{x \in \tilde{\mathcal{A}}_{\sigma, A} \mid \sigma \in \mathcal{N}(\mathcal{U}), A_i \in \mathcal{A}(\sigma(i))\}.$$

Denote by $\langle \Lambda \rangle$ the free abelian group generated by Λ . Given $\Psi \in \mathbf{C}_{\mathcal{U}}^n(\tilde{\mathcal{A}}, \tilde{M})$ and $x = \sum_{\sigma, A} x_{\sigma, A} \in \langle \Lambda \rangle$ where $x_{\sigma, A} \in \tilde{\mathcal{A}}_{\sigma, A}$, then we set

$$\Psi(x) = \sum_{\sigma, A} \Psi(x_{\sigma, A})$$

in which $\Psi(x_{\sigma, A}) = 0$ if $\sigma \notin \mathcal{N}_n(\mathcal{U})$.

Let $\sigma = (u_1, \dots, u_n)$ and $\gamma = (v_1, \dots, v_m)$ be simplices as in (3.8). Let $A = (A_i)_{i=0}^n$ where $A_i \in \tilde{\mathcal{A}}(U_i)$ and $B = (B_i)_{i=0}^m$ where $B_i \in \tilde{\mathcal{A}}(V_i)$. Given $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n) \in \tilde{\mathcal{A}}_{\sigma, A}$ and $\tilde{b} = (\tilde{b}_1, \dots, \tilde{b}_m) \in \tilde{\mathcal{A}}_{\gamma, B}$ as in (4.2), we have

$$\text{simp}(\tilde{a}) = \sigma; \text{simp}(\tilde{b}) = \gamma.$$

Assume $A_n = B_0$ and $U_n = V_0$, we define the concatenation

$$\tilde{b} \sqcup \tilde{a} = (\tilde{b}_1, \dots, \tilde{b}_m, \tilde{a}_1, \dots, \tilde{a}_n);$$

$$\text{simp}(\tilde{b} \sqcup \tilde{a}) = \text{simp}(\tilde{b}) \sqcup \text{simp}(\tilde{a}) = \gamma \sqcup \sigma.$$

We have $\text{simp}(\tilde{a}_i) = (U_{n-i} \xrightarrow{u_{n-i+1}} U_{n-i+1})$, and so

$$\text{simp}(\tilde{a}) = \sigma = \text{simp}(\tilde{a}_1) \sqcup \cdots \sqcup \text{simp}(\tilde{a}_n).$$

We use the following notations

$$\partial_0(\tilde{a}) = (\tilde{a}_2, \dots, \tilde{a}_n), \text{simp}(\partial_0(\tilde{a})) = \partial_n \sigma;$$

$$\partial_i(\tilde{a}) = (\tilde{a}_1, \dots, \tilde{a}_{i-1}, \mu(\tilde{a}_i, \tilde{a}_{i+1}), \tilde{a}_{i+2}, \dots, \tilde{a}_n), \text{simp}(\partial_i(\tilde{a})) = \partial_{n-i} \sigma;$$

$$\begin{aligned}\partial_n(\tilde{a}) &= (\tilde{a}_1, \dots, \tilde{a}_{n-1}), \quad \text{simp}(\partial_n(\tilde{a})) = \partial_0\sigma; \\ R_p\tilde{a} &= (\tilde{a}_1, \dots, \tilde{a}_{n-p}); \\ \bar{a}_{n+1-p, \dots, n} &= \tilde{a}_{n+1-p, \dots, n}, \quad \text{simp}(\bar{a}_{n+1-p, \dots, n}) = (U_0 \xrightarrow{1} U_0).\end{aligned}$$

In the abelian group $\langle \Lambda \rangle$, we put

$$\begin{aligned}\omega_{n,p}(\sigma, \tilde{a}) &= \sum_{\substack{\bar{m} \in \text{Part}(n-p) \\ \xi \in \text{Seq}(R_p\sigma, R_p\tilde{a}, \bar{m})}} \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta \in S_{n-p,p} \\ \zeta \in \text{Seq}(L_p\sigma, \bar{m}')}} (-1)^{\bar{m}+\xi+\zeta+\beta} \xi *_{\beta} \zeta \sqcup \bar{a}_{n+1-p, \dots, n}; \\ \omega_n(\sigma, \tilde{a}) &= \sum_{p=1}^n \omega_{n,p}(\sigma, \tilde{a}); \\ \Delta_n(\sigma, \tilde{a}) &= \sum_{\substack{\bar{m} \in \text{Part}(n) \\ \xi \in \text{Seq}(\sigma, \bar{m})}} (-1)^{\bar{m}+\xi} \xi - (\tilde{a}_1, \dots, \tilde{a}_n).\end{aligned}$$

By induction, we define

$$(4.12) \quad \Omega_n(\sigma, \tilde{a}) = (-1)^{n+1} \omega_n(\sigma, \tilde{a}) + \Omega_{n-1}(\partial_0\sigma, \partial_n\tilde{a}) \sqcup \tilde{a}_n, \quad \text{for } n \geq 2,$$

if $n = 1$ then (σ, \tilde{a}) is represented as

$$(\sigma, \tilde{a}) = \left(\begin{array}{ccc} A_0 & \xrightarrow{\tilde{a}} & A_1 \\ U_0 & \xrightarrow[u_1]{} & U_1 \end{array} \right)$$

and we set

$$\Omega_1(\sigma, \tilde{a}) = \left(\begin{array}{ccccc} A_0 & \xrightarrow{\tilde{a}} & u_1^* A_1 & \xrightarrow{1_{u_1^* A_1}} & A_1 \\ U_0 & \xrightarrow[1_{U_0}]{} & U_0 & \xrightarrow[u_1]{} & U_1 \end{array} \right).$$

Now we define the homotopy maps $\{T_{n+1} : \mathbf{C}_{\mathcal{U}}^{n+1}(\tilde{\mathcal{A}}, \tilde{M}) \longrightarrow \mathbf{C}_{\mathcal{U}}^n(\tilde{\mathcal{A}}, \tilde{M})\}$ as follows:

$$\begin{aligned}T_1 &= 0, \\ (T_{n+1}\Psi)^\sigma(\tilde{a}) &= \Psi(\Omega_n(\sigma, \tilde{a})), \quad n \geq 1.\end{aligned}$$

From now on, for simplicity we write $\Omega_n(\tilde{a})$ and $\omega_n(\tilde{a})$ instead of $\Omega_n(\sigma, \tilde{a})$ and $\omega_n(\sigma, \tilde{a})$.

Theorem 4.6. *The maps T_n constitute a homotopy $T : \mathcal{FG} \longrightarrow 1$. More precisely, given a cochain Ψ in $\mathbf{C}_{\mathcal{U}}^n(\tilde{\mathcal{A}}, \tilde{M})$, we have*

$$(4.13) \quad \mathcal{FG}(\Psi) - \Psi = \delta T_n \Psi + T_{n+1} \delta \Psi.$$

Proof. We have

$$\begin{aligned}(\mathcal{FG}\Psi)^\sigma(\tilde{a}) &= (\mathcal{F}_0\mathcal{G}\Psi)^\sigma(\tilde{a}) + \sum_{p=1}^n (\mathcal{F}_p\mathcal{G}\Psi)^\sigma(\tilde{a}) \\ &= (\mathcal{F}_0\mathcal{G}\Psi)^\sigma(\tilde{a}) + \sum_{\substack{\bar{m} \in \text{Part}(n-p) \\ \xi \in \text{Seq}(R_p\sigma, R_p\tilde{a}, \bar{m})}} \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta \in S_{n-p,p} \\ \zeta \in \text{Seq}(L_p\sigma, \bar{m}')}} (-1)^{\xi+\zeta+\beta} \Psi(\xi *_{\beta} \zeta) \tilde{a}_{n+1-p, \dots, n} \\ &= (\mathcal{F}_0\mathcal{G}\Psi)^\sigma(\tilde{a}) + \delta_{n+1} \Psi(\omega_n(\sigma, \tilde{a})).\end{aligned}$$

Moreover, we have

$$\begin{aligned}(-1)^{n+1} (T_{n+1} \delta_{n+1} \Psi)^\sigma(\tilde{a}) &= \delta_{n+1} \Psi(\omega_n(\sigma, \tilde{a})) + (-1)^{n+1} \delta_{n+1} \Psi(\Omega_{n-1}(\partial_0\sigma, \partial_n\tilde{a}) \sqcup \tilde{a}_n) \\ &= \delta_{n+1} \Psi(\omega_n(\sigma, \tilde{a})) + (-1)^{n+1} \Psi(\Omega_{n-1}(\partial_0\sigma, \partial_n\tilde{a})) \tilde{a}_n\end{aligned}$$

and

$$(-1)^n(\delta_n T_n \Psi)^\sigma(\tilde{a}) = (-1)^n(T_n \Psi)^{\partial_0 \sigma}(\partial_n \tilde{a})\tilde{a}_n = (-1)^n \Psi(\Omega_{n-1}(\partial_0 \sigma, \partial_n \tilde{a}))\tilde{a}_n.$$

This implies

$$(-1)^{n+1}(T_{n+1} \delta_{n+1} \Psi)^\sigma(\tilde{a}) + (-1)^n(\delta_n T_n \Psi)^\sigma(\tilde{a}) = \delta_{n+1} \Psi(\omega_n(\sigma, \tilde{a})).$$

So the equation (4.13) is equivalent to the equation

$$(\mathcal{F}_0 \mathcal{G} \Psi)^\sigma(\tilde{a}) - \Psi^\sigma(\tilde{a}) = \sum_{i=0}^n (-1)^i (T_{n+1} \delta_i \Psi)^\sigma(\tilde{a}) + \sum_{i=0}^{n-1} (-1)^i (\delta_i T_n \Psi)^\sigma(\tilde{a}).$$

This equation holds true due to Lemma 4.7 right below. \square

Lemma 4.7. *Let $\sigma = (u_1, \dots, u_n)$ be a simplex and $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_n)$ as in (4.2), then the following equation holds true*

$$(4.14) \sum_{i=0}^n (-1)^i \partial_i \Omega_n(\tilde{a}_1, \dots, \tilde{a}_n) + \sum_{i=0}^{n-1} (-1)^i \Omega_{n-1}(\partial_i(\tilde{a}_1, \dots, \tilde{a}_n)) = \Delta_n(\tilde{a}).$$

Proof. We prove this lemma by induction on n . The equation (4.14) is equivalent to

$$\sum_{i=0}^{n-1} (-1)^i \Omega_{n-1}(\partial_i \tilde{a}) = \Delta_n(\tilde{a}) - \sum_{i=0}^n (-1)^{i+n+1} \partial_i \omega_n(\tilde{a}) - \sum_{i=0}^n (-1)^i \partial_i (\Omega_{n-1}(\tilde{a}_1, \dots, \tilde{a}_{n-1}) \sqcup \tilde{a}_n).$$

Assume that the equation (4.14) holds true for n . We prove it holds true for $n+1$.

Assume $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_{n+1})$. Let

$$B = \sum_{i=0}^{n+1} (-1)^i \partial_i \Omega_{n+1}(\tilde{a}_1, \dots, \tilde{a}_{n+1}), \text{ and } C = \sum_{i=0}^n (-1)^i \Omega_n(\partial_i(\tilde{a}_1, \dots, \tilde{a}_{n+1})).$$

We need to prove

$$(4.15) \quad B + C = \Delta(\tilde{a}_1, \dots, \tilde{a}_{n+1}).$$

By definition, we have

$$\begin{aligned} B &= \sum_{i=0}^{n+1} (-1)^{i+n+2} \partial_i \omega_{n+1}(\tilde{a}_1, \dots, \tilde{a}_{n+1}) + \sum_{i=0}^{n+1} (-1)^i \partial_i (\Omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}) \\ &= \sum_{i=0}^{n+1} (-1)^{i+n+2} \partial_i \omega_{n+1}(\tilde{a}_1, \dots, \tilde{a}_{n+1}) + B_1 + B_2 \end{aligned}$$

where

$$\begin{aligned} B_1 &= \sum_{i=0}^{n+1} (-1)^{i+n+1} \partial_i (\omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}) \\ B_2 &= \sum_{i=0}^{n+1} (-1)^i \partial_i (\Omega_{n-1}(\tilde{a}_1, \dots, \tilde{a}_{n-1}) \sqcup \tilde{a}_n \sqcup \tilde{a}_{n+1}). \end{aligned}$$

We also have

$$\begin{aligned} C &= \sum_{i=0}^n (-1)^{i+n+1} \omega_n(\partial_i(\tilde{a}_1, \dots, \tilde{a}_{n+1})) + \sum_{i=0}^{n-1} \Omega_{n-1}(\partial_i(\tilde{a}_1, \dots, \tilde{a}_n)) \sqcup \tilde{a}_{n+1} \\ &\quad + (-1)^n \Omega_{n-1}(\tilde{a}_1, \dots, \tilde{a}_{n-1}) \sqcup \tilde{a}_{n,n+1}. \end{aligned}$$

By induction hypothesis, we have

$$\sum_{i=0}^{n-1} (-1)^i \Omega_{n-1}(\partial_i(\tilde{a}_1, \dots, \tilde{a}_n)) \sqcup \tilde{a}_{n+1}$$

$$\begin{aligned}
&= \Delta_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1} - \sum_{i=0}^n (-1)^{i+n+1} \partial_i \omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1} \\
&\quad - \sum_{i=0}^n (-1)^i \partial_i (\Omega_{n-1}(\tilde{a}_1, \dots, \tilde{a}_{n-1}) \sqcup \tilde{a}_n) \sqcup \tilde{a}_{n+1} \\
&= \Delta_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1} - (B_1 - \partial_{n+1}(\omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1})) \\
&\quad - (B_2 - (-1)^{n+1} \Omega_{n-1}(\tilde{a}_1, \dots, \tilde{a}_{n-1}) \sqcup \tilde{a}_{n,n+1}).
\end{aligned}$$

This implies

$$\begin{aligned}
B + C &= \sum_{i=0}^{n+1} (-1)^{i+n+2} \partial_i \omega_{n+1}(\tilde{a}_1, \dots, \tilde{a}_{n+1}) + \sum_{i=0}^n (-1)^{i+n+1} \omega_n(\partial_i(\tilde{a}_1, \dots, \tilde{a}_{n+1})) \\
&\quad + \partial_{n+1}(\omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}) + \Delta_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}.
\end{aligned}$$

Thus, by Lemma (4.8), we obtain the equation (4.15). \square

Lemma 4.8. *Let $\sigma = (u_1, \dots, u_{n+1})$ be a simplex and $\tilde{a} = (\tilde{a}_1, \dots, \tilde{a}_{n+1})$ as in (4.2), then we have*

$$\begin{aligned}
&\sum_{i=0}^{n+1} (-1)^{i+n+2} \partial_i \omega_{n+1}(\tilde{a}_1, \dots, \tilde{a}_{n+1}) + \sum_{i=0}^n (-1)^{i+n+1} \omega_n(\partial_i(\tilde{a}_1, \dots, \tilde{a}_{n+1})) \\
&\quad + \partial_{n+1}(\omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}) + \Delta_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1} - \Delta(\tilde{a}_1, \dots, \tilde{a}_{n+1}) = 0.
\end{aligned}$$

Proof. We denote

$$\begin{aligned}
B &= \sum_{i=0}^{n+1} (-1)^{i+n+2} \partial_i \omega_{n+1}(\tilde{a}_1, \dots, \tilde{a}_{n+1}); \quad C = \sum_{i=0}^n (-1)^{i+n+1} \omega_n(\partial_i(\tilde{a}_1, \dots, \tilde{a}_{n+1})); \\
D &= \partial_{n+1}(\omega_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}); \quad E = \Delta_n(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1} - \Delta(\tilde{a}_1, \dots, \tilde{a}_{n+1}).
\end{aligned}$$

We prove that each of the terms appearing in the expansion of B is cancelled out against a unique term in C, D , or E , and vice-versa. The cancellation is as follows:

$$\begin{array}{ccc}
B & \longleftrightarrow & C \\
\updownarrow & \searrow & \\
D & & E
\end{array}$$

We write

$$\begin{aligned}
B_p &= \sum_{i=0}^{n+1} (-1)^{i+n+2} \partial_i \omega_{n+1,p}(\tilde{a}) \\
&= \sum_{i=0}^{n+1} \sum_{\substack{\bar{m} \in \text{Part}(n+1-p) \\ \xi \in \text{Seq}(R_p \sigma, R_p \tilde{a}, \bar{m})}} \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta \in S_{n+1-p,p} \\ \zeta \in \text{Seq}(L_p \sigma, \bar{m}')}} (-1)^{i+n+2} (-1)^{\bar{m} + \xi + \zeta + \beta} \partial_i B_p(\tilde{a}, \xi, \zeta, \beta)
\end{aligned}$$

where $B_p(\tilde{a}, \xi, \zeta, \beta) = \xi *_{\beta} \zeta \sqcup \bar{a}_{n+2-p, \dots, n+1}$, and write

$$\begin{aligned}
C_p &= \sum_{i=0}^n (-1)^{i+n+1} \omega_{n,p}(\partial_i \tilde{a}) \\
&= \sum_{i=0}^n \sum_{\substack{\bar{m} \in \text{Part}(n-p) \\ \xi \in \text{Seq}(R_p \partial_{n+1-i} \sigma, R_p \partial_i \tilde{a}, \bar{m})}} \sum_{\substack{\bar{m}' \in \text{Part}(p), \beta \in S_{n-p,p} \\ \zeta \in \text{Seq}(L_p \partial_{n+1-i} \sigma, \bar{m}')}} (-1)^{i+n+1} (-1)^{\bar{m} + \xi + \zeta + \beta} C_p(\partial_i \tilde{a}, \xi, \zeta, \beta)
\end{aligned}$$

where $C_p(\partial_i \tilde{a}, \xi, \zeta, \beta) = \xi *_{\beta} \zeta \sqcup \overline{(\partial_i \tilde{a})}_{n+1-p, \dots, n}$.

We also write

$$E_0 = -\Delta(\tilde{a}_1, \dots, \tilde{a}_{n+1}) = \sum_{\substack{\tilde{m} \in \text{Part}(n+1) \\ \xi \in \text{Seq}(\sigma, \tilde{m})}} (-1)^{\tilde{m} + \xi + 1} \xi;$$

$$E_1 = \Delta(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1} = \sum_{\substack{\tilde{m} \in \text{Part}(n) \\ \xi \in \text{Seq}(\partial_0 \sigma, \tilde{m})}} (-1)^{\tilde{m} + \xi} \xi \sqcup \tilde{a}_{n+1}.$$

and

$$D_p = \partial_{n+1} \omega_{n,p}(\tilde{a}_1, \dots, \tilde{a}_n) \sqcup \tilde{a}_{n+1}$$

$$= \sum_{\substack{\tilde{m} \in \text{Part}(n-p) \\ \xi \in \text{Seq}(R_p \partial_0 \sigma, R_p \partial_{n+1} \tilde{a}, \tilde{m})}} \sum_{\substack{\tilde{m}' \in \text{Part}(p), \beta \in S_{n-p,p} \\ \zeta \in \text{Seq}(L_p \partial_0 \sigma, \tilde{m}')}} (-1)^{\tilde{m} + \xi + \zeta + \beta} \partial_{n+1} D_p((\tilde{a}_1, \dots, \tilde{a}_n), \xi, \zeta, \beta)$$

where $D_p(\partial_{n+1} \tilde{a}, \xi, \zeta, \beta) = \xi *_{\beta} \zeta \sqcup \tilde{a}_{n+1-p, \dots, n} \sqcup \tilde{a}_{n+1}$.

Assume that $\tilde{m} = (m_l, \dots, m_1) \in \text{Part}(n+1-p)$ and $\tilde{m}' = (m'_k, \dots, m'_1) \in \text{Part}(p)$. Let $\xi \in \text{Seq}(R_p \sigma, \tilde{m})$, $\zeta \in \text{Seq}(L_p \sigma, \tilde{m}')$ and $\beta \in S_{n+1-p,p}$. We denote

$$B_p(\tilde{a}, \xi, \zeta, \beta) = (b_1, \dots, b_{n+2}),$$

and denote $(\underline{b}_1, \dots, \underline{b}_{n+2})$ the formal sequence of $B_p(\tilde{a}, \xi, \zeta, \beta)$.

Step 1. Consider the case $i = 0$, then $\partial_0(B_p(\tilde{a}, \xi, \zeta, \beta)) = (b_2, \dots, b_{n+2})$. There are only the following three cases:

$$b_1 = \begin{cases} \tilde{a}_1; \\ r_1^{m_t} & \text{where } r^{m_t} = (r_1^{m_t}, \dots, r_{m_t-1}^{m_t}) \in \mathcal{P}(\sigma[m_t]); \\ 1_{(L_p \sigma)[m'_1]^*}. \end{cases}$$

- (i) Assume $\underline{b}_1 = \tilde{a}_1$. Then $m_1 = 1$, and we choose $\tilde{m} = (m_l, \dots, m_2) \in \text{Part}(n-p)$. There exists a unique element $\xi' \in \text{Seq}(R_p(\partial_{n+1} \sigma), \tilde{m})$ such that $b_1 \sqcup \xi' = \xi$. There exists a unique $\beta' \in S_{n-p,p}$ such that

$$(b_1 \sqcup \xi') *_{\beta'} \zeta = \xi *_{\beta} \zeta.$$

We get

$$(-1)^{n+2} (-1)^{\tilde{m} + \xi + \zeta + \beta} \partial_0 B_p(\tilde{a}, \xi, \zeta, \beta) + (-1)^{n+1} (-1)^{\tilde{m} + \xi' + \zeta + \beta'} C_p(\partial_0 \tilde{a}, \xi', \zeta, \beta') = 0.$$

- (ii) Assume $\underline{b}_1 = r_1^{m_t}$ for some t , where $r^{m_t} = (r_1^{m_t}, \dots, r_{m_t-1}^{m_t}) \in \mathcal{P}(\sigma[m_t])$. Then $r_1^{m_t} = c^{(R_p \sigma)[m_t], j}$ for some j . Set $\Delta = j$, $\Delta' = m_t - j$. Using the analogous argument as in Case 2 of Step 2 in the proof of Proposition 4.2, considering the partition

$$\tilde{m} = (m_l, \dots, m_{t+1}, \Delta, \Delta', m_{t-1}, \dots, m_1) \in \text{Part}(n+1-p)$$

we find unique $\xi' \in \text{Seq}(R_p \sigma, \tilde{m})$ and $1 \leq j_0 \leq n+1$ such that

$$(-1)^{n+2} (-1)^{\tilde{m} + \xi + \zeta + \beta} \partial_0 B_p(\tilde{a}, \xi, \zeta, \beta) + (-1)^{j_0 + n+2} (-1)^{\tilde{m} + \xi' + \zeta + \beta} \partial_{j_0} B_p(\tilde{a}, \xi, \zeta, \beta) = 0.$$

- (iii) Assume $\underline{b}_1 = 1_{(L_p \sigma)[m'_1]^*}$.

If $m'_1 < p$, choose $\tilde{m}' = (m'_k, \dots, m'_2) \in \text{Part}(p-m'_1)$ and $\tilde{m} = (m'_1, m_l, \dots, m_1) \in \text{Part}(n+1-p+m'_1)$. There exists unique $\xi' \in \text{Seq}(R_{p-m'_1} \sigma, \tilde{m})$, $\zeta' \in \text{Seq}(L_{p-m'_1} \sigma, \tilde{m}')$ and $\beta' \in S_{n+1-p+m'_1, p-m'_1}$ such that

$$\xi' *_{\beta'} \zeta' \sqcup \tilde{a}_{n+2-p+m'_1, \dots, n+1} = (b_2, \dots, b_{n+1}) \sqcup \tilde{a}_{n+2-p, \dots, n+1-p+m'_1} \sqcup \tilde{a}_{n+2-p+m'_1, \dots, n+1}.$$

So we get $\partial_{n+1}(\xi' *_{\beta'} \zeta' \sqcup \tilde{a}_{n+2-p+m'_1, \dots, n+1}) = (b_2, \dots, b_{n+2})$. Then, we obtain

$$(-1)^{n+2} (-1)^{\tilde{m} + \xi + \zeta + \beta} \partial_0 B_p(\tilde{a}, \xi, \zeta, \beta) - (-1)^{\tilde{m} + \xi' + \zeta' + \beta'} \partial_{n+1} B_{p-m'_1}(\tilde{a}, \xi', \zeta', \beta') = 0.$$

If $m'_1 = p$, then $\bar{m}' = (m'_1) \in \text{Part}(p)$ and thus $\text{simp}(b_i) = (U_0 \xrightarrow{1} U_0)$ for $i = 2, \dots, (n+1)$. Let $\bar{m} = (m'_1, m_l, \dots, m_1) \in \text{Part}(n+1)$. It is seen that $\xi' = (b_2, \dots, b_{n+1}) \in \text{Seq}(\sigma, \bar{m})$ and

$$(-1)^{n+2}(-1)^{\bar{m}+\xi+\zeta+\beta} \partial_0 B_p(\bar{a}, \xi, \zeta, \beta) = (-1)^{\bar{m}+\xi'} \xi'.$$

Hence the term $(-1)^{n+2}(-1)^{\bar{m}+\xi+\zeta+\beta} \partial_0 B_p(\bar{a}, \xi, \zeta, \beta)$ is killed by the term $(-1)^{1+\bar{m}+\xi'} \xi'$ in E_0 . In this way, when p runs through $\{1, \dots, n+1\}$, every term in E_0 is eliminated except the term $(-\bar{a}_1, \dots, \bar{a}_{n+1})$ which is eliminated by the term $(\bar{a}_1, \dots, \bar{a}_n) \sqcup \bar{a}_{n+1}$ in E_1 .

Step 2. Consider the case $1 \leq i \leq n$. We write $\xi = (\xi_1, \dots, \xi_{n+1-p})$ and $\zeta = (\zeta_1, \dots, \zeta_p)$. We have

$$\partial_i B_p(\bar{a}, \xi, \zeta, \beta) = (b_1, \dots, b_{i-1}, \mu(b_i, b_{i+1}), b_{i+2}, \dots, b_{n+2}).$$

There are only the following three cases:

$$\left[\begin{array}{l} \{b_i, b_{i+1}\} = \{\xi_j, \zeta_{j'}\} \text{ for some } j, j'; \\ \{b_i, b_{i+1}\} \subseteq \{\xi_1, \dots, \xi_{n+1-p}\}; \\ \{b_i, b_{i+1}\} \subseteq \{\zeta_1, \dots, \zeta_p\}. \end{array} \right.$$

- Assume $\{b_i, b_{i+1}\} = \{\xi_j, \zeta_{j'}\}$. Choose $\beta' = (i, i+1) \circ \beta$ then

$$(-1)^{i+n+2}(-1)^{\bar{m}+\xi+\zeta+\beta} \partial_i B_p(\bar{a}, \xi, \zeta, \beta) + (-1)^{i+n+2}(-1)^{\bar{m}+\xi+\zeta+\beta'} \partial_i B_p(\bar{a}, \xi, \zeta, \beta') = 0.$$

- Assume $\{b_i, b_{i+1}\} \subseteq \{\xi_1, \dots, \xi_{n+1-p}\}$. We repeat the arguments of Step 2 in the Proposition 4.2.
- Assume $\{b_i, b_{i+1}\} \subseteq \{\zeta_1, \dots, \zeta_p\}$. We repeat the arguments of Step 3 in the Proposition 4.5.

Step 3. Consider the case $i = n+1$. We have

$$\begin{aligned} \partial_{n+1} B_p(\bar{a}, \xi, \zeta, \beta) &= \partial_{n+1}((b_1, \dots, b_{n+1}) \sqcup \bar{a}_{n+2-p, \dots, n+1}) \\ &= (b_1, \dots, b_n, \mu(b_{n+1}, \bar{a}_{n+2-p, \dots, n+1})). \end{aligned}$$

There are only the following three cases for \underline{b}_{n+1} :

$$\underline{b}_{n+1} = \begin{cases} \bar{a}[m_l]; \\ r_{m'_t-1}^{m'_t} & \text{where } r^{m'_t} = (r_1^{m'_t}, \dots, r_{m'_t-1}^{m'_t}) \in \mathcal{P}(L_p \sigma[m'_t]); \\ 1_{(L_p \sigma)[m'_k]^*} & \text{where } m'_k = 1 \text{ and } \bar{m}' = (1, m'_{k-1}, \dots, m'_1) \in \text{Part}(p). \end{cases}$$

- Assume $\underline{b}_{n+1} = \bar{a}[m_l]$. We apply the argument in (iii) of Step 1, then every term of this form is killed.
- Assume $\underline{b}_{n+1} = r_{m'_t-1}^{m'_t}$. We assume that $r_{m'_t-1}^{m'_t} = \epsilon^{L_p \sigma[m'_t].j}$ for some j . Then

$$\mu(r_1^{m'_t}, \bar{a}_{n+2-p, \dots, n+1}) = \overline{(\partial_j \bar{a})}_{n+2-p, \dots, n+1}.$$

In C we consider terms $C_p(\partial_j \bar{a}, \xi', \zeta', \beta')$. There exists unique (ξ', ζ', β') such that

$$-(-1)^{\bar{m}+\xi+\zeta+\beta} \partial_{n+1} B_p(\bar{a}, \xi, \zeta, \beta) + (-1)^{n+1+j}(-1)^{\bar{m}+\xi'+\zeta'+\beta'} C_p(\partial_j \bar{a}, \xi', \zeta', \beta') = 0$$

Combining with Step 2 and (i) in Step 1, we see that every term in C is killed.

- Assume that $\underline{b}_{n+1} = 1_{(L_p \sigma)[m'_k]^*}$ where $\bar{m}' = (1, m'_{k-1}, \dots, m'_1) \in \text{Part}(p)$. Thus we have $\underline{b}_{n+1} = 1_{u_1^*}$ and $\text{simp}(b_{n+1}) = (U_0 \xrightarrow{u_1} U_1)$. If $p = 1$, then we have $\bar{m}' = (1)$, $\zeta = 1_{L_p \sigma[1]^*} = b_{n+1}$, $\beta = 1$ and $\text{simp}(b_i) = (U_1 \xrightarrow{1} U_1)$ for $i \leq n$. So $\partial_{n+1} B_1(\bar{a}, \xi, \zeta, \beta) = \xi \sqcup \bar{a}_{n+1}$. We have $(-1)^{\bar{m}+\xi} \xi \sqcup \bar{a}_{n+1}$ is a term in E_1 and

$$-(-1)^{\bar{m}+\xi+\zeta+\beta} \partial_{n+1} B_1(\bar{a}, \xi, \zeta, \beta) + (-1)^{\bar{m}+\xi} \xi \sqcup \bar{a}_{n+1} = 0.$$

So we see that every term in E_1 is killed.

If $p > 1$, recall that $\tilde{m} = (m_l, \dots, m_1)$. We show that the term

$$-(-1)^{\tilde{m}+\xi+\zeta+\beta} \partial_{n+1} B_p(\tilde{a}, \xi, \zeta, \beta)$$

is killed by a term in D . Thus in the expression of D_p , we choose $\tilde{m}' = (m'_{k-1}, \dots, m'_1) \in \text{Part}(p-1)$, $\tilde{m} = \tilde{m} \in \text{Part}(n+1-p)$, $\xi' = \xi$ and $\zeta' = (\zeta_1, \dots, \zeta_{p-1})$. There exists a unique $\beta' \in S_{n+1-p, p-1}$ such that

$$-(-1)^{\tilde{m}+\xi+\zeta+\beta} \partial_{n+1} B_p(\tilde{a}, \xi, \zeta, \beta) + (-1)^{\tilde{m}+\xi'+\zeta'+\beta'} D_{p-1}(\partial_{n+1} \tilde{a}, \xi', \zeta', \beta') = 0.$$

When p varies, we see that every term in D is killed. □

Proposition 4.9. *Let ϕ be a normalized reduced cochain in $\bar{\mathbf{C}}_{\text{GS}}^n(\mathcal{A}, M)$ then we have*

$$\mathcal{GF}(\phi) = \phi.$$

Proof. The computations are straightforward. □

REFERENCES

- [1] *Revêtements étales et groupe fondamental (SGA 1)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 3, Société Mathématique de France, Paris, 2003, Séminaire de géométrie algébrique du Bois Marie 1960–61. [Geometric Algebra Seminar of Bois Marie 1960-61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)]. MR MR2017446 (2004g:14017)
- [2] H. Dinh Van, *On the differential of the Yetter complex for monoidal categories*, in preparation.
- [3] H. Dinh Van, L. Liu, and W. Lowen, *Non-commutative deformations and quasi-coherent modules*, *Selecta Mathematica* **23** (2016), no. 2, 1061–1119.
- [4] H. Dinh Van and W. Lowen, *On higher structure on the Gerstenhaber-Schack complex for prestacks*, in preparation.
- [5] M. Doubek, *Gerstenhaber-Schack diagram cohomology from the operadic point of view*, *J. Homotopy Relat. Struct.* **7** (2012), no. 2, 165–206. MR 2988945
- [6] J. Elgueta, *Cohomology and deformation theory of monoidal 2-categories i*, *Adv. Math.* **182** (2004), 204–277.
- [7] Y. Frégier, M. Markl, and D. Yau, *The L_∞ -deformation complex of diagrams of algebras*, *New York J. Math.* **15** (2009), 353–392. MR 2530153 (2011b:16039)
- [8] M. Gerstenhaber and S. D. Schack, *On the deformation of algebra morphisms and diagrams*, *Trans. Amer. Math. Soc.* **279** (1983), no. 1, 1–50. MR MR704600 (85d:16021)
- [9] ———, *Algebraic cohomology and deformation theory*, *Deformation theory of algebras and structures and applications (Il Ciocco, 1986)*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 247, Kluwer Acad. Publ., Dordrecht, 1988, pp. 11–264. MR MR981619 (90c:16016)
- [10] ———, *The cohomology of presheaves of algebras. I. Presheaves over a partially ordered set*, *Trans. Amer. Math. Soc.* **310** (1988), no. 1, 135–165. MR MR965749 (89k:16052)
- [11] J. L. Loday and B. Vallette, *Algebraic operads*, *Grundlehren der mathematischen Wissenschaften*, Vol. 346, Springer, 2012.
- [12] W. Lowen, *Hochschild cohomology of presheaves as map-graded categories*, *Int. Math. Res. Not. IMRN* (2008), Art. ID rnn118, 32. MR 2449052 (2009i:18015)
- [13] W. Lowen and M. Van den Bergh, *A local-to-global spectral sequence for Hochschild cohomology*, in preparation.
- [14] ———, *A Hochschild cohomology comparison theorem for prestacks*, *Trans. Amer. Math. Soc.* **363** (2011), no. 2, 969–986. MR 2728592 (2012c:16033)
- [15] S. Merkulov and B. Vallette, *Deformation theory of representations of prop(erad)s*, <http://arxiv.org/abs/0707.0889>.
- [16] T. Shrestha, *Algebraic deformation of a monoidal category*, PhD. Thesis, Kansas State University 2010, <http://krex.k-state.edu/dspace/handle/2097/6393>.
- [17] D. N. Yetter, *On deformations of pasting diagrams*, *Theory and Application of Category.* **22** (2009), 23–54.

(Dinh Van Hoang) UNIVERSITEIT ANTWERPEN, DEPARTEMENT WISKUNDE-INFORMATICA, MIDDELHEIMCAMPUS, MIDDELHEIMLAAN 1, 2020 ANTWERP, BELGIUM

E-mail address: hoang.dinhvan@uantwerpen.be

(Wendy Lowen) UNIVERSITEIT ANTWERPEN, DEPARTEMENT WISKUNDE-INFORMATICA, MIDDELHEIMCAMPUS, MIDDELHEIMLAAN 1, 2020 ANTWERP, BELGIUM

E-mail address: wendy.lowen@uantwerpen.be