

Hochschild Cohomology with Support

Wendy Lowen

Departement Wiskunde-Informatica, Universiteit Antwerpen,
Middelheimcampus, Middelheimlaan 1, 2020 Antwerp, Belgium

Correspondence to be sent to: e-mail: wendy.lowen@uantwerpen.be

In this paper, we investigate the functoriality properties of map-graded Hochschild complexes. We show that the category Map of map-graded categories is naturally a stack over the category of small categories endowed with a certain Grothendieck topology of 3-covers. For a related topology of ∞ -covers on the Cartesian morphisms in Map , we prove that taking map-graded Hochschild complexes defines a sheaf. From the functoriality related to "injections" between map-graded categories, we obtain Hochschild complexes "with support". We revisit Keller's arrow category argument from this perspective, and introduce and investigate a general Grothendieck construction which encompasses both the map-graded categories associated to presheaves of algebras and certain generalized arrow categories, which together constitute a pair of complementary tools for deconstructing Hochschild complexes.

1 Introduction

Hochschild cohomology originated as a cohomology theory for algebras A . While the degree 0 cohomology is the center of A and the degree 1 cohomology corresponds to the derivations of A , the degree 2 cohomology group parameterizes first-order deformations of A . The relation between Hochschild cohomology and deformation theory is in fact more profound, and can be understood in terms of the structured Hochschild complex

Received November 12, 2013; Accepted February 27, 2014
Communicated by Prof. Dmitry Kaledin

$\mathbf{C}(A)$. In the mean time, Hochschild cohomology and Hochschild complexes have been defined for a wide range of objects of algebro-geometric nature, from schemes [21] and presheaves of algebras [6] to differential graded, exact [12] and abelian categories [15], and various links with deformation theory have been established in these contexts [5, 14, 16].

An important shortcoming of Hochschild cohomology when compared, for instance, with Hochschild homology, is its lack of functoriality. Even for algebras, a morphism $f: A \rightarrow B$ does not naturally give rise to a map between the Hochschild cohomologies $\mathrm{HH}^*(A)$ and $\mathrm{HH}^*(B)$ in either direction. When we turn from algebras to linear categories, that is, algebras with several objects in the sense of [20], the situation becomes somewhat better. More precisely, the inclusion of a full subcategory $\mathfrak{b} \subseteq \mathfrak{a}$ naturally gives rise to a “restriction” map $\mathbf{C}(\mathfrak{a}) \rightarrow \mathbf{C}(\mathfrak{b})$ between the Hochschild complexes. Furthermore, this observation of *limited functoriality* provides a way to relate the Hochschild complexes of categories related by a bimodule, through an intermediate *arrow category*. In [12], the *arrow category argument* for structured (B_∞ -algebra) Hochschild complexes is developed by Keller in the context of dg categories. The main applications of this argument can be divided into two types:

- (1) Relating the Hochschild complex of a more involved object (the arrow category) to the Hochschild complexes of its easier building blocks (two smaller categories and a bimodule relating them).
- (2) Proving that two objects have isomorphic Hochschild complexes in the homotopy category of B_∞ -algebras, by relating them via a suitable bimodule.

Applications of type (1) can be seen as generalizations of results on the Hochschild cohomology of triangular matrix algebras, a topic which, since the work of Happel [11], has received a lot of attention [2, 3, 7–10, 19]. An application of type (2) is the relation between the Hochschild complexes of Koszul dual algebras obtained in [12]. The argument is also extensively used in [15] to compare various candidate Hochschild complexes of abelian categories and ringed spaces. The paper [15] also contains some results more in the spirit of (1), like the existence of Mayer–Vietoris sequences for Hochschild cohomology of ringed spaces.

The main aim of the current work is to provide a comprehensive treatment of the natural tools for “breaking down” complicated Hochschild complexes into easier pieces. In subsequent work, we will apply our results both to improve our understanding of deformation theory, for instance of schemes, and to the computation of Hochschild

cohomology groups of various origins, for instance for singular schemes in the absence of the classical Hochschild-Kostant-Rosenberg decomposition.

The framework we choose for our exposition is that of map-graded categories [14]. Although a more powerful theory can be obtained in the combined context of “map-graded differential graded categories”, for simplicity we present our work in the context of linear map-graded categories. A map-graded category \mathfrak{a} can be viewed as a monoid-graded algebra with several objects. It has an underlying grading category \mathcal{U} , object sets \mathfrak{a}_U for $U \in \mathcal{U}$, and morphism modules $\mathfrak{a}_u(A, A')$ for $u: U \rightarrow U'$ in \mathcal{U} , $A \in \mathfrak{a}_U$ and $A' \in \mathfrak{a}_{U'}$. Thus, the grading category \mathcal{U} can be seen as prescribing a certain shape for \mathfrak{a} . The following are examples of naturally map-graded categories:

- (i) For a presheaf of k -algebras $\mathcal{A}: \mathcal{U} \rightarrow \text{Alg}(k)$, there is an associated \mathcal{U} -graded category \mathfrak{a} obtained as a kind of Grothendieck construction from \mathcal{A} in the spirit of [1] (where the input is a pseudofunctor $\mathcal{U} \rightarrow \text{Cat}$ and the output is a category fibered over \mathcal{U}). The structured Hochschild complex $\mathbf{C}(\mathfrak{a})$ controls the deformation theory of \mathcal{A} as a *twisted* presheaf of algebras [14] and calculates the Hochschild cohomology of \mathcal{A} from [4] as shown in [17].
- (ii) For two linear categories \mathfrak{a} and \mathfrak{b} and an \mathfrak{a} - \mathfrak{b} -bimodule M , the arrow category $(\mathfrak{b} \rightarrow_M \mathfrak{a})$ is naturally graded over the path category of $\bullet \rightarrow \bullet$.

Our point of view is that limited functoriality is determined by grading categories in a fundamental way. In Section 2, we first endow the category Cat of small categories with the Grothendieck pretopology of n -covers (for $n \in \mathbb{N}$) for which a collection of functors $(\varphi_i: \mathcal{V}_i \rightarrow \mathcal{U})_{i \in I}$ is an n -cover provided that it induces a jointly surjective collection of maps $(\mathcal{N}(\varphi_i): \mathcal{N}_n(\mathcal{V}_i) \rightarrow \mathcal{N}(\mathcal{U}))_{i \in I}$ between n -simplices of the simplicial nerves. We show that the category Map of map-graded categories is naturally fibered over Cat (Proposition 2.25) and constitutes a stack for $n \geq 3$ (Corollary 2.40). Let $\text{Map}_c \subseteq \text{Map}$ denote the full subcategory of Cartesian morphisms with respect to the fibered category $\text{Map} \rightarrow \text{Cat}$. In Section 4, we show that taking Hochschild complexes defines a functor (Proposition 4.2)

$$\mathbf{C}: \text{Map}_c \rightarrow B_\infty: (\mathcal{U}, \mathfrak{a}) \mapsto \mathbf{C}_{\mathcal{U}}(\mathfrak{a}).$$

Now endow Map_c with the pretopology of ∞ -covers for which the collection $((\varphi_i, F_i): (\mathcal{V}_i, \mathfrak{b}_i) \rightarrow (\mathcal{U}, \mathfrak{a}))_{i \in I}$ is an ∞ -cover provided that the collection $(\varphi_i: \mathcal{V}_i \rightarrow \mathcal{U})_{i \in I}$ is an ∞ -cover in Cat , that is, an n -cover for every $n \geq 0$. Our Theorem 4.7 implies the following theorem.

Theorem 1.1. The functor $\mathbf{C} : \text{Map}_c \longrightarrow B_\infty$ is a sheaf for the pretopology of ∞ -covers on Map . \square

As an application of the theorem, in Section 4.5 we obtain a Mayer–Vietoris sequence of Hochschild complexes for a \mathcal{U} -graded category \mathfrak{a} and two subcategories $\varphi_i : \mathcal{V}_i \subseteq \mathcal{U}$ for $i \in \{1, 2\}$ that constitute an ∞ -cover of \mathcal{U} :

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{V}_1}(\mathfrak{a}^{\varphi_1}) \oplus \mathbf{C}_{\mathcal{V}_2}(\mathfrak{a}^{\varphi_2}) \longrightarrow \mathbf{C}_{\mathcal{V}_1 \cap \mathcal{V}_2}(\mathfrak{a}^\varphi) \longrightarrow 0.$$

Here $(\mathcal{V}_i, \mathfrak{a}^{\varphi_i}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ and $(\mathcal{V}_1 \cap \mathcal{V}_2, \mathfrak{a}^\varphi) \longrightarrow (\mathcal{U}, \mathfrak{a})$ are chosen to be Cartesian.

In Section 5, we start from a single subcategory $\varphi : \mathcal{V} \subseteq \mathcal{U}$, and a Cartesian functor $(\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$. This gives rise to a surjective morphism between Hochschild complexes $\mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b})$ of which we investigate the kernel $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a})$. The results we obtain depend upon the assumption that the \mathcal{U} -morphisms outside of \mathcal{V} constitute an ideal \mathcal{Z} in \mathcal{U} . In this case, we show in Proposition 5.3 that

$$\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a}) \cong \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, (1_{\mathfrak{a}})_{\mathcal{Z}}),$$

where $(1_{\mathfrak{a}})_{\mathcal{Z}}$ is the natural restriction of $1_{\mathfrak{a}}$ to an \mathfrak{a} -bimodule supported on \mathcal{Z} (i.e., with zero values outside of \mathcal{Z}). An example where our setup applies is the situation where \mathcal{U} is the category associated to a collection of open subsets of a topological space X ordered by inclusion, \mathcal{V} is the full subcategory of subsets $U \subseteq V$ for a fixed subset V , and \mathcal{Z} contains the inclusions $U' \subseteq U$ with $U \not\subseteq V$.

In Section 5.7, we revisit the arrow category construction from [12] in the map-graded context. For a $(\mathcal{U}, \mathfrak{a})$ - $(\mathcal{V}, \mathfrak{b})$ -bimodule (S, M) , we take the natural inclusion $\mathcal{V} \coprod \mathcal{U} \longrightarrow (\mathcal{V} \rightarrow_S \mathcal{U})$ and corresponding Cartesian functor

$$\left(\mathcal{V} \coprod \mathcal{U}, \mathfrak{b} \coprod \mathfrak{a} \right) \longrightarrow (\mathcal{V} \rightarrow_S \mathcal{U}, \mathfrak{b} \rightarrow_M \mathfrak{a}),$$

as starting point for obtaining map-graded analogs of some of the main results from [12]. Sections 5.9 and 5.10 are entirely modeled upon the treatment in [12], and mainly formulate results from [12] in the map-graded context, making use of the natural Hom and tensor functors from Section 3.2. Further, in Section 5.8, we give an intrinsic characterization of arrow categories based upon the *thin ideals* introduced in Section 5.6.

In Section 6, we present a unified framework for constructing map-graded categories and deconstructing their Hochschild complexes. Our main observation is that

both examples (i) and (ii) that we gave of map-graded categories can be viewed as special cases of a generalized Grothendieck construction for map-graded categories. The classical Grothendieck construction from [1] takes a pseudofunctor $\mathcal{U} \rightarrow \text{Cat}$ as input and turns it into a category fibered over \mathcal{U} . The construction from [14] of which we gave an example in (i) is clearly a k -linearized version of this construction, using the category $\text{Cat}(k)$ of k -linear categories instead of Cat . If we relax $\text{Cat}(k)$ to the bicategory $\underline{\text{Cat}}(k)$ of k -linear categories and bimodules, we can in fact describe *any* \mathcal{U} -graded category as a kind of Grothendieck construction of a naturally associated pseudofunctor

$$\mathfrak{a} : \mathcal{U} \rightarrow \underline{\text{Cat}}(k) : U \mapsto \mathfrak{a}_U.$$

In the paper, we go yet another step further and start from a pseudofunctor

$$(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \rightarrow \underline{\text{Map}} : C \mapsto (\mathcal{U}_C, \mathfrak{a}_C),$$

where \mathcal{C} is a small category and $\underline{\text{Map}}$ is the bicategory of map-graded categories and bimodules described in Section 3.2. Allowing arbitrary bimodules rather than functors between map-graded categories allows us to capture the arrow category with respect to a bimodule from (ii). In general, we can now deconstruct the Hochschild complex of the Grothendieck construction $(\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ of $(\mathcal{U}, \mathfrak{a})$ based upon the internal structure of \mathcal{C} . Here, the strategy is to cover $(\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ by other Grothendieck constructions, based upon base change for pseudofunctors from Section 6.5. For instance, in Proposition 6.7, we prove the following proposition.

Proposition 1.2. Suppose that \mathcal{C} has finite products. Let \mathcal{C}^* be the category \mathcal{C} with terminal object $*$ adjoined. Put $(\tilde{\mathcal{U}}|_*, \tilde{\mathfrak{a}}|_*) = (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ and let $(\tilde{\mathcal{U}}|_C, \tilde{\mathfrak{a}}|_C)$ be the Grothendieck construction of the restriction of $(\mathcal{U}, \mathfrak{a})$ to \mathcal{C}/\mathcal{C} . Let $(C_i)_{i \in I}$ be a collection of objects in \mathcal{C} such that for every $C \in \mathcal{C}$ there exists a map $C \rightarrow C_i$ for some i . There is a natural functor

$$\mathbf{C} : \mathcal{C}^* \rightarrow B_\infty : C \mapsto \mathbf{C}_{\tilde{\mathcal{U}}|_C}(\tilde{\mathfrak{a}}|_C),$$

which satisfies the sheaf property with respect to the collection of maps $(C_i \rightarrow *)_{i \in I}$ in \mathcal{C}^* . □

Let us now restrict our attention to the case where \mathcal{C} is the category associated to a poset. Then it is not hard to see that \mathcal{C} can be covered by path categories of:

$$\bullet \rightarrow \bullet \rightarrow \dots \rightarrow \dots \rightarrow \bullet \rightarrow \bullet.$$

In Section 6.6, we consider the Grothendieck constructions of such categories as “generalized arrow categories”, and deconstruct them using iterated arrow category constructions. As such, we see in Section 6.7 how the sheaf property for Hochschild complexes on the one hand, and the arrow category construction on the other hand, can be seen as complementary tools for deconstructing Hochschild cohomology.

In Section 6.8, we consider a pseudofunctor $(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \rightarrow \text{Map}_c$ with some extension $(\mathcal{U}^*, \mathfrak{a}^*) : \mathcal{C}^* \rightarrow \text{Map}_c$, which we compare with the natural pseudofunctor of Grothendieck constructions

$$(\mathcal{U}^*, \mathfrak{a}^*) : \mathcal{C}^* \rightarrow \text{Map}_c : \mathcal{C} \rightarrow (\tilde{\mathcal{U}}|_{\mathcal{C}}, \tilde{\mathfrak{a}}|_{\mathcal{C}}).$$

Our main Theorem 6.14 encompasses the following comparison result.

Theorem 1.3. There is a morphism of pseudofunctors landing in the homotopy category of B_∞ -algebras

$$\mathbf{G}_{\mathcal{U}^*}(\mathfrak{a}^*) \rightarrow \mathbf{G}_{\mathcal{U}}(\mathfrak{a}),$$

in which every component morphism is a quasi-isomorphism except possibly the top one. The situation where the top morphism is a quasi-isomorphism too can be characterized by the usual bimodule criterion from [12]. \square

The proof of the theorem is heavily based upon Keller’s arrow category argument in the case of a fully faithful functor $\mathfrak{b} \rightarrow \mathfrak{a}$, which is in fact a special case of our theorem. A combination of Proposition 1.2 and Theorem 1.3 allows us to pull a deconstruction of Hochschild complexes of Grothendieck constructions back to a deconstruction of the Hochschild complexes of the categories $(\mathcal{U}_*^*, \mathfrak{a}_*^*)$ and $(\mathcal{U}_c, \mathfrak{a}_c)$ in which we are primarily interested—at least in the homotopy category of B_∞ -algebras. In particular, exact sequences of Mayer–Vietoris type now naturally give rise to Mayer–Vietoris exact triangles, inducing the desired long exact cohomology sequences. As an application, we recover the Mayer–Vietoris triangles for ringed spaces from [15, Section 7.9].

2 Map-Graded Categories

Let k be a commutative ground ring. In this section, we introduce the category Map of small k -linear map-graded categories and functors, which is naturally fibered over the category Cat of small categories and functors. We also introduce the intermediate category Mas of map-graded sets. A map-graded set $(\mathcal{U}, \mathfrak{a})$ consists of a small category \mathcal{U} and for every object $U \in \mathcal{U}$, a set \mathfrak{a}_U . A map-graded category $(\mathcal{U}, \mathfrak{a})$ is a map-graded

set with additionally, for every morphism $u: U \rightarrow U'$ in \mathcal{U} , $A \in \mathfrak{a}_U$ and $A' \in \mathfrak{a}_{U'}$, a k -module $\mathfrak{a}_u(A, A')$. These modules are endowed with category-like composition and identity morphism [14]. Every map-graded set or category $(\mathcal{U}, \mathfrak{a})$ has an associated object $(\mathcal{U}^\sharp, \mathfrak{a}^\sharp)$ of the same kind, with “ungrouped object sets”, that is, with $\text{Ob}(\mathcal{U}^\sharp) = \coprod_{U \in \mathcal{U}} \mathfrak{a}_U$ and $(\mathfrak{a}^\sharp)_A = \{A\}$. We define the *nerve* $\mathcal{N}(\mathfrak{a})$ of $(\mathcal{U}, \mathfrak{a})$ to be the simplicial nerve $\mathcal{N}(\mathcal{U}^\sharp)$. For small categories, map-graded sets, and map-graded categories, we define pretopologies of n -covers by declaring a collection of functors to be an n -cover provided that for $p \leq n$ the induced collection of maps between p -simplices of the nerves is jointly surjective (Definition 2.30). We prove that for $n \geq 0$, Mas is a stack over Cat (Theorem 2.36) and for $n \geq 3$, Map is a stack over Mas (Theorem 2.38) and over Cat (Corollary 2.40).

2.1 Fibered categories and stacks

In this section, we recall the basic concepts concerning fibered categories and stacks [1, 22], and give some results we will use later on.

Let \mathcal{U} be a category. A *category over \mathcal{U}* is a functor $F: \mathcal{A} \rightarrow \mathcal{U}$. The functor is perceived from the point of view of its fibers. That is, for every $U \in \mathcal{U}$ we consider the fiber of objects over U :

$$\mathcal{A}_U = \{A \in \mathcal{A} \mid F(A) = U\},$$

and for $u: V \rightarrow U$, $A \in \mathcal{A}_U$, $B \in \mathcal{A}_V$, we consider the fiber of morphisms over u :

$$\mathcal{A}_u(B, A) = \{a \in \mathcal{A}(B, A) \mid F(a) = u\}.$$

A morphism $a: B \rightarrow A$ in \mathcal{A} with $F(a) = u: V \rightarrow U$ is *Cartesian (with respect to F)* provided that for every $v: W \rightarrow V$ in \mathcal{U} and $C \in \mathcal{A}_W$, the composition map

$$a- : \mathcal{A}_v(C, B) \rightarrow \mathcal{A}_{uv}(C, A)$$

is an isomorphism.

The category \mathcal{A} is called *fibered over \mathcal{U}* provided that for every $u: V \rightarrow U$ in \mathcal{U} and $A \in \mathcal{A}_U$, there is an object $u^*A \in \mathcal{A}_V$ and a Cartesian morphism $\delta^{u,A} \in \mathcal{A}_u(u^*A, A)$. The choice, for every u and A , of such a Cartesian morphism $\delta^{u,A}$ is called a *cleavage*. For a fibered category with a cleavage, there is an associated pseudofunctor

$$\mathcal{A}: \mathcal{U}^{\text{op}} \rightarrow \text{Cat}: U \mapsto \mathcal{A}(U),$$

where \mathbf{Cat} denotes the category of small categories. Here $\mathcal{A}(U)$ is the category with object set given by \mathcal{A}_U and $\mathcal{A}(U)(B, A) = \mathcal{A}_{1_U}(B, A)$. For a map $u: V \rightarrow U$ in \mathcal{U} , the corresponding functor $u^*: \mathcal{A}(U) \rightarrow \mathcal{A}(V)$ is naturally determined by the chosen Cartesian morphisms.

Proposition 2.1. Let $F: \mathcal{A} \rightarrow \mathcal{U}$ be a fibered category over \mathcal{U} with a cleavage. Suppose for $v: W \rightarrow V, u: V \rightarrow U$ in \mathcal{U} and $A \in \mathcal{A}_U$, and for the Cartesian morphisms $\delta^{u,A}: u^*A \rightarrow A, \delta^{v,u^*A}: v^*u^*A \rightarrow u^*A$, and $\delta^{uw,A}: (uw)^*A \rightarrow A$, we have $v^*u^*A = (uw)^*A$ and $\delta^{u,A}\delta^{v,u^*A} = \delta^{uw,A}$. Then the associated pseudofunctor $\mathcal{A}: \mathcal{U}^{\text{op}} \rightarrow \mathbf{Cat}$ is actually a functor. \square

Next we recall the definition of a pretopology. Let \mathcal{U} be a category with pullbacks. A *pretopology* on \mathcal{U} consists of the notion of a *covering collection* of maps $(U_i \rightarrow U)_{i \in I}$, also called a *cover* of the object U , satisfying the following axioms:

- (1) The collection consisting of $1_U: U \rightarrow U$ is a cover of U .
- (2) If $(U_i \rightarrow U)_{i \in I}$ is a cover of U and $u: V \rightarrow U$ an arbitrary map, the collection of pullbacks $(V \times_U U_i \rightarrow V)_{i \in I}$ is a cover of V .
- (3) If $(U_i \rightarrow U)_{i \in I}$ is a cover of U and for every $i \in I, (U_{ij} \rightarrow U_i)_{j \in J_i}$ is a cover of U_i , then the collection of compositions $(U_{ij} \rightarrow U_i \rightarrow U)_{i \in I, j \in J_i}$ is a cover of U .

On a category \mathcal{U} with pullbacks, the notion of pretopology can be seen as describing somewhat more tangible input data for a Grothendieck topology. More precisely, a Grothendieck topology is obtained by defining a sieve $R \subseteq \mathcal{U}(-, U)$ to be covering if and only if it contains a covering collection of maps for the given pretopology.

Consider a fibered category \mathcal{A} over \mathcal{U} with a cleavage and associated pseudofunctor. Let there be given a pretopology on \mathcal{U} and let $S = (U_i \rightarrow U)_{i \in I}$ be a cover of $U \in \mathcal{U}$. The *descent category* $\text{Des}(S, \mathcal{A})$ is defined in the following way. An object, called a *descent datum*, consists of a collection $(A_i)_{i \in I}$ of objects with $A_i \in \mathcal{A}(U_i)$, together with for every $i, j \in I$ an isomorphism $\alpha_1^*(A_i) \cong \alpha_2^*(A_j)$ in $\mathcal{A}(U_{ij})$ for the pullback

$$\begin{array}{ccc}
 U_i & \longrightarrow & U \\
 \alpha_1 \uparrow & & \uparrow \\
 U_{ij} & \longrightarrow & U_j \\
 & \alpha_2 &
 \end{array}$$

These isomorphisms have to satisfy the natural compatibility requirement on triple pullbacks. A morphism between descent data $(A_i)_i \rightarrow (B_i)_i$ consists of compatible

morphisms $A_i \rightarrow B_i$ in $\mathcal{A}(U_i)$. The fibered category \mathcal{A} is called a *stack* (respectively, a *prestack*) provided that the natural comparison functor

$$\mathcal{A}(U) \rightarrow \text{Des}(S, \mathcal{A}) : A \mapsto (u_i^* A)_i$$

is an equivalence of categories (respectively, fully faithful) for every cover $S = (u_i : U_i \rightarrow U)_{i \in I}$.

Next we collect some useful facts concerning composable functors $G : \mathcal{A} \rightarrow \mathcal{X}$ and $F : \mathcal{X} \rightarrow \mathcal{U}$. We suppose that all three categories have pullbacks and the functors F and G preserve pullbacks.

Proposition 2.2. (1) If a morphism $a : A \rightarrow A'$ in \mathcal{A} is Cartesian with respect to G and $G(a)$ is Cartesian with respect to F , then a is Cartesian with respect to FG .

(2) If both F and G are fibered, then so is FG . □

Proposition 2.3. Suppose that \mathcal{U} is endowed with a pretopology. There is a pretopology on \mathcal{X} for which $(x_i : X_i \rightarrow X)_i$ is a cover of X if and only if $(F(x_i) : F(X_i) \rightarrow F(X))_i$ is a cover of $F(X)$. □

Proposition 2.4. Consider morphisms in \mathcal{X} :

$$X_2 \xrightarrow{x_2} X_1 \xrightarrow{x_1} X .$$

- (1) If x_2 and x_1 are Cartesian, then so is $x_1 x_2$.
- (2) If x_1 and $x_1 x_2$ are Cartesian, then so is x_2 . □

Proposition 2.5. Consider a commutative square in \mathcal{X} :

$$\begin{array}{ccc} X_1 & \xrightarrow{x_1} & X \\ y_1 \uparrow & & \uparrow x_2 \\ X_{12} & \xrightarrow{y_2} & X_2 \end{array}$$

- (1) If the square is a pullback and x_2 is Cartesian, then so is y_1 .
- (2) If the image of the square under F is a pullback, and the morphisms x_1, x_2, y_1, y_2 are Cartesian, then the square is a pullback. □

Proposition 2.6. There is a pretopology on \mathcal{X} for which $(x_i : X_i \rightarrow X)_i$ is a cover of X if and only if the morphisms x_i are Cartesian with respect to F . \square

Proposition 2.7. Suppose that \mathcal{U} is endowed with a pretopology. There is a pretopology on \mathcal{X} for which $(x_i : X_i \rightarrow X)_i$ is a cover of X if and only if the following two conditions hold:

- (1) $(F(x_i) : F(X_i) \rightarrow F(X))_i$ is a cover of $F(X)$;
- (2) every x_i is Cartesian with respect to F . \square

Now suppose that both F and G are fibered, and consider a cleavage for both functors. For a morphism $u : V \rightarrow U$ in \mathcal{U} and $A \in \mathcal{A}_U$ with respect to FG , we obtain the Cartesian morphism $x = \delta^{u, G(A)} : u^*G(A) \rightarrow G(A)$ in \mathcal{X} . Next we obtain the Cartesian morphism $\delta^{x, A} : x^*A \rightarrow A$. We obtain a cleavage for FG by putting $u^*A = x^*A$ and $\delta^{u, A} = \delta^{x, A}$. Let \mathcal{U} be endowed with a pretopology and endow \mathcal{X} with the pretopology described in Proposition 2.7. Let \mathcal{X}^F be the pseudofunctor associated to $F : \mathcal{X} \rightarrow \mathcal{U}$, \mathcal{A}^G be the one associated to $G : \mathcal{A} \rightarrow \mathcal{X}$, and \mathcal{A}^{FG} be the one associated to $FG : \mathcal{A} \rightarrow \mathcal{U}$.

Proposition 2.8. If \mathcal{X}^F and \mathcal{A}^G are stacks (respectively, prestacks), then so is \mathcal{A}^{FG} . \square

2.2 Map-graded categories and functors

Let \mathcal{U} be a base category. A \mathcal{U} -graded set \mathfrak{r} consists of the datum, for every $U \in \mathcal{U}$, of a set \mathfrak{r}_U . A (k -linear) \mathcal{U} -graded category (see [14]) \mathfrak{a} consists of the following data:

- (1) For every $U \in \mathcal{U}$, a set \mathfrak{a}_U of *objects over* U .
- (2) For every $u : V \rightarrow U$ in \mathcal{U} , $B \in \mathfrak{a}_V$, $A \in \mathfrak{a}_U$, a k -module $\mathfrak{a}_u(B, A)$ of *morphisms over* u .
- (3) For a further $v : W \rightarrow V$ in \mathcal{U} and $C \in \mathfrak{a}_W$, a *composition map*

$$\mathfrak{a}_u(B, A) \otimes_k \mathfrak{a}_v(C, B) \longrightarrow \mathfrak{a}_{uv}(C, A).$$

- (1) For $A \in \mathfrak{a}_U$, an *identity element*

$$1_A \in \mathfrak{a}_{1_U}(A, A).$$

These data should satisfy the obvious category-type axioms, that is, the composition has to be associative and the identity elements have to act identically under composition. There is an associated k -linear category $\tilde{\mathfrak{a}}$ with $\text{Ob}(\tilde{\mathfrak{a}}) = \coprod_{U \in \mathcal{U}} \mathfrak{a}_U$ and

$\mathfrak{a}(A_V, A_U) = \bigoplus_{u \in \mathcal{U}(V, U)} \mathfrak{a}_u(A_V, A_U)$. Clearly, a \mathcal{U} -graded category has an underlying \mathcal{U} -graded set given by the object sets.

Remark 2.9. The definition of a k -linear \mathcal{U} -graded category is a k -linearized version of the notion of a category over \mathcal{U} from Section 2.1. Indeed, if we drop k -linearity from the definition, a \mathcal{U} -graded category \mathcal{A} now has a natural associated category $\tilde{\mathcal{A}}$ with $\tilde{\mathcal{A}}(A_V, A_U) = \coprod_{u \in \mathcal{U}(V, U)} \mathfrak{a}_u(A_V, A_U)$. Hence, there is a natural functor $\tilde{\mathcal{A}} \rightarrow \mathcal{U}$, and the datum of this functor is equivalent to the datum of \mathcal{A} . To avoid confusion, for us a *\mathcal{U} -graded category* will implicitly mean a k -linear \mathcal{U} -graded category over some fixed k , whereas we will refer to the nonlinear variant explicitly as a *nonlinear \mathcal{U} -graded category*. \square

Example 2.10. Suppose that \mathcal{U} has a single object $*$ and $\mathfrak{a}_* = \{*\}$. Then $G = \mathcal{U}(*, *)$ is a monoid, and \mathfrak{a} corresponds to a G -graded algebra A with $A_g = \mathfrak{a}_g(*, *)$. Thus, in the spirit of [20], we can view map-graded categories as monoid-graded algebras with several objects. \square

Example 2.11. A k -linear category \mathfrak{a} can be made into a graded category in several natural ways:

- (1) Let $\mathcal{U}_\mathfrak{a}$ be the category with $\text{Ob}(\mathcal{U}_\mathfrak{a}) = \text{Ob}(\mathfrak{a})$ and $\mathcal{U}_\mathfrak{a}(B, A) = \{*\}$ for all $B, A \in \mathfrak{a}$. Then we can make \mathfrak{a} into a $\mathcal{U}_\mathfrak{a}$ -graded category \mathfrak{a}_{st} with $(\mathfrak{a}_{\text{st}})_A = \{A\}$ and $(\mathfrak{a}_{\text{st}})_*(B, A) = \mathfrak{a}(B, A)$. We refer to this grading as the *standard grading* on \mathfrak{a} .
- (2) Let e be the category with one object $*$ and one morphism 1_* . Then we can make \mathfrak{a} into an e -graded category \mathfrak{a}_{tr} with $(\mathfrak{a}_{\text{tr}})_* = \text{Ob}(\mathfrak{a})$ and $(\mathfrak{a}_{\text{tr}})_{1_*}(B, A) = \mathfrak{a}(B, A)$. We refer to this grading as the *trivial grading* on \mathfrak{a} .
- (3) Clearly, every partition of $\text{Ob}(\mathfrak{a})$ will give rise to a graded incarnation of \mathfrak{a} “in between” the two extremes described in (1) and (2). \square

Example 2.12. Let \mathcal{U} be a category. The *free \mathcal{U} -graded category* $k\mathcal{U}$ is defined by putting $k\mathcal{U}_U = \{U\}$ and $k\mathcal{U}_u(V, U) = k$ for all $u: V \rightarrow U \in \mathcal{U}$. Compositions are defined by means of the multiplication of k , and identity elements are provided by the unit of k . Instead of k , we can actually use an arbitrary k -algebra A and perform a similar construction. \square

Example 2.13. Let \mathfrak{r} be a \mathcal{U} -graded set. There is an associated small category \mathcal{U}^\sharp and a \mathcal{U}^\sharp -graded set \mathfrak{r}^\sharp involving only singleton sets above the objects of \mathcal{U}^\sharp . Precisely, we put $\text{Ob}(\mathcal{U}^\sharp) = \coprod_{U \in \mathcal{U}} \mathfrak{r}_U$ and $\mathcal{U}^\sharp(X_U, X_{U'}) = \mathcal{U}(U, U')$, and $\mathfrak{r}^\sharp_{X_U} = \{\mathfrak{r}_U\}$. If \mathfrak{a} is a \mathcal{U} -graded

category, we further obtain the \mathcal{U}^\sharp -graded category \mathfrak{a}^\sharp with in addition $\mathfrak{a}_u^\sharp(A_U, A_{U'}) = \mathfrak{a}_u(A_U, A_{U'})$. \square

Example 2.14. With the notation of Example 2.11, for a k -linear category, we have $(\mathcal{U}_\mathfrak{a}, \mathfrak{a}_{\text{st}}) = (e, \mathfrak{a}_{\text{tr}})^\sharp$. \square

Let $\varphi : \mathcal{V} \rightarrow \mathcal{U}$ be a functor, \mathfrak{x} be a \mathcal{U} -graded set, and η be a \mathcal{V} -graded set. A φ -graded map F consists of maps $F_V : \eta_V \rightarrow \mathfrak{x}_{\varphi(V)}$ for every $V \in \mathcal{V}$.

Let \mathfrak{a} be a \mathcal{U} -graded category and \mathfrak{b} be a \mathcal{V} -graded category. A φ -graded functor F consists of the following data:

- (1) For every $V \in \mathcal{V}$, a map $F_V : \mathfrak{b}_V \rightarrow \mathfrak{a}_{\varphi(V)}$.
- (2) For every $v : V \rightarrow V'$ in \mathcal{V} , $B \in \mathfrak{b}_V$ and $B' \in \mathfrak{b}_{V'}$, a k -linear map

$$\mathfrak{b}_v(B, B') \rightarrow \mathfrak{a}_{\varphi(v)}(F(B), F(B')).$$

These data should satisfy the obvious functoriality-type axioms, that is, F respects compositions and identity elements. Clearly, a φ -graded functor has an underlying φ -graded map between object sets. Let an underlying ground ring k be fixed. Map-graded categories (respectively, sets) and functors (respectively, maps) constitute a category Map (respectively, Mas) in the following way. An object of Map (respectively, Mas) is given by a small category \mathcal{U} and a \mathcal{U} -graded category (respectively, set) \mathfrak{a} . A morphism $(\mathcal{U}, \mathfrak{a}) \mapsto (\mathcal{V}, \mathfrak{b})$ is given by a functor $\varphi : \mathcal{U} \rightarrow \mathcal{V}$ and a φ -graded functor (respectively, map) $F : \mathfrak{a} \rightarrow \mathfrak{b}$.

Example 2.15. Let \mathfrak{a} be a linear category. Let $(\mathfrak{a}_{\text{st}}, \mathcal{U}_\mathfrak{a})$ be the graded category resulting from the standard grading on \mathfrak{a} as in Example 2.11(1) and let $(\mathfrak{a}_{\text{tr}}, e)$ be the graded category resulting from the trivial grading on \mathfrak{a} as in Example 2.11(2). Let $\varphi : \mathcal{U}_\mathfrak{a} \rightarrow e$ be the unique functor. The map $(\mathfrak{a}_{\text{st}})_A = \{A\} \rightarrow \text{Ob}(\mathfrak{a}) = (\mathfrak{a}_{\text{tr}})_* : A \mapsto A$ and the maps $1 : (\mathfrak{a}_{\text{st}})_*(A, A') = \mathfrak{a}(A, A') \rightarrow \mathfrak{a}(A, A') = (\mathfrak{a}_{\text{tr}})_*(A, A')$ define a φ -graded functor. \square

Example 2.16. Let \mathfrak{a} be a \mathcal{U} -graded category and \mathfrak{b} be a linear category with associated e -graded category \mathfrak{b}_{tr} with trivial grading as in Example 2.11(2) and let $\varphi : \mathcal{U} \rightarrow e$ be the unique functor. There is a one-to-one correspondence between φ -graded functors $\mathfrak{a} \rightarrow \mathfrak{b}_{\text{tr}}$ and k -linear functors $\tilde{\mathfrak{a}} \rightarrow \mathfrak{b}$. \square

Example 2.17. With the notation of Example 2.13, let \mathfrak{a} be a \mathcal{U} -graded set, respectively, category and let $\varphi_\mathfrak{a} : \mathcal{U}^\sharp \rightarrow \mathcal{U}$ be the forgetful functor. There is a $\varphi_\mathfrak{a}$ -graded map, respectively, functor $F_\mathfrak{a} : \mathfrak{a}^\sharp \rightarrow \mathfrak{a}$ given by $F(A_U) = A_U \in \mathfrak{a}_U$ (and $1 : \mathfrak{a}_u^\sharp(A, A') \rightarrow \mathfrak{a}_u(A, A')$ in the

category case). A graded map, respectively, functor $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ gives rise to a commutative square

$$\begin{array}{ccc} (\mathcal{V}, \mathfrak{b}) & \xrightarrow{(\varphi, F)} & (\mathcal{U}, \mathfrak{a}) \\ (\varphi_{\mathfrak{b}}, F_{\mathfrak{b}}) \uparrow & & \uparrow (\varphi_{\mathfrak{a}}, F_{\mathfrak{a}}) \\ (\mathcal{V}^{\sharp}, \mathfrak{b}^{\sharp}) & \xrightarrow{(\varphi^{\sharp}, F^{\sharp})} & (\mathcal{U}^{\sharp}, \mathfrak{a}^{\sharp}) \end{array}$$

We thus obtain a natural functor

$$(-)^{\sharp} : \text{Map} \longrightarrow \text{Map} : (\mathcal{U}, \mathfrak{a}) \longmapsto (\mathcal{U}^{\sharp}, \mathfrak{a}^{\sharp}),$$

and similarly in the case of Mas. □

Example 2.18. Denote the category of small k -linear categories by $\text{Cat}(k)$. There are natural functors $\text{Map} \longrightarrow \text{Cat}(k) : \mathfrak{a} \longmapsto \tilde{\mathfrak{a}}$ and $\text{Cat}(k) \longrightarrow \text{Map} : \mathfrak{a} \longmapsto \mathfrak{a}_{\text{tr}}$. By Example 2.16, the first functor is left adjoint to the second functor. □

2.3 Two fibered categories of map-graded categories

Let k be a fixed commutative ground ring. We consider the categories Map and Mas as defined in Section 2.2, as well as the category Cat of small categories. There are natural forgetful functors

$$\psi_1 : \text{Map} \longrightarrow \text{Mas} : (\mathcal{U}, \mathfrak{a}) \longmapsto (\mathcal{U}, \mathfrak{r}_{\mathfrak{a}}),$$

where $\mathfrak{r}_{\mathfrak{a}}$ is the underlying \mathcal{U} -graded set of objects of \mathfrak{a} and

$$\psi_0 : \text{Mas} \longrightarrow \text{Cat} : (\mathcal{U}, \mathfrak{a}) \longmapsto \mathcal{U}.$$

Further, all three categories Map , Mas , and Cat have pullbacks and the two functors ψ_1 and ψ_0 preserve them. We first look at Cat . For functors $\varphi_1 : \mathcal{V}_1 \longrightarrow \mathcal{U}$ and $\varphi_2 : \mathcal{V}_2 \longrightarrow \mathcal{U}$, the pullback

$$\begin{array}{ccc} \mathcal{V}_1 & \xrightarrow{\varphi_1} & \mathcal{U} \\ \alpha_1 \uparrow & & \uparrow \varphi_2 \\ \mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2 & \xrightarrow{\alpha_2} & \mathcal{V}_2 \end{array} \tag{1}$$

is given by the category $\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2$ where

$$\text{Ob}(\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2) = \text{Ob}(\mathcal{V}_1) \times_{\text{Ob}(\mathcal{U})} \text{Ob}(\mathcal{V}_2)$$

and

$$(\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2)((V_1, V_2), (V'_1, V'_2)) = \mathcal{V}_1(V_1, V'_1) \times_{\mathcal{U}(\varphi_1(V_1), \varphi_2(V_2))} \mathcal{V}_2(V_2, V'_2)$$

are given by the pullbacks in the category of sets. For a collection of functors $\varphi_i : \mathcal{V}_i \rightarrow \mathcal{U}$, we similarly obtain a limit category $\prod_{i, \mathcal{U}} \mathcal{V}_i$.

Example 2.19. Consider subcategories $\mathcal{V}_1 \subseteq \mathcal{U}$ and $\mathcal{V}_2 \subseteq \mathcal{U}$. We define the category $\mathcal{V}_1 \cap \mathcal{V}_2$ with $\text{Ob}(\mathcal{V}_1 \cap \mathcal{V}_2) = \text{Ob}(\mathcal{V}_1) \cap \text{Ob}(\mathcal{V}_2)$ and $(\mathcal{V}_1 \cap \mathcal{V}_2)(V, V') = \mathcal{V}_1(V, V') \cap \mathcal{V}_2(V, V')$. The canonical inclusion functors $\mathcal{V}_1 \cap \mathcal{V}_2 \subseteq \mathcal{V}_1$ and $\mathcal{V}_2 \subseteq \mathcal{V}_1 \cap \mathcal{V}_2$ induce an isomorphism of categories $\mathcal{V}_1 \cap \mathcal{V}_2 \cong \mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2$. \square

Pullbacks in Mas and Map are described in a similar fashion. For instance, the pullback of graded categories is described by

$$\begin{array}{ccc} (\mathcal{V}_1, \mathfrak{b}_1) & \xrightarrow{(\varphi_1, F_1)} & (\mathcal{U}, \mathfrak{a}) \\ \uparrow (\alpha_1, G_1) & & \uparrow (\varphi_2, F_2) \\ (\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2, \mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2) & \xrightarrow{(\alpha_2, G_2)} & (\mathcal{V}_2, \mathfrak{b}_2) \end{array}$$

with underlying pullback of categories described by (1) and with

$$(\mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2)_{(V_1, V_2)} = (\mathfrak{b}_1)_{V_1} \times_{\mathfrak{a}_{\varphi_1(V_1)}} (\mathfrak{b}_2)_{V_2}$$

and

$$(\mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2)_{(v_1, v_2)}((B_1, B_2), (B'_1, B'_2)) = (\mathfrak{b}_1)_{v_1}(B_1, B'_1) \times_{\mathfrak{a}_{\varphi_1(v_1)}(F_1(B_1), F_1(B'_1))} (\mathfrak{b}_2)_{v_2}(B_2, B'_2).$$

Example 2.20. The functors $(-)^{\sharp} : \text{Map} \rightarrow \text{Map}$ and $(-)^{\sharp} : \text{Mas} \rightarrow \text{Mas}$ from Example 2.17 preserve pullbacks. \square

Thus, the results of Section 2.1 apply to the composable functors Ψ_0 and Ψ_1 . In the remainder of this section, we show that both these functors are fibered.

We start with Ψ_0 . Let $\varphi : \mathcal{V} \longrightarrow \mathcal{U}$ be a functor and \mathfrak{r} be a \mathcal{U} -graded set. We define the \mathcal{V} -graded set \mathfrak{r}^φ with

$$\mathfrak{r}_V^\varphi = \mathfrak{r}_{\varphi(V)},$$

and the morphism of graded sets $(\varphi, \delta^{\varphi, \mathfrak{r}}) : (\mathcal{V}, \mathfrak{r}^\varphi) \longrightarrow (\mathcal{U}, \mathfrak{r})$ with $\delta_V^{\varphi, \mathfrak{r}} : \mathfrak{r}_V^\varphi = \mathfrak{r}_{\varphi(V)} \longrightarrow \mathfrak{r}_{\varphi(V)}$ given by the identity morphism.

Proposition 2.21. A morphism $(\varphi, F) : (\mathcal{V}, \eta) \longrightarrow (\mathcal{U}, \mathfrak{r})$ of graded sets is Cartesian with respect to Ψ_0 if and only if for every $V \in \mathcal{V}$ the map $F_V : \eta_V \longrightarrow \mathfrak{r}_{\varphi(V)}$ is an isomorphism of sets. \square

Proposition 2.22. (1) The morphisms $(\varphi, \delta^{\varphi, \mathfrak{r}})$ are Cartesian with respect to Ψ_0 .

(2) The category **Mas** is fibered over **Cat** through Ψ_0 . \square

Next we look at Ψ_1 . Let $(\varphi, F) : (\mathcal{V}, \eta) \longrightarrow (\mathcal{U}, \mathfrak{r})$ be a morphism of graded sets, and let \mathfrak{a} be a \mathcal{U} -graded category with underlying graded set \mathfrak{r} . We define the \mathcal{V} -graded category $\mathfrak{a}^{\varphi, F}$ with

$$\mathfrak{a}_V^{\varphi, F} = \eta_V,$$

and with, for $v : V \longrightarrow V'$ in \mathcal{V} , $Y \in \eta_V$, $Y' \in \eta_{V'}$:

$$\mathfrak{a}_v^{\varphi, F}(Y, Y') = \mathfrak{a}_{\varphi(v)}(F(Y), F(Y')),$$

and with, for another $v' : V' \longrightarrow V''$ in \mathcal{V} and $Y'' \in \eta_{V''}$, the composition in $\mathfrak{a}^{\varphi, F}$

$$\mathfrak{a}_{\varphi(v')}(F(Y'), F(Y'')) \otimes \mathfrak{a}_{\varphi(v)}(F(Y), F(Y')) \longrightarrow \mathfrak{a}_{\varphi(v'v)}(F(Y), F(Y''))$$

defined by the composition in \mathfrak{a} , and similarly for the identity elements in $\mathfrak{a}_{1_V}^{\varphi, F}(Y, Y) = \mathfrak{a}_{1_{\varphi(V)}}(F(Y), F(Y))$. We define the morphism of graded categories $(\varphi, \delta = \delta^{\varphi, F, \mathfrak{a}}) : (\mathcal{V}, \mathfrak{a}^{\varphi, F}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ by the morphisms $\delta_V = F_V : \eta_V \longrightarrow \mathfrak{a}_{\varphi(V)}$ and the identity morphisms

$$\mathfrak{a}_v^{\varphi, F}(Y, Y') = \mathfrak{a}_{\varphi(v)}(F(Y), F(Y')) \longrightarrow \mathfrak{a}_{\varphi(v)}(F(Y), F(Y')).$$

Proposition 2.23. A morphism $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ of graded categories is Cartesian with respect to Ψ_1 if and only if for every $v : V \longrightarrow V'$ in \mathcal{V} , $B \in \mathfrak{b}_V$, $B' \in \mathfrak{b}_{V'}$, the map $\mathfrak{b}_v(B, B') \longrightarrow \mathfrak{a}_{\varphi(v)}(F(B), F(B'))$ is an isomorphism. \square

Proposition 2.24. (1) The morphisms $(\varphi, \delta^{\varphi, F, \mathfrak{a}})$ are Cartesian with respect to Ψ_1 .

(2) The category **Map** is fibered over **Mas** through Ψ_1 . \square

By Proposition 2.2, as a consequence of Propositions 2.22 and 2.24, the category Map is fibered over Cat through Ψ . We end this section by describing the cleavage which follows from the higher cleavages for Ψ_0 and Ψ_1 .

Let \mathfrak{a} be a \mathcal{U} graded category and let $\varphi : \mathcal{V} \rightarrow \mathcal{U}$ be a functor. We define the \mathcal{V} -graded category \mathfrak{a}^φ with

$$\mathfrak{a}_V^\varphi = \mathfrak{a}_{\varphi(V)},$$

and with, for $v : V \rightarrow V'$ in \mathcal{V} , $A \in \mathfrak{a}_V^\varphi$ and $A' \in \mathfrak{a}_{V'}^\varphi$

$$\mathfrak{a}_v^\varphi(A, A') = \mathfrak{a}_{\varphi(v)}(A, A').$$

For another $v' : V' \rightarrow V''$ in \mathcal{V} and $A'' \in \mathfrak{a}_{V''}^\varphi$, the composition in \mathfrak{a}^φ

$$\mathfrak{a}_{\varphi(v')}^\varphi(A', A'') \otimes \mathfrak{a}_{\varphi(v)}^\varphi(A, A') \rightarrow \mathfrak{a}_{\varphi(v'v)}^\varphi(A, A'')$$

is defined by the composition in \mathfrak{a} , and similarly for the identity elements in $\mathfrak{a}_{\varphi(1_V)}^\varphi(A, A) = \mathfrak{a}_{1_{\varphi(V)}}(A, A)$. We further define the morphism of graded categories $(\varphi, \delta^{\varphi, \mathfrak{a}}) : (\mathcal{V}, \mathfrak{a}^\varphi) \rightarrow (\mathcal{U}, \mathfrak{a})$ for which the maps

$$\mathfrak{a}_V^\varphi \rightarrow \mathfrak{a}_{\varphi(V)}$$

are given by identities and the maps

$$\mathfrak{a}_v^\varphi(A, A') \rightarrow \mathfrak{a}_{\varphi(v)}(A, A'),$$

as well.

Proposition 2.25. (1) The morphisms $(\varphi, \delta^{\varphi, \mathfrak{a}})$ are Cartesian with respect to Ψ_1 .

(2) The category Map is fibered over Cat through Ψ . □

Proposition 2.26. A morphism $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \rightarrow (\mathcal{U}, \mathfrak{a})$ of graded categories is Cartesian with respect to Ψ if and only if the following hold:

- (1) For $V \in \mathcal{V}$, the map $F_V : \mathfrak{b}_V \rightarrow \mathfrak{a}_{\varphi(V)}$ is an isomorphism.
- (2) For $v : V \rightarrow V'$ in \mathcal{V} , $B \in \mathfrak{b}_V$, $B' \in \mathfrak{b}_{V'}$, the map $\mathfrak{b}_v(B, B') \rightarrow \mathfrak{a}_{\varphi(v)}(F(B), F(B'))$ is an isomorphism. □

To distinguish between the notions arising in Propositions 2.23 and 2.26, we make the following definition.

Definition 2.27. Consider a morphism $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ between graded categories.

- (1) (φ, F) is called *Cartesian* if it is Cartesian with respect to Ψ .
- (2) (φ, F) is called *sub-Cartesian* if it is Cartesian with respect to Ψ_1 . □

Remark 2.28. (1) If (φ, F) is Cartesian, then it is sub-Cartesian.

- (2) (φ, F) is sub-Cartesian if and only if $(\varphi^\sharp, F^\sharp)$ as defined in Example 2.17 is Cartesian. □

Example 2.29. Let $\mathfrak{b} \longrightarrow \mathfrak{a}$ be a fully faithful k -linear functor between k -linear categories. With the notation from Example 2.11, the natural graded functor $(e, \mathfrak{b}_{\text{tr}}) \longrightarrow (e, \mathfrak{a}_{\text{tr}})$ is sub-Cartesian and the functor $(\mathcal{U}_{\mathfrak{b}}, \mathfrak{b}_{\text{st}}) \longrightarrow (\mathcal{U}_{\mathfrak{a}}, \mathfrak{a}_{\text{st}})$ is Cartesian (see also Example 2.14). □

2.4 Sites of categories, graded sets, and graded categories

In this section, we will introduce pretopologies on the categories Cat , Mas , and Map . We start by introducing appropriate nerves. For a small category \mathcal{U} , we denote by $\mathcal{N}(\mathcal{U})$ the simplicial nerve of \mathcal{U} . Concretely, for $n \geq 1$, $\mathcal{N}_n(\mathcal{U})$ consists of the n -simplices

$$U_0 \xrightarrow[u_0]{} U_1 \xrightarrow[u_1]{} \cdots \xrightarrow[u_{n-1}]{} U_n ,$$

with $U_i \in \text{Ob}(\mathcal{U})$ and $u_i \in \text{Mor}(\mathcal{U})$, and $\mathcal{N}_0(\mathcal{U}) = \text{Ob}(\mathcal{U})$. Note that $\mathcal{N}_1(\mathcal{U}) = \text{Mor}(\mathcal{U})$. For a functor $\varphi : \mathcal{V} \longrightarrow \mathcal{U}$, we obtain an induced map $\mathcal{N}(\varphi) : \mathcal{N}(\mathcal{V}) \longrightarrow \mathcal{N}(\mathcal{U})$ between the nerves, with components $\mathcal{N}_n(\varphi) : \mathcal{N}_n(\mathcal{V}) \longrightarrow \mathcal{N}_n(\mathcal{U})$.

For a \mathcal{U} -graded category or set \mathfrak{a} , we consider the associated category \mathcal{U}^\sharp from Example 2.13 with $\text{Ob}(\mathcal{U}^\sharp) = \coprod_{U \in \mathcal{U}} \mathfrak{a}_U$ and $\mathcal{U}^\sharp(A_U, A_{U'}) = \mathcal{U}(U, U')$. We define the *nerve* of \mathfrak{a} to be $\mathcal{N}(\mathfrak{a}) = \mathcal{N}(\mathcal{U}, \mathfrak{a}) = \mathcal{N}(\mathcal{U}^\sharp)$. By Example 2.17, a graded map or functor $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ induces an associated map $\mathcal{N}(F) : \mathcal{N}(\mathfrak{b}) \longrightarrow \mathcal{N}(\mathfrak{a})$ given by $\mathcal{N}(\varphi^\sharp) : \mathcal{N}(\mathcal{V}^\sharp) \longrightarrow \mathcal{N}(\mathcal{U}^\sharp)$.

Definition 2.30. Let $n \in \mathbb{N} \cup \{\infty\}$. A collection of functors, graded maps, or graded functors $(F_i : \mathfrak{b}_i \longrightarrow \mathfrak{a})_{i \in I}$ is an n -cover of \mathfrak{a} in Cat , in Mas , or in Map , respectively, if for all $k \in \mathbb{N}$ with $k \leq n$, the collection of maps $(\mathcal{N}_k(F_i))_{i \in I}$ is jointly surjective. □

Thus, a collection of graded maps (respectively, graded functors) $(\varphi_i, F_i) : (\mathcal{V}_i, \mathfrak{b}_i) \longrightarrow (\mathcal{U}, \mathfrak{a})$ is an n -cover of \mathfrak{a} in Mas (respectively, in Map) if and only if the collection of functors $(\varphi^\sharp : \mathcal{V}_i^\sharp \longrightarrow \mathcal{U}^\sharp)_{i \in I}$ is an n -cover in Cat .

Lemma 2.31. Let $n \in \mathbb{N}$. Consider a collection of functors, graded maps, or graded functors, respectively, $S = (F_i : \mathfrak{b}_i \longrightarrow \mathfrak{a})_{i \in I}$. If the collection of maps $(\mathcal{N}_n(F_i))_{i \in I}$ is jointly surjective, then S is an n -cover in Cat , in Mas , or in Map , respectively. \square

Proof. Let $k < n$. Let us look at the graded cases, and let \mathfrak{a} be graded over \mathcal{U} and \mathfrak{b} over \mathcal{V}_i . Every k -simplex u for \mathcal{U}^\sharp gives rise to an n -simplex u' by adding $n - k$ identity maps $1_{U_0} : U_0 \longrightarrow U_0$. If $u' = \varphi_i(v')$ for some i and $v' \in \mathcal{V}_i^\sharp$, then removing the first $n - k$ maps from v' yields a k -simplex v with $\varphi(v) = u$. \blacksquare

Lemma 2.32. Let $n \in \mathbb{N}$. Consider a pullback in Cat , Mas , or Map :

$$\begin{array}{ccc}
 \mathfrak{b}_1 & \xrightarrow{F_1} & \mathfrak{a} \\
 G_1 \uparrow & & \uparrow F_2 \\
 \mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2 & \xrightarrow{G_2} & \mathfrak{b}_2
 \end{array}$$

We have $\mathcal{N}_n(\mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2) \cong \mathcal{N}_n(\mathfrak{b}_1) \times_{\mathcal{N}(\mathfrak{a})} \mathcal{N}(\mathfrak{b}_2)$. \square

Proposition 2.33. Let $n \in \mathbb{N} \cup \{\infty\}$ be fixed. The n -covers from Definition 2.30 define pre-topologies on Cat , Mas , and Map . \square

Proof. The identity and glueing properties are immediate. Concerning the pullback property, we note that in Set , the pullback of a jointly surjective collection of maps is jointly surjective. Hence, the result follows from Example 2.20. \blacksquare

Example 2.34. (1) A functor $\varphi : \mathcal{V} \longrightarrow \mathcal{U}$ which is full and surjective on objects constitutes an ∞ -cover of \mathcal{U} in Cat .

(2) Consider a collection of objects $(U_i)_{i \in I}$ in the category \mathcal{U} and the associated collection of functors $(\varphi_i : \mathcal{U}/U_i \longrightarrow \mathcal{U})_{i \in I}$. The collection $(\varphi_i)_i$ is a 1-cover in

Cat if for every object $U \in \mathcal{U}$, there exists a morphism $U \rightarrow U_i$ for some i . In this case, the collection $(\varphi_i)_i$ is automatically an ∞ -cover of \mathcal{U} . \square

2.5 The stack of map-graded sets

Consider the functor $\Psi_0 : \text{Mas} \rightarrow \text{Cat}$ and consider the pretopology \mathcal{T}_n of n -covers from Proposition 2.33 on Cat for some $n \in \mathbb{N} \cup \{\infty\}$.

Proposition 2.35. The pseudofunctor Mas associated to Ψ_0 is a functor. \square

Let $S = (\varphi_i : \mathcal{V}_i \rightarrow \mathcal{U})_i$ be an n -cover of \mathcal{U} in Cat . We introduce some notation to be able to describe the descent category $\text{Des}(S, \text{Mas})$. We put $\mathcal{V}_{ij} = \mathcal{V}_i \times_{\mathcal{U}} \mathcal{V}_j$ and $\mathcal{V}_{ijk} = \mathcal{V}_i \times_{\mathcal{U}} \mathcal{V}_j \times_{\mathcal{U}} \mathcal{V}_k$. We denote the canonical maps from a k -fold pullback by $\alpha_1, \dots, \alpha_k$ to avoid confusion when some of the indices i, j, \dots coincide. For example, we have $\alpha_1 : \mathcal{V}_{ij} \rightarrow \mathcal{V}_i$ and $\alpha_2 : \mathcal{V}_{ij} \rightarrow \mathcal{V}_j$. For $\mathfrak{b}_i \in \text{Mas}(\mathcal{V}_i)$, we use notation like $\mathfrak{b}_i|_{ij}^1 = \mathfrak{b}_i^{\alpha_1} \in \text{Map}(\mathcal{V}_{ij})$ and $\mathfrak{b}_j|_{ijk}^2 = \mathfrak{b}_j^{\alpha_2} \in \text{Map}(\mathcal{V}_{ijk})$. The descent category $\text{Des}(S, \text{Mas})$ has objects given by $(\mathfrak{b}_i)_i$ with $\mathfrak{b}_i \in \text{Mas}(\mathcal{V}_i)$ along with compatible isomorphisms

$$\rho_{ij} : \mathfrak{b}_i|_{ij}^1 \cong \mathfrak{b}_j|_{ij}^2.$$

Precisely, we require that the cocycle condition

$$(\rho_{jk}|_{ijk}^{23})(\rho_{ij}|_{ijk}^{12}) = \rho_{ik}|_{ijk}^{13} \tag{2}$$

on triple pullbacks holds. Consider $(V_1, V_2, V_3) \in \mathcal{V}_{ijk}$. We have $(\rho_{jk}|_{ijk}^{23})_{(V_1, V_2, V_3)} = (\rho_{jk})_{(V_2, V_3)} : (\mathfrak{b}_j)_{V_2} \rightarrow (\mathfrak{b}_k)_{V_3}$, $(\rho_{ij}|_{ijk}^{12})_{(V_1, V_2, V_3)} = (\rho_{ij})_{(V_1, V_2)} : (\mathfrak{b}_i)_{V_1} \rightarrow (\mathfrak{b}_j)_{V_2}$, $(\rho_{ik}|_{ijk}^{13})_{(V_1, V_2, V_3)} = (\rho_{ik})_{(V_1, V_3)} : (\mathfrak{b}_i)_{V_1} \rightarrow (\mathfrak{b}_k)_{V_3}$. Hence, from (2), we obtain

$$(\rho_{jk})_{(V_2, V_3)}(\rho_{ij})_{(V_1, V_2)} = (\rho_{ik})_{(V_1, V_3)}. \tag{3}$$

In particular, we have

$$(\rho_{ii})_{(V, V)} = 1 : (\mathfrak{b}_i)_V \rightarrow (\mathfrak{b}_i)_V. \tag{4}$$

Theorem 2.36. The functor Mas is a stack. \square

Proof. We may take $n = 0$. Let $S = (\varphi_i : \mathcal{V}_i \rightarrow \mathcal{U})_i$ be a 0-cover of \mathcal{U} in Cat .

First, consider $\mathfrak{a}, \mathfrak{b} \in \text{Mas}(\mathcal{U})$ and a compatible collection of morphism $F_i : \mathfrak{b}^{\varphi_i} \rightarrow \mathfrak{a}^{\varphi_i}$. For the pullback

$$\begin{array}{ccc} \mathcal{V}_i & \xrightarrow{\varphi_i} & \mathcal{U} \\ \alpha_1 \uparrow & & \uparrow \varphi_j \\ \mathcal{V}_{ij} & \xrightarrow{\alpha_2} & \mathcal{V}_j \end{array}$$

we have $(\mathfrak{a}^{\varphi_i})^{\alpha_1} = (\mathfrak{a}^{\varphi_j})^{\alpha_2}$ and similarly for \mathfrak{b} , and compatibility amounts to the fact that $(F_i)^{\alpha_1} = (F_j)^{\alpha_2}$. We are to define a unique glueing $F : \mathfrak{b} \rightarrow \mathfrak{a}$. This consists of maps $F_U : \mathfrak{b}_U \rightarrow \mathfrak{a}_U$ for every $U \in \mathcal{U}$ such that for every i and $V \in \mathcal{V}_i$ we have $(F_i)_V = (F^{\varphi_i})_V = F_{\varphi_i(V)} : \mathfrak{b}_{\varphi_i(V)} \rightarrow \mathfrak{a}_{\varphi_i(V)}$. Since we have a 0-cover, for $U \in \mathcal{U}$ there is some i and some $V \in \mathcal{V}_i$ with $\varphi_i(V) = U$, so we put $F_U = (F_i)_V$. If for another j and $V' \in \mathcal{V}_j$ we also have $\varphi_j(V') = U$, then $(V, V') \in \mathcal{V}_{ij}$ and $(F_i)_V = (F_i)^{\alpha_1}_{(V, V')} = (F_j)^{\alpha_2}_{(V, V')} = (F_j)_{V'}$.

Next we consider a descent datum $(\mathfrak{b}_i)_i$ with $\mathfrak{b}_i \in \text{Mas}(\mathcal{V}_i)$ with compatible isomorphisms $\rho_{ij} : \mathfrak{b}_i|_{\mathcal{V}_{ij}}^1 \cong \mathfrak{b}_j|_{\mathcal{V}_{ij}}^2$. We define the \mathcal{U} -graded set \mathfrak{b} with

$$\mathfrak{b}_U = \coprod_{\varphi_i(V)=U} (\mathfrak{b}_i)_V / \sim,$$

where for $X_1 \in (\mathfrak{b}_i)_{V_1}$ and $X_2 \in (\mathfrak{b}_j)_{V_2}$ we have $X_1 \sim X_2$ if and only if the two elements correspond through the isomorphism $(\rho_{ij})_{(V_1, V_2)} : (\mathfrak{b}_i)_{V_1} \rightarrow (\mathfrak{b}_j)_{V_2}$. The relation \sim is obviously symmetric, it is transitive by (3) and reflexive by (4). For every i and $V \in \mathcal{V}_i$ with $\varphi_i(V) = U$, the canonical map $(\mathfrak{b}_i)_V \rightarrow \mathfrak{b}_U$ is an isomorphism, giving rise to a compatible collection of isomorphisms $F_i : \mathfrak{b}_i \rightarrow \mathfrak{b}^{\varphi_i}$ constituting an isomorphism $F : (\mathfrak{b}_i)_i \rightarrow (\mathfrak{b}^{\varphi_i})_i$ in $\text{Des}(\mathcal{S}, \text{Mas})$. ■

2.6 The stack of map-graded categories

Consider the functor $\Psi_1 : \text{Map} \rightarrow \text{Mas}$ and consider the pretopology \mathcal{T}_n of n -covers from Proposition 2.33 on Mas for some $n \in \mathbb{N} \cup \{\infty\}$.

Proposition 2.37. The pseudofunctor Map associated to Ψ_1 is a functor. □

Let $S = ((\varphi_i, F_i) : (\mathcal{V}_i, \eta_i) \rightarrow (\mathcal{U}, \mathfrak{r}))_i$ be an n -cover of \mathfrak{r} in Mas . We use notation that is similar to the notation introduced in Section 2.5. Pullbacks are denoted by $\eta_{ij} = \eta_i \times_{\mathfrak{r}} \eta_j$ and $\eta_{ijk} = \eta_i \times_{\mathfrak{r}} \eta_j \times_{\mathfrak{r}} \eta_k$ and for $\mathfrak{b}_i \in \text{Map}(\eta_i)$, restrictions are denoted by $\mathfrak{b}_i|_{\mathcal{V}_{ij}}^1 \in \text{Map}(\eta_{ij})$, etc. The descent category $\text{Des}(\mathcal{S}, \text{Map})$ has objects given by $(\mathfrak{b}_i)_i$ with $\mathfrak{b}_i \in \text{Map}(\eta_i)$ along

with compatible isomorphisms

$$\rho_{ij} : \mathfrak{b}_i|_{ij}^1 \cong \mathfrak{b}_j|_{ij}^2,$$

for which the cocycle condition (2) holds. To unravel this condition, we now consider maps $v_\kappa : (V_\kappa, Y_\kappa) \rightarrow (V'_\kappa, Y'_\kappa)$ in $\mathcal{V}_\kappa^\sharp$ for $\kappa \in \{1, 2, 3\}$. Suppose $(v_1, v_2, v_3) \in \eta_{ijk}$. We have $(\rho_{ij})_{v_1, v_2} : (\mathfrak{b}_i)_{v_1}(Y_1, Y'_1) \rightarrow (\mathfrak{b}_j)_{v_2}(Y_2, Y'_2)$, $(\rho_{jk})_{v_2, v_3} : (\mathfrak{b}_j)_{v_2}(Y_2, Y'_2) \rightarrow (\mathfrak{b}_k)_{v_3}(Y_3, Y'_3)$, $(\rho_{ik})_{v_1, v_3} : (\mathfrak{b}_i)_{v_1}(Y_1, Y'_1) \rightarrow (\mathfrak{b}_k)_{v_3}(Y_3, Y'_3)$ and by (2)

$$(\rho_{jk})_{v_2, v_3}(\rho_{ij})_{v_1, v_2} = (\rho_{ik})_{v_1, v_3}. \tag{5}$$

In particular, we have

$$(\rho_{ii})_{v_1, v_1} = 1 : (\mathfrak{b}_i)_{v_1}(Y_1, Y'_1) \rightarrow (\mathfrak{b}_i)_{v_1}(Y_1, Y'_1). \tag{6}$$

Theorem 2.38. (1) If $n \geq 2$, then Map is a prestack.

(2) If $n \geq 3$, then Map is a stack. □

Proof. Let $\mathcal{S} = ((\varphi_i, F_i) : (\mathcal{V}_i, \eta_i) \rightarrow (\mathcal{U}, \mathfrak{r}))_i$ be a 2-cover of \mathfrak{r} in Mas .

First, consider $\mathfrak{a}, \mathfrak{b} \in \text{Map}(\mathfrak{r})$ and a compatible family of morphisms $G_i : \mathfrak{b}|_i \rightarrow \mathfrak{a}|_i$ in $\text{Map}(\eta_i)$. Thus, for every $Y \in (\eta_i)_V$, $Y' \in (\eta_i)_{V'}$, and $v \in \mathcal{V}_i(V, V')$, we have a map $(G_i)_{v, Y, Y'} : (\mathfrak{b}|_i)_v(Y, Y') \rightarrow (\mathfrak{a}|_i)_v(Y, Y')$, that is, a map $(G_i)_{v, Y, Y'} : \mathfrak{b}_{\varphi_i(v)}(F_i(Y), F_i(Y')) \rightarrow \mathfrak{a}_{\varphi_i(v)}(F_i(Y), F_i(Y'))$. To define a unique glueing $G : \mathfrak{b} \rightarrow \mathfrak{a}$ in $\text{Map}(\mathfrak{r})$, we consider $u : U \rightarrow U'$ in \mathcal{U} , $X \in \mathfrak{r}_U$, $X' \in \mathfrak{r}_{U'}$. Since \mathcal{S} is a 1-cover, there is an i and $v : V \rightarrow V'$ in \mathcal{V}_i , $Y \in (\eta_i)_V$, $Y' \in (\eta_i)_{V'}$ with $\varphi_i(v) = u$, $F_i(Y) = X$, $F_i(Y') = X'$. We put

$$G_{u, X, X'} = (G_i)_{v, Y, Y'} : \mathfrak{b}_u(X, X') \rightarrow \mathfrak{a}_u(X, X').$$

Compatibility of the G_i ensures that G is well defined and the fact that \mathcal{S} is a 2-cover can be used to show that G is a \mathcal{U} -graded functor.

Next we consider a descent datum $(\mathfrak{b}_i)_i$ with $\mathfrak{b}_i \in \text{Map}(\eta_i)$ with compatible isomorphisms $\rho_{ij} : \mathfrak{b}_i|_{ij}^1 \cong \mathfrak{b}_j|_{ij}^2$ in $\text{Map}(\eta_{ij})$. We are to define a \mathcal{U} -graded category \mathfrak{b} with underlying graded set \mathfrak{r} . For $u : (U, X) \rightarrow (U', X')$ in \mathcal{U}^\sharp , we first define the set

$$\mathfrak{r}_u(X, X') = \coprod \left[(\mathfrak{b}_i)_v(B, B'), \right]$$

where the coproduct is taken over all $v : (V, B) \rightarrow (V', B')$ in \mathcal{V}_i^\sharp with $\varphi_i(v) = u$, $X = F_i(B)$, and $X' = F_i(B')$. For $w : (W, C) \rightarrow (W', C')$ in \mathcal{V}_j^\sharp with $\varphi_j(w) = u$, $X = F_j(C)$ and $X' = F_j(C')$,

we use the isomorphism

$$(\rho_{ij})_{v,w} : (\mathfrak{b}_i)_v(B, B') \cong (\mathfrak{b}_j)_w(C, C'),$$

to declare when morphisms are equivalent. We thus obtain an equivalence relation on $\mathfrak{r}_u(X, X')$ such that the quotient $\mathfrak{b}_u(X, X')$ has a natural k -module structure, and the canonical morphisms

$$(\mathfrak{b}_i)_v(B, B') \longrightarrow \mathfrak{b}_u(X, X')$$

are k -linear isomorphisms. To define the composition on \mathfrak{b} , we use the fact that S is a 2-cover to choose, for $u : (U, X) \longrightarrow (U', X')$ in \mathcal{U}^\sharp and $u' : (U', X') \longrightarrow (U'', X'')$ a single $(\varphi_i, F_i) : (\mathcal{V}_i, \eta_i) \longrightarrow (\mathcal{U}, \mathfrak{r})$ and $v : (V, B) \longrightarrow (V', B')$, $v' : (V', B') \longrightarrow (V'', B'')$ with $\varphi_i^\sharp(v) = u$, $\varphi_i^\sharp(v') = u'$ and to use the composition

$$(\mathfrak{b}_i)_{v'}(B', B'') \otimes (\mathfrak{b}_i)_v(B, B') \longrightarrow (\mathfrak{b}_i)_{v'v}(B, B'').$$

Finally, to show that the composition is associative, we use the fact that S is a 3-cover. ■

Lemma 2.39. Consider the functor $\Psi_0 : \text{Mas} \longrightarrow \text{Cat}$. On Cat , consider the pretopology of n -covers for some $n \in \mathbb{N} \cup \{\infty\}$. On Mas , consider the pretopology of collections consisting of Cartesian morphisms with respect to Ψ_0 , that are mapped to an n -cover under Ψ_0 (as described in Proposition 2.7). Every cover for this pretopology is an n -cover in the sense of Proposition 2.33. □

Corollary 2.40. Consider the functor $\Psi = \Psi_0 \Psi_1 : \text{Map} \longrightarrow \text{Cat}$ and consider the pretopology of n -covers from Proposition 2.33 on Cat for some $n \in \mathbb{N} \cup \{\infty\}$. The pseudofunctor Map' associated to Ψ is a functor.

- (1) If $n \geq 2$, then Map' is a prestack.
- (2) If $n \geq 3$, then Map' is a stack. □

Proof. This follows from Theorems 2.36, 2.38, Lemma 2.39, and Proposition 2.8. ■

3 Bimodules

In this section, we introduce the bicategory $\underline{\text{Map}}$ of map-graded categories and bimodules between them. We develop the usual machinery of tensor (Section 3.2) and Hom

(Section 3.3) functors in the map-graded context. Some attention is given to the fact that in this context, the notion of bimodule is very natural, while there seems to be no natural notion of module available which does not implicitly or explicitly use bimodules.

3.1 Bifunctors

Let \mathcal{U} and \mathcal{V} be small categories. A \mathcal{U} - \mathcal{V} -bifunctor S is by definition a functor

$$\mathcal{V}^{\text{op}} \times \mathcal{U} \longrightarrow \mathbf{Set} : (V, U) \longmapsto S(V, U).$$

Functoriality translates into the existence of action maps

$$\mathcal{U}(U, U') \times S(V, U) \times \mathcal{V}(V', V) \longrightarrow S(V', U')$$

satisfying the natural associativity and identity axioms. For \mathcal{U} , we have the *identity \mathcal{U} -bifunctor*

$$1_{\mathcal{U}} : \mathcal{U}^{\text{op}} \times \mathcal{U} \longrightarrow \mathbf{Set} : (V, U) \longmapsto \mathcal{U}(V, U).$$

A *morphism* between \mathcal{U} - \mathcal{V} -bifunctors S and S' is a natural transformation $S \longrightarrow S'$. There is a natural category $\mathbf{Bifun}(\mathcal{U}, \mathcal{V})$ of \mathcal{U} - \mathcal{V} -bifunctors and their morphisms.

Example 3.1. (1) Let $\varphi : \mathcal{V} \longrightarrow \mathcal{U}$ be a functor between small categories. There is an associated \mathcal{U} - \mathcal{V} -bifunctor S_{φ} with $S_{\varphi}(V, U) = \mathcal{U}(\varphi(V), U)$.

(2) Let $\varphi : \mathcal{U} \longrightarrow \mathcal{V}$ be a functor between small categories. There is an associated \mathcal{U} - \mathcal{V} -bifunctor S^{φ} with $S^{\varphi}(V, U) = \mathcal{U}(V, \varphi(U))$. \square

Bifunctors can be composed in the following way. Consider an additional small category \mathcal{W} and a \mathcal{V} - \mathcal{W} bifunctor T . We define $S \circ T$ to be the \mathcal{U} - \mathcal{W} -bifunctor with

$$S \circ T(W, U) = \coprod_{V \in \mathcal{V}} S(V, U) \times T(W, V) / \sim,$$

where for $s \in S(V, U)$, $v \in \mathcal{V}(V', V)$, and $t \in T(W, V')$ we have $(sv, t) \sim (s, vt)$. There are canonical isomorphisms $(R \circ S) \circ T \cong R \circ (S \circ T)$, $1_{\mathcal{U}} \circ S \cong S$, and $S \circ 1_{\mathcal{V}} \cong S$. These give rise to a bicategory $\underline{\mathbf{Cat}}$ of categories, bifunctors, and natural transformations.

3.2 Bimodules and tensor functors

Consider a \mathcal{U} -graded category \mathfrak{a} , a \mathcal{V} -graded category \mathfrak{b} , and a \mathcal{U} - \mathcal{V} -bifunctor S . An \mathfrak{a} - S - \mathfrak{b} -bimodule consists of k -modules $M_s(B, A)$ for $s \in S(V, U)$, $B \in \mathfrak{b}_V$, $A \in \mathfrak{a}_U$ with actions

$$\mathfrak{a}_u(A, A') \otimes M_s(B, A) \otimes \mathfrak{b}_v(B', B) \longrightarrow M_{usv}(B', A')$$

satisfying the natural associativity and identity axioms.

Example 3.2. (1) Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be a graded functor. Let S_φ be the \mathcal{U} - \mathcal{V} -bifunctor from Example 3.1(1). For $s \in S_\varphi(V, U) = \mathcal{U}(\varphi(V), U)$ and $B \in \mathfrak{b}_V$, $A \in \mathfrak{a}_U$, we put $(M_F)_s(B, A) = \mathfrak{a}_s(F(B), A)$. This defines an \mathfrak{a} - S_φ - \mathfrak{b} -bimodule M_F .

(2) Let $(\varphi, F) : (\mathcal{U}, \mathfrak{a}) \longrightarrow (\mathcal{V}, \mathfrak{b})$ be a graded functor. Let S^φ be the \mathcal{U} - \mathcal{V} -bifunctor from Example 3.1(2). For $s \in S^\varphi(V, U) = \mathcal{V}(V, \varphi(U))$ and $B \in \mathfrak{b}_V$, $A \in \mathfrak{a}_U$, we put $(M^F)_s(B, A) = \mathfrak{b}_s(B, F(A))$. This defines an \mathfrak{a} - S^φ - \mathfrak{b} -bimodule M^F . \square

The \mathfrak{a} - S - \mathfrak{b} -bimodules form an abelian category $\text{Bimod}_S(\mathfrak{a}, \mathfrak{b})$ with the natural choice of morphisms. If $\mathcal{U} = \mathcal{V}$, we can take $S = 1_{\mathcal{U}}$ and \mathfrak{a} - $1_{\mathcal{U}}$ - \mathfrak{b} -bimodules are simply called \mathfrak{a} - \mathfrak{b} -bimodules. The corresponding category is denoted by $\text{Bimod}_{\mathcal{U}}(\mathfrak{a}, \mathfrak{b})$. As usual, \mathfrak{a} - \mathfrak{a} -bimodules are called \mathfrak{a} -bimodules and the corresponding category is denoted by $\text{Bimod}_{\mathcal{U}}(\mathfrak{a})$. In $\text{Bimod}_{\mathcal{U}}(\mathfrak{a})$, we have the *identity \mathfrak{a} -bimodule* $1_{\mathfrak{a}}$ with $(1_{\mathfrak{a}})_u(A, A') = \mathfrak{a}_u(A, A')$.

Similar to [14, Proposition 2.11], the category $\text{Bimod}_S(\mathfrak{a}, \mathfrak{b})$ can be described as a module category over a linear category. To do so, we define the linear category $\mathfrak{a}^{\text{op}} \otimes_S \mathfrak{b}$ with

$$\text{Ob}(\mathfrak{a}^{\text{op}} \otimes_S \mathfrak{b}) = \coprod_{s \in S(V, U)} \mathfrak{a}_V \times \mathfrak{b}_U$$

and

$$\text{Hom}((s, A, B), (s', A', B')) = \bigoplus_{s' = usv} \mathfrak{a}_u(A, A') \otimes \mathfrak{b}_v(B', B).$$

Proposition 3.3. There is an isomorphism of linear categories

$$\text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \cong \text{Mod}(\mathfrak{a}^{\text{op}} \otimes_S \mathfrak{b}). \quad \square$$

Consider an additional \mathcal{W} -graded category \mathfrak{c} and a \mathcal{V} - \mathcal{W} -bifunctor T . There is a natural tensor product

$$\text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \times \text{Bimod}_T(\mathfrak{b}, \mathfrak{c}) \longrightarrow \text{Bimod}_{S \circ T}(\mathfrak{a}, \mathfrak{c}) : (M, N) \longmapsto M \otimes_{\mathfrak{b}} N,$$

with

$$(M \otimes_b N)_r(C, A) = \bigoplus_{[(s,t)]=r,B} M_s(B, A) \otimes_k N_t(C, B) / \sim,$$

where for $s \in S(V, U)$, $v \in \mathcal{V}(V', V)$, and $t \in T(W, V')$, $B \in \mathfrak{b}_V$, $B' \in \mathfrak{b}_{V'}$, $m \in M_s(B, A)$, $b \in \mathfrak{b}_v(B, B')$, and $n \in N_t(C, B')$ we have $(mb, n) \sim (m, bn)$.

As an application of the tensor product, we obtain tensor actions

$$\otimes_b : \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \times \text{Bimod}_{\mathcal{V}}(\mathfrak{b}) \longrightarrow \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) : (M, X) \longmapsto M \otimes_b X$$

and

$$\otimes_a : \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) \times \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \longrightarrow \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) : (X, M) \longmapsto X \otimes_a M.$$

Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be a graded functor. There is an induced functor

$$F^* : \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \text{Bimod}_{\mathcal{V}}(\mathfrak{b}) : M \longmapsto F^* M,$$

with $(F^* M)_v(B, B') = M_{\varphi(v)}(F(B), F(B'))$.

Example 3.4. For a functor $\varphi : \mathcal{V} \longrightarrow \mathcal{U}$ and a \mathcal{U} -graded \mathfrak{a} , consider the Cartesian φ -graded functor $\delta^{\varphi, \mathfrak{a}} : \mathfrak{a}^{\varphi} \longrightarrow \mathfrak{a}$. We obtain the induced functor

$$(-)^{\varphi} = (\delta^{\varphi, \mathfrak{a}})^* : \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \text{Bimod}_{\mathcal{V}}(\mathfrak{a}^{\varphi}) : M \longmapsto M^{\varphi},$$

with $M_v^{\varphi}(A, A') = M_{\varphi(v)}(A, A')$. □

Letting S vary, we obtain a category $\text{Bimod}(\mathfrak{a}, \mathfrak{b})$ of \mathfrak{a} - \mathfrak{b} bimodules in the following way. An \mathfrak{a} - \mathfrak{b} bimodule consist of a \mathcal{U} - \mathcal{V} -bifunctor S and an \mathfrak{a} - S - \mathfrak{b} -bimodule N . A morphism of bimodules $(\phi, F) : (S, M) \longrightarrow (T, N)$ consists of a natural transformation $\phi : S \longrightarrow T$ of bifunctors and, for every $A \in \mathfrak{a}_{\mathcal{U}}$, $B \in \mathfrak{b}_{\mathcal{V}}$, and $s \in S(V, U)$, a morphism

$$F_{s,B,A} : M_s(B, A) \longrightarrow N_{\phi(s)}(B, A),$$

such that the morphisms $F_{s,B,A}$ are compatible with the actions of \mathfrak{a} and \mathfrak{b} .

Combining the composition of bifunctors and the tensor product of bimodules, we obtain a tensor product

$$\text{Bimod}(\mathfrak{a}, \mathfrak{b}) \times \text{Bimod}(\mathfrak{b}, \mathfrak{c}) \longrightarrow \text{Bimod}(\mathfrak{a}, \mathfrak{c}) : ((S, M), (T, N)) \longmapsto (S \circ T, M \otimes_b N).$$

If we consider a further \mathcal{Z} -graded category \mathfrak{z} and $(R, P) \in \text{Bimod}(\mathfrak{z}, \mathfrak{a})$, then there are natural isomorphisms

$$((R \circ S) \circ T, (P \otimes_{\mathfrak{a}} M) \otimes_{\mathfrak{b}} N) \cong (R \circ (S \circ T), P \otimes_{\mathfrak{a}} (M \otimes_{\mathfrak{b}} N)),$$

and $(1_{\mathcal{U}} \circ S, 1_{\mathfrak{a}} \otimes_{\mathfrak{a}} M) \cong (S, M) \cong (S \circ 1_{\mathcal{V}}, M \otimes_{\mathfrak{b}} 1_{\mathfrak{b}})$. These give rise to a bicategory Map of map-graded categories, bimodules, and bimodule morphisms.

3.3 One-sided bimodules and Hom functors

Bimodules are the natural notion when working with graded categories, but some bimodules can be considered to be more “one-sided” than others. Let \mathcal{U} and \mathcal{V} be categories with a \mathcal{U} - \mathcal{V} -bimodule S . Consider a \mathcal{U} -graded category \mathfrak{a} and a \mathcal{V} -graded category \mathfrak{b} . Furthermore, consider the free \mathcal{U} -graded category $k\mathcal{U}$ and the free \mathcal{V} -graded category $k\mathcal{V}$ as in Example 2.12. Then there are natural bimodule categories

$$\text{Mod}_S(\mathfrak{a}) = \text{Bimod}_S(\mathfrak{a}, k\mathcal{V})$$

and

$$\text{Mod}_S(\mathfrak{b}) = \text{Bimod}_S(k\mathcal{U}, \mathfrak{b}).$$

Example 3.5. If we take, in the first case, $\mathcal{V} = e$, then there is a unique \mathcal{U} - e -bimodule S with $S(*, U) = \{*\}$ for every $U \in \mathcal{U}$. Thus, we obtain the category of left \mathfrak{a} -modules

$$\text{Mod}^l(\mathfrak{a}) = \text{Mod}_S(\mathfrak{a}).$$

Similarly, taking $\mathcal{U} = e$ there is a unique e - \mathcal{V} -bifunctor S with $S(V, *) = \{*\}$ for every $V \in \mathcal{V}$, yielding the category of right \mathfrak{b} -modules

$$\text{Mod}^r(\mathfrak{b}) = \text{Mod}_S(\mathfrak{b}). \quad \square$$

Now we return to the general situation of an underlying \mathcal{U} - \mathcal{V} -bimodule S . Then fixing one argument in an \mathfrak{a} - \mathfrak{b} -bimodule M yields one-sided bimodules in the following sense. Fix $U \in \mathcal{U}$ and $A \in \mathfrak{a}_U$. Then S yields an e - \mathcal{V} -bimodule S_U with

$$S_U(V, *) = S(V, U),$$

and M yields a ke - \mathfrak{b} -bimodule $M_A \in \text{Bimod}_{S_U}(ke, \mathfrak{b})$ with

$$(M_A)_S(B, *) = M_S(B, A).$$

Furthermore, the categories $\text{Bimod}_{S_U}(ke, \mathfrak{b})$ are connected in the following way. Consider a morphism $u: U \rightarrow U'$ in \mathcal{U} . Then there is an associated functor

$$u^* : \text{Bimod}_{S_{U'}}(ke, \mathfrak{b}) \rightarrow \text{Bimod}_{S_U}(ke, \mathfrak{b}) : M \mapsto u^* M,$$

with

$$(u^* M)_S(B, *) = M_{uS}(B, *).$$

Now we are ready to define the Hom functor

$$\text{Hom}_{\mathfrak{b}} : \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \times \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \rightarrow \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) : (M, N) \mapsto \text{Hom}_{\mathfrak{b}}(M, N),$$

where for $u: U \rightarrow U'$ in \mathcal{U} and $A \in \mathfrak{a}_U, A' \in \mathfrak{a}_{U'}$

$$\text{Hom}_{\mathfrak{b}}(M, N)_u(A, A') = \text{Bimod}_{S_U}(ke, \mathfrak{b})(M_A, u^* N_{A'}).$$

There are natural morphisms

$$1_{\mathfrak{a}} \rightarrow \text{Hom}_{\mathfrak{b}}(M, M),$$

given by the natural action maps

$$\mathfrak{a}_u(A, A') \otimes M_S(B, A) \rightarrow M_{uS}(B, A').$$

In a similar way, we define

$$\text{Hom}_{\mathfrak{a}^{\text{op}}} : \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \times \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \rightarrow \text{Bimod}_{\mathcal{V}}(\mathfrak{b}) : (M, N) \mapsto \text{Hom}_{\mathfrak{a}^{\text{op}}}(M, N).$$

Proposition 3.6. (1) The functor $- \otimes_{\mathfrak{a}} M : \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) \rightarrow \text{Bimod}_S(\mathfrak{a}, \mathfrak{b})$ is left adjoint to $\text{Hom}_{\mathfrak{b}}(M, -) : \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \rightarrow \text{Bimod}_{\mathcal{U}}(\mathfrak{a})$.

(2) The functor $M \otimes_{\mathfrak{b}} - : \text{Bimod}_{\mathcal{V}}(\mathfrak{b}) \rightarrow \text{Bimod}_S(\mathfrak{a}, \mathfrak{b})$ is left adjoint to $\text{Hom}_{\mathfrak{a}^{\text{op}}}(M, -) : \text{Bimod}_S(\mathfrak{a}, \mathfrak{b}) \rightarrow \text{Bimod}_{\mathcal{V}}(\mathfrak{b})$. □

4 Functoriality of Map-Graded Hochschild Complexes

The Hochschild complex $\mathbf{C}_{\mathcal{U}}(\mathfrak{a})$ of a map-graded category $(\mathcal{U}, \mathfrak{a})$ is defined in analogy with the Hochschild complex of an algebra [14], naturally making use of the simplicial structure of the nerve $\mathcal{N}(\mathfrak{a}) = \mathcal{N}(\mathcal{U}^{\sharp})$. In this section, we investigate the functoriality properties of map-graded Hochschild complexes. A morphism of map-graded categories is called *sub-Cartesian* if it is Cartesian with respect to $\text{Map} \rightarrow \text{Mas}$. This amounts to the fact that all maps occurring between Hom-modules are isomorphisms. For instance, if we consider a fully faithful functor $\mathfrak{b} \rightarrow \mathfrak{a}$ between k -linear categories as a graded functor between trivially graded categories, it is sub-Cartesian (Example 2.29). In Proposition 4.2, we show that taking Hochschild complexes is functorial with respect to sub-Cartesian functors. This constitutes a natural generalization of the *limited functoriality* of Hochschild complexes of linear categories (see [12] in the differential graded context). Let $\text{Map}_{\text{sc}} \subseteq \text{Map}$ denote the full subcategory of sub-Cartesian morphisms. We endow Map_{sc} with the pretopology of n -covers from Definition 2.30. In Theorem 4.7, we show that the functor

$$\mathbf{C}^n : \text{Map}_{\text{sc}} \longrightarrow \text{Mod}(k) : (\mathcal{U}, \mathfrak{a}) \longmapsto \mathbf{C}_{\mathcal{U}}^n(\mathfrak{a})$$

is a sheaf. As an application of the theorem, in Section 4.5 we obtain a Mayer–Vietoris sequence of Hochschild complexes

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{V}_1}(\mathfrak{a}^{\varphi_1}) \oplus \mathbf{C}_{\mathcal{V}_2}(\mathfrak{a}^{\varphi_2}) \longrightarrow \mathbf{C}_{\mathcal{V}_1 \cap \mathcal{V}_2}(\mathfrak{a}^{\varphi}) \longrightarrow 0,$$

for a map-graded category $(\mathcal{U}, \mathfrak{a})$ and two Cartesian morphisms $(\mathcal{V}_i, \mathfrak{a}^{\varphi_i}) \rightarrow (\mathcal{U}, \mathfrak{a})$ and $(\mathcal{V}_1 \cap \mathcal{V}_2, \mathfrak{a}^{\varphi}) \rightarrow (\mathcal{U}, \mathfrak{a})$ associated to subcategories $\varphi_i : \mathcal{V}_i \subseteq \mathcal{U}$ constituting an n -cover of \mathcal{U} for all $n \geq 0$. Finally, in Section 4.7, we discuss censoring subcategories as a natural generalization of the censoring relations from [15, Section 4.3].

4.1 The map-graded Hochschild complex

Let \mathfrak{a} be a \mathcal{U} -graded category. Let \mathcal{U}^{\sharp} be the category defined in Example 2.13. Recall from Section 2.4 that the nerve of \mathfrak{a} is defined to be the simplicial set $\mathcal{N}(\mathfrak{a}) = \mathcal{N}(\mathcal{U}^{\sharp})$ with n -simplices $\sigma = (u, A)$ given by data

$$\begin{array}{ccccccc} A_0 & & A_1 & & \cdots & & A_n \\ \\ U_0 & \xrightarrow{u_0} & U_1 & \xrightarrow{u_1} & \cdots & \xrightarrow{u_{n-1}} & U_n \end{array}$$

with $u_i \in \mathcal{U}$, $A_i \in \mathfrak{a}_{U_i}$. For $u \in \mathcal{N}(\mathcal{U})_n$, we will use the notation

$$|u| = u_{n-1} \cdots u_1 u_0.$$

A graded functor $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ induces a functor $\mathcal{V}^\sharp \longrightarrow \mathcal{U}^\sharp$ and hence a map $\mathcal{N}(F) : \mathcal{N}(\mathfrak{b}) \longrightarrow \mathcal{N}(\mathfrak{a})$.

Let M be an \mathfrak{a} -bimodule. The *Hochschild complex of \mathfrak{a} with values in M* naturally arises from this simplicial structure as the complex $\mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M)$ with

$$\mathbf{C}_{\mathcal{U}}^n(\mathfrak{a}) = \prod_{(u, A) \in \mathcal{N}(\mathfrak{a})} \text{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), M_{|u|}(A_0, A_n)),$$

with the simplicial Hochschild differential. We put $\mathbf{C}_{\mathcal{U}}(\mathfrak{a}) = \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, 1_{\mathfrak{a}})$. This complex is in fact a B_∞ -algebra [14].

Example 4.1. Let \mathfrak{a} be a linear category. For all the \mathcal{U} -gradings on \mathfrak{a} of Example 2.11, the corresponding Hochschild complexes $\mathbf{C}_{\mathcal{U}}(\mathfrak{a})$ are canonically isomorphic to $\mathbf{C}(\mathfrak{a})$. This results from the fact that all the nerves of these graded categories are canonically isomorphic to $\mathcal{N}(\mathfrak{a})$. □

4.2 Limited functoriality

It is well known that the Hochschild complex of linear categories satisfies so-called “limited functoriality” with respect to inclusions of full subcategories, see [12] for the more general statement for differential graded categories. In this section, we discuss a limited functoriality property for map-graded categories.

Recall that by Proposition 2.23, a graded functor $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ is sub-Cartesian in the sense of Definition 2.27 provided that for every $v : V \longrightarrow V'$ in \mathcal{V} , $B \in \mathfrak{b}_V$, $B' \in \mathfrak{b}_{V'}$, the map

$$\mathfrak{b}_v(B, B') \longrightarrow \mathfrak{a}_{\varphi(v)}(F(B), F(B'))$$

is an isomorphism.

Proposition 4.2. Consider a graded functor $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$. Let M be an \mathfrak{a} -bimodule with induced \mathfrak{b} -bimodule F^*M .

(1) There is a canonical map

$$\prod_{(u,A) \in \mathcal{N}(a)_n} \text{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), M_{|u|}(A_0, A_n))$$

$$\longrightarrow (F_M^*)^n$$

$$\prod_{(v,B) \in \mathcal{N}(b)_n} \text{Hom}_k(\mathfrak{b}_{v_{n-1}}(B_{n-1}, B_n) \otimes \cdots \otimes \mathfrak{b}_{v_0}(B_0, B_1), (F^*M)_{|v|}(B_0, B_n)),$$

given by

$$(F_M^*)^n((\phi_{(u,A)})(u,A)) = (\phi_{(\varphi(v), F(B))} \circ F^{\otimes n})_{(v,B)}.$$

(2) The maps $(F_M^*)^n$ determine a morphism of complexes

$$F_M^* : \mathbf{C}_{\mathcal{U}}(a, M) \longrightarrow \mathbf{C}_{\mathcal{V}}(b, F^*M).$$

(3) If (φ, F) is sub-Cartesian, we have $F^*1_a \cong 1_b$ and the maps $(F_{1_a}^*)^n$ determine a morphism of B_∞ -algebras

$$F^* : \mathbf{C}_{\mathcal{U}}(a) \longrightarrow \mathbf{C}_{\mathcal{V}}(b),$$

with

$$(F^*)^n((\phi_{(u,A)})(u,A)) = (F^{-1} \circ \phi_{(\varphi(v), F(B))} \circ F^{\otimes n})_{(v,B)}. \quad \square$$

Clearly, graded categories with sub-Cartesian graded functors constitute a sub-category $\text{Map}_{\text{sc}} \subseteq \text{Map}$. Let B_∞ denote the category of B_∞ -algebras and morphisms. By Proposition 4.2(3), we obtain a contravariant functor

$$\mathbf{C} : \text{Map}_{\text{sc}} \longrightarrow B_\infty : (\mathcal{U}, a) \longmapsto \mathbf{C}_{\mathcal{U}}(a). \quad (7)$$

Definition 4.3. Let $(\varphi, F) : (\mathcal{V}, b) \longrightarrow (\mathcal{U}, a)$ be a graded functor between graded categories (respectively, a graded map between graded sets). Let $n \in \mathbb{N} \cup \{\infty\}$.

- (1) (φ, F) is *n-surjective* if the canonical $\mathcal{N}_k(F) : \mathcal{N}_k(b) \longrightarrow \mathcal{N}_k(a)$ is surjective for $k \leq n$.
- (2) (φ, F) is *n-injective* if $\mathcal{N}_k(F) : \mathcal{N}_k(b) \longrightarrow \mathcal{N}_k(a)$ is injective for $k \leq n$.
- (3) (φ, F) is *injective* if $\mathcal{N}_1(\varphi) : \mathcal{N}_1(\mathcal{V}) \longrightarrow \mathcal{N}_1(\mathcal{U})$ is injective and every $F : \mathfrak{b}_V \longrightarrow \mathfrak{a}_{\varphi(V)}$ is injective. □

Remark 4.4. (1) (φ, F) is n -surjective if and only if the collection containing (φ, F) as single element is an n -cover in Map (respectively, Mas) in the sense of Definition 2.30.

- (2) If (φ, F) is 1-injective, then it is ∞ -injective.
- (3) If (φ, F) is injective, then it is 1-injective.
- (4) (φ, F) is 1-injective if and only if $(\varphi^\sharp, F^\sharp)$ defined as in Example 2.17 is injective. □

Proposition 4.5. Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \rightarrow (\mathcal{U}, \mathfrak{a})$ be a sub-Cartesian graded functor and let M be an \mathfrak{a} -bimodule. Let $n \in \mathbb{N}$.

- (1) If (φ, F) is n -injective, then $(F^*)^n : \mathbf{C}_{\mathcal{U}}^n(\mathfrak{a}, M) \rightarrow \mathbf{C}_{\mathcal{V}}^n(\mathfrak{b}, F^*M)$ is surjective.
- (2) If (φ, F) is n -surjective, then $(F^*)^n : \mathbf{C}_{\mathcal{U}}^n(\mathfrak{a}, M) \rightarrow \mathbf{C}_{\mathcal{V}}^n(\mathfrak{b}, F^*M)$ is injective. □

Proof. (1) If $\mathcal{N}_n(F) : \mathcal{N}_n(\mathfrak{b}) \rightarrow \mathcal{N}_n(\mathfrak{a})$ is injective, the map $(F^*)^n$ is isomorphic to a projection on a subproduct, whence surjective. (2) Looking at the prescription for $(F^*)^n$, if $\phi \in \mathbf{C}_{\mathcal{U}}^n(\mathfrak{a}, M)$ is such that $\phi_{(\varphi(v), F(B))} = 0$ for every $(v, B) \in \mathcal{N}_n(\mathfrak{b})$, then using n -surjectivity we can write every $(u, A) \in \mathcal{N}(\mathfrak{a})$ as $(u, A) = (\varphi(v), F(B))$ for some (v, B) whence $\phi_{(u,A)} = 0$. ■

Example 4.6. Consider the graded functor $(\varphi, F) : (\mathcal{U}^\sharp, \mathfrak{a}^\sharp) \rightarrow (\mathcal{U}, \mathfrak{a})$ as in Example 2.17. Clearly, (φ, F) is sub-Cartesian and $\mathcal{N}(F) : \mathcal{N}(\mathfrak{a}^\sharp) = \mathcal{N}((\mathcal{U}^\sharp)^\sharp) \cong \mathcal{N}(\mathcal{U}^\sharp) \rightarrow \mathcal{N}(\mathcal{U}^\sharp)$ is an isomorphism. Consequently, $F^* : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \cong \mathbf{C}_{\mathcal{U}^\sharp}(\mathfrak{a}^\sharp)$ is an isomorphism. □

4.3 The sheaf of Hochschild complexes

Consider the presheaf

$$\mathbf{C} : \text{Map}_{\text{sc}} \rightarrow B_\infty : (\mathcal{U}, \mathfrak{a}) \mapsto \mathbf{C}_{\mathcal{U}}(\mathfrak{a}),$$

from (7). Let $n \in \mathbb{N} \cup \{\infty\}$ be fixed. We endow $\text{Map}_{\text{sc}} \subseteq \text{Map}$ with the pretopology of n -covers, as described in Proposition 2.33.

Theorem 4.7. The presheaf $\mathbf{C}^n : \text{Map}_{\text{sc}} \rightarrow \text{Mod}(k) : (\mathcal{U}, \mathfrak{a}) \mapsto \mathbf{C}_{\mathcal{U}}^n(\mathfrak{a})$ is a sheaf. If $n = \infty$, then $\mathbf{C} : \text{Map}_{\text{sc}} \rightarrow B_\infty$ is a sheaf of B_∞ -algebras. □

Proof. Let $((\varphi_i, F_i) : (\mathcal{V}_i, \mathfrak{b}_i) \rightarrow (\mathcal{U}, \mathfrak{a}))_i$ be an n -cover in Map_{sc} . A compatible family of elements for this cover consists of n -cocycles

$$\phi_i \in \mathbf{C}_{\mathcal{V}_i}(\mathfrak{b}_i),$$

such that for every pullback diagram

$$\begin{array}{ccc}
 (\mathcal{V}_i, \mathfrak{b}_i) & \xrightarrow{(\varphi_i, F_i)} & (\mathcal{U}, \mathfrak{a}) \\
 \uparrow (\alpha_1, G_1) & & \uparrow (\varphi_j, F_j) \\
 (\mathcal{V}_i \times_{\mathcal{U}} \mathcal{V}_j, \mathfrak{b}_i \times_{\mathfrak{a}} \mathfrak{b}_j) & \xrightarrow{(\alpha_2, G_2)} & (\mathcal{V}_j, \mathfrak{b}_j)
 \end{array}$$

we have

$$G_i^*(\phi_i) = G_j^*(\phi_j).$$

To define the unique glueing of this family on \mathfrak{a} , we must define for every $(u, A) \in \mathcal{N}_n(\mathfrak{a})$ a corresponding cocycle

$$\phi_{(u, A)} \in \text{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), \mathfrak{a}_{|u|}(A_0, A_n)).$$

Since the collection $(\mathcal{N}_n(F_i))_i$ is jointly surjective, there is an i and $(v, B) \in \mathcal{N}_n(\mathfrak{b}_i)$ for which $\varphi_i(v) = u$, $F_i(B) = A$. We thus have isomorphisms $F_i : (\mathfrak{b}_i)_{v_i}(B_i, B_{i+1}) \rightarrow \mathfrak{a}_{u_i}(A_i, A_{i+1})$ and $F_i : (\mathfrak{b}_i)_{|v|}(B_0, B_n) \rightarrow \mathfrak{a}_{|u|}(A_0, A_n)$. We put

$$\phi_{(u, A)} = F_i \circ \phi_{i(v, B)} \circ (F_i^{-1})^{\otimes n}.$$

It remains to show that this is well defined. Suppose that there is another j and $(w, C) \in \mathcal{N}_n(\mathfrak{b}_j)$ for which $\varphi_j(w) = u$, $F_j(C) = A$. We are to show that $F_i \circ \phi_{i(v, B)} \circ (F_i^{-1})^{\otimes n} = F_j \circ \phi_{j(w, C)} \circ (F_j^{-1})^{\otimes n}$. Now G_1^* maps the collection ϕ_i to a collection ψ_i with

$$\psi_{i((v, w), (B, C))} = (G_1)^{-1} \circ \phi_{i(v, B)} \circ (G_1)^{\otimes n},$$

and G_2^* maps the collection ϕ_j to a collection ψ_j with

$$\psi_{j((v, w), (B, C))} = (G_2)^{-1} \circ \phi_{j(w, C)} \circ (G_2)^{\otimes n}.$$

We thus have

$$F_i \circ \phi_{i(v, B)} \circ (F_i^{-1})^{\otimes n} = (F_i G_1) \circ \psi_{i((v, w), (B, C))} \circ ((F_i G_1)^{-1})^{\otimes n}$$

and

$$F_j \circ \phi_{j(w, C)} \circ (F_j^{-1})^{\otimes n} = (F_j G_2) \circ \psi_{j((v, w), (B, C))} \circ ((F_j G_2)^{-1})^{\otimes n}.$$

By the commutativity of the pullback square and the compatibility assumption $\psi_i = \psi_j$, these two expressions are equal as desired. ■

Let \mathfrak{a} be a \mathcal{U} -graded category with underlying graded set \mathfrak{r} . There are natural functors

$$\Phi_1 : \text{Mas}/\mathfrak{r} \longrightarrow \text{Map} : ((\varphi, F) : (\mathcal{V}, \eta) \rightarrow (\mathcal{U}, \mathfrak{r})) \longmapsto \mathfrak{a}^{(\varphi, F)}$$

and

$$\Phi_2 : \text{Cat}/\mathcal{U} \longrightarrow \text{Map} : (\varphi : \mathcal{V} \rightarrow \mathcal{U}) \longmapsto \mathfrak{a}^\varphi.$$

Let $n \in \mathbb{N} \cup \infty$. Endow Mas/\mathfrak{r} and Cat/\mathcal{U} with the pretopologies of n -covers induced from the ones on Mas and Cat , and also endow Map with the pretopology of n -covers.

Proposition 4.8. The functors Φ_1 and Φ_2 both map covers to covers. □

Corollary 4.9. The presheaves

$$\mathbf{C}_1^n = \mathbf{C}^n \circ \Phi_1 : \text{Mas}/\mathfrak{r} \longrightarrow \text{Mod}(k) : ((\varphi, F) : (\mathcal{V}, \eta) \rightarrow (\mathcal{U}, \mathfrak{r})) \longmapsto \mathbf{C}_{\mathfrak{V}}^n(\mathfrak{a}^{(\varphi, F)})$$

and

$$\mathbf{C}_2^n = \mathbf{C}^n \circ \Phi_2 : \text{Cat}/\mathcal{U} \longrightarrow \text{Mod}(k) : (\varphi : \mathcal{V} \rightarrow \mathcal{U}) \longmapsto \mathbf{C}_{\mathcal{V}}^n(\mathfrak{a}^\varphi)$$

are sheaves. □

We end this section by noting that for a fixed \mathfrak{a} -bimodule M , one similarly has the following proposition.

Proposition 4.10. There are natural sheaves

$$\mathbf{C}_{1,M}^n : \text{Mas}/\mathfrak{r} \longrightarrow \text{Mod}(k) : ((\varphi, F) : (\mathcal{V}, \eta) \rightarrow (\mathcal{U}, \mathfrak{r})) \longmapsto \mathbf{C}_{\mathfrak{V}}^n(\mathfrak{a}^{(\varphi, F)}, F^*M)$$

and

$$\mathbf{C}_{2,M}^n = \mathbf{C}^n \circ \Phi_2 : \text{Cat}/\mathcal{U} \longrightarrow \text{Mod}(k) : (\varphi : \mathcal{V} \rightarrow \mathcal{U}) \longmapsto \mathbf{C}_{\mathcal{V}}^n(\mathfrak{a}^\varphi, M^\varphi). \quad \square$$

Remark 4.11. It is possible to formulate a version of Theorem 4.7 taking map-graded categories endowed with a bimodule as input data for \mathbf{C} , such that Proposition 4.10 is obtained as a corollary. The details are left to the reader. □

4.4 ∞ -Surjections

Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be an ∞ -surjective morphism in Map_{sc} . To formulate the sheaf property of \mathbf{C}^n for the corresponding ∞ -cover, we look at the pullback

$$\begin{array}{ccc} (\mathcal{V}, \mathfrak{b}) & \xrightarrow{(\varphi, F)} & (\mathcal{U}, \mathfrak{a}) \\ (\alpha_1, G_1) \uparrow & & \uparrow (\varphi, F) \\ (\mathcal{V} \times_{\mathcal{U}} \mathcal{V}, \mathfrak{b} \times_{\mathfrak{a}} \mathfrak{b}) & \xrightarrow{(\alpha_2, G_2)} & (\mathcal{V}, \mathfrak{b}) \end{array}$$

According to Theorem 4.7, we obtain an exact sequence of B_∞ -algebras

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \xrightarrow{F^*} \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \xrightarrow{G_2^* - G_1^*} \mathbf{C}_{\mathcal{V} \times_{\mathcal{U}} \mathcal{V}}(\mathfrak{b} \times_{\mathfrak{a}} \mathfrak{b}).$$

Define the complex

$$\mathbf{C}_{\mathcal{V}/\mathcal{U}}(\mathfrak{b}/\mathfrak{a}) = \text{Im}(G_2^* - G_1^*).$$

Then we obtain a long exact cohomology sequence

$$\cdots \longrightarrow HH_{\mathcal{U}}^i(\mathfrak{a}) \xrightarrow{F^*} HH_{\mathcal{V}}^i(\mathfrak{b}) \xrightarrow{G_2^* - G_1^*} HH_{\mathcal{V}/\mathcal{U}}^i(\mathfrak{b}/\mathfrak{a}) \longrightarrow \cdots .$$

4.5 Mayer–Vietoris sequences

Let \mathfrak{a} be a \mathcal{U} -graded category. Consider two sub-Cartesian 1-injections $(\varphi_1, F_1) : (\mathcal{V}_1, \mathfrak{b}_1) \longrightarrow (\mathcal{U}, \mathfrak{a})$ and $(\varphi_2, F_2) : (\mathcal{V}_2, \mathfrak{b}_2) \longrightarrow (\mathcal{U}, \mathfrak{a})$ that together constitute an ∞ -cover of \mathfrak{a} . It is our aim to formulate the sheaf property, so we have to look at all possible pullbacks between (φ_1, F_1) and (φ_2, F_2) . For $i \in \{1, 2\}$, we first look at the pullback

$$\begin{array}{ccc} (\mathcal{V}_i, \mathfrak{b}_i) & \xrightarrow{(\varphi_i, F_i)} & (\mathcal{U}, \mathfrak{a}) \\ (\alpha_1, G_1) \uparrow & & \uparrow (\varphi_i, F_i) \\ (\mathcal{V}_i \times_{\mathcal{U}} \mathcal{V}_i, \mathfrak{b}_i \times_{\mathfrak{a}} \mathfrak{b}_i) & \xrightarrow{(\alpha_2, G_2)} & (\mathcal{V}_i, \mathfrak{b}_i) \end{array}$$

The functor $(-)^{\sharp}$ from Example 2.17 maps this pullback to the pullback of $(\varphi_i^{\sharp}, F_i^{\sharp}) : (\mathcal{V}_i^{\sharp}, \mathfrak{b}_i^{\sharp}) \rightarrow (\mathcal{U}^{\sharp}, \mathfrak{a}^{\sharp})$ with itself. Now $(\varphi_i^{\sharp}, F_i^{\sharp})$ is an injection by Remark 4.4(4), whence it is easily seen to be a monomorphism in Map . Thus, the pullback of this morphism with itself is given by identity morphisms, inducing identity morphisms between Hochschild complexes. Using Example 4.6, we conclude that an element $\phi \in \mathbf{C}_{\mathcal{V}_i}(\mathfrak{b}_i)$ satisfies

$$G_1^*(\phi) = G_2^*(\phi) \in \mathbf{C}_{\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2}(\mathfrak{b}_i \times_{\mathfrak{a}} \mathfrak{b}_i).$$

It thus remains to look at the pullback

$$\begin{array}{ccc} (\mathcal{V}_1, \mathfrak{b}_1) & \xrightarrow{(\varphi_1, F_1)} & (\mathcal{U}, \mathfrak{a}) \\ (\alpha_1, G_1) \uparrow & & \uparrow (\varphi_2, F_2) \\ (\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2, \mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2) & \xrightarrow{(\alpha_2, G_2)} & (\mathcal{V}_2, \mathfrak{b}_2) \end{array}$$

from which we obtain an exact sequence of complexes:

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{V}_1}(\mathfrak{b}_1) \oplus \mathbf{C}_{\mathcal{V}_2}(\mathfrak{b}_2) \longrightarrow \mathbf{C}_{\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2}(\mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2) \longrightarrow 0$$

$$\begin{pmatrix} F_1^* \\ F_2^* \end{pmatrix} \qquad \qquad \qquad \begin{pmatrix} -G_1^* & G_2^* \end{pmatrix}$$

Here, the sequence is left exact by the sheaf property Theorem 4.7 and moreover right exact by Proposition 4.5(2). We thus obtain a long exact cohomology sequence

$$\cdots \longrightarrow HH_{\mathcal{U}}^i(\mathfrak{a}) \longrightarrow HH_{\mathcal{V}_1}^i(\mathfrak{b}_1) \oplus HH_{\mathcal{V}_2}^i(\mathfrak{b}_2) \longrightarrow HH_{\mathcal{V}_1 \times_{\mathcal{U}} \mathcal{V}_2}^i(\mathfrak{b}_1 \times_{\mathfrak{a}} \mathfrak{b}_2) \longrightarrow \cdots$$

$$\begin{pmatrix} F_1^* \\ F_2^* \end{pmatrix} \qquad \qquad \qquad \begin{pmatrix} -G_1^* & G_2^* \end{pmatrix}$$

Example 4.12. Let \mathfrak{a} be a \mathcal{U} -graded category and let $\varphi_1 : \mathcal{V}_1 \subseteq \mathcal{U}$ and $\varphi_2 : \mathcal{V}_2 \subseteq \mathcal{U}$ be subcategories that together constitute an ∞ -cover of \mathcal{U} in Cat . Then the induced Cartesian morphisms $(\mathcal{V}_1, \mathfrak{a}^{\varphi_1}) \rightarrow (\mathcal{U}, \mathfrak{a})$ and $(\mathcal{V}_2, \mathfrak{a}^{\varphi_2}) \rightarrow (\mathcal{U}, \mathfrak{a})$ in Map constitute an ∞ -cover by 1-injections. Put $\varphi : \mathcal{V}_1 \cap \mathcal{V}_2 \subseteq \mathcal{U}$ the inclusion. By Example 2.19, we obtain an exact

sequence of complexes

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{V}_1}(\mathfrak{a}^{\varphi_1}) \oplus \mathbf{C}_{\mathcal{V}_2}(\mathfrak{a}^{\varphi_2}) \longrightarrow \mathbf{C}_{\mathcal{V}_1 \cap \mathcal{V}_2}(\mathfrak{a}^{\varphi}) \longrightarrow 0$$

$$\begin{pmatrix} \varphi_1^* \\ \varphi_2^* \end{pmatrix} \qquad \qquad \qquad \begin{pmatrix} -\alpha_1^* & \alpha_2^* \end{pmatrix}$$

Based upon Proposition 4.10, one obtains a version of this sequence involving bimodules. \square

Example 4.13. Let (X, \mathcal{O}) be a ringed space with an acyclic basis \mathcal{B} of open sets, that is, $H^i(U, \mathcal{O}_U) = 0$ for $i > 0$ and $U \in \mathcal{B}$ (see [15]). Consider open sets U_0, U_1 , and U_2 with $U_0 = U_1 \cup U_2$ and put $U_{12} = U_1 \cap U_2$. Put

$$\mathcal{B}_\star = \{B \in \mathcal{B} \mid B \subseteq U_\star\},$$

for $\star = 1, 2, 12$. Then \mathcal{B}_\star is an acyclic basis for U_\star and $\mathcal{B}_{12} = \mathcal{B}_1 \cap \mathcal{B}_2$. Further, $\mathcal{B}_0 = \mathcal{B}_1 \cup \mathcal{B}_2$ is an acyclic basis for U_0 (which is smaller than the basis U_0 immediately inherits from \mathcal{B}). Now consider all the posets $(\mathcal{B}_\star, \subseteq)$ as small categories. Then we have a cover of \mathcal{B}_0 consisting of the injections $\varphi_i : \mathcal{B}_i \rightarrow \mathcal{B}_0$ for $i = 1, 2$ and we have $\mathcal{B}_{12} = \mathcal{B}_1 \cap \mathcal{B}_2$ as categories. On each of these categories, we consider the corresponding \mathcal{B}_\star -graded category \mathfrak{b}_\star with $(\mathfrak{b}_\star)_V = \{V\}$ and

$$\mathfrak{b}_\star(W, V) = \mathcal{O}(W),$$

for $W \subseteq V$, that is, \mathfrak{b}_\star is the map-graded category associated to the restricted structure sheaf on \mathcal{B}_\star . Clearly, $\mathfrak{b}_1 = (\mathfrak{b}_0)^{\varphi_1}$ and similarly for the other injections. According to [15], $\mathbf{C}_{\mathcal{B}_\star}(\mathfrak{b}_\star)$ computes the Hochschild cohomology of the ringed space $(U_\star, \mathcal{O}|_{U_\star})$. From the above, we obtain an exact sequence

$$0 \longrightarrow \mathbf{C}_{\mathcal{B}_0}(\mathfrak{b}_0) \longrightarrow \mathbf{C}_{\mathcal{B}_1}(\mathfrak{b}_1) \oplus \mathbf{C}_{\mathcal{B}_2}(\mathfrak{b}_2) \longrightarrow \mathbf{C}_{\mathcal{B}_{12}}(\mathfrak{b}_{12}) \longrightarrow 0,$$

and an induced long exact cohomology sequence

$$\cdots \longrightarrow HH^i(U_0) \longrightarrow HH^i(U_1) \oplus HH^i(U_2) \longrightarrow HH^i(U_{12}) \longrightarrow \cdots$$

The more subtle problem of defining a sheaf of Hochschild complexes on quasi-compact opens of a quasi-compact separated scheme was solved in [13]. The existence of Mayer-Vietoris sequences for arbitrary ringed spaces was shown in [15, Section 7.9] making use

of the definition of the Hochschild complex of X as the Hochschild complex of the linear category of injectives in the category of sheaves $\text{Mod}(X)$. We come back to this approach in Example 6.16. \square

4.6 1-Injections

Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be a 1-injective sub-Cartesian graded functor and let M be an \mathfrak{a} -bimodule. We thus have injections $\mathcal{N}_n(\varphi^\sharp) : \mathcal{N}_n(\mathcal{V}^\sharp) \longrightarrow \mathcal{N}_n(\mathcal{U}^\sharp)$ for all $n \in \mathbb{N}$. By Proposition 4.5, we obtain a surjective morphism

$$F^* : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, F^*M).$$

Define the complex

$$\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}^\sharp}(\mathfrak{a}, M) = \text{Ker}(F^*).$$

Then $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}^\sharp}^n(\mathfrak{a}, M)$ is isomorphic to

$$\prod_{(u, A) \notin \text{Im}(\mathcal{N}_n(\varphi^\sharp))} \text{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), M_{|u|}(A_0, A_n)). \tag{8}$$

Proposition 4.14. The following are equivalent:

- (1) $(u, A) \in \text{Im}(\mathcal{N}_n(\varphi^\sharp))$.
- (2) For every $u_i : A_i \longrightarrow A_{i+1}$ in $\mathcal{N}_1(\mathcal{U}^\sharp)$ occurring in (u, A) , we have $u_i \in \text{Im}(\mathcal{N}_1(\varphi^\sharp))$.

Proof. Suppose that (2) holds. Let $u_i = \varphi^\sharp(v) : A_i \longrightarrow A_{i+1}$ for $v : B \longrightarrow B'$ in $\mathcal{N}_1(\mathcal{V}^\sharp)$ and let $u_{i+1} = \varphi^\sharp(w) : A_{i+1} \longrightarrow A_{i+2}$ for $w : C \longrightarrow C'$ in $\mathcal{N}_1(\mathcal{V}^\sharp)$. Then $\varphi^\sharp(1_{B'}) = 1_{A_{i+1}} = \varphi^\sharp(1_C)$ whence $B' = C$. \blacksquare

We call $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}^\sharp}(\mathfrak{a}, M)$ the *Hochschild complex of \mathfrak{a} with support outside \mathcal{V}^\sharp (and values in M)*. We thus obtain an exact sequence of complexes

$$0 \longrightarrow \mathbf{C}_{\mathcal{U} \setminus \mathcal{V}^\sharp}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, F^*M) \longrightarrow 0, \tag{9}$$

and a long exact cohomology sequence

$$\cdots \longrightarrow HH_{\mathcal{U} \setminus \mathcal{V}^\sharp}^i(\mathfrak{a}, M) \longrightarrow HH_{\mathcal{U}}^i(\mathfrak{a}, M) \xrightarrow{F^*} HH^i(\mathfrak{b}, F^*M) \longrightarrow \cdots$$

Example 4.15. Let \mathfrak{a} be a \mathcal{U} -graded category and $\varphi : \mathcal{V} \subseteq \mathcal{U}$ be a subcategory. Consider the Cartesian morphism $\delta^{\varphi, \mathfrak{a}} : \mathfrak{a}^\varphi \longrightarrow \mathfrak{a}$. Then $(\varphi, \delta^{\varphi, \mathfrak{a}}) : (\mathcal{V}, \mathfrak{a}^\varphi) \longrightarrow (\mathcal{U}, \mathfrak{a})$ is injective whence 1-injective. In this case, the following are equivalent:

- (1) $(u, A) \in \text{Im}(\mathcal{N}_n(\varphi^\sharp))$.
- (2) For every $u_i : U_i \longrightarrow U_{i+1}$ in $\mathcal{N}_1(\mathcal{U})$ occurring in u , we have $u \in \text{Im}(\mathcal{N}_1(\varphi))$.

Consequently, we will denote $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a}, M) = \mathbf{C}_{\mathcal{U} \setminus \mathcal{V}^\sharp}(\mathfrak{a}, M)$ and call this complex the *Hochschild complex of \mathfrak{a} with support outside \mathcal{V}* . □

4.7 Censoring subcategories

Let \mathfrak{a} be a \mathcal{U} -graded category. A subcategory $\mathcal{V} \subseteq \mathcal{U}$ is called *censoring* if for all $u : U \longrightarrow U', A \in \mathfrak{a}_U, A' \in \mathfrak{a}_{U'}$ with $u \notin \mathcal{V}$, we have $\mathfrak{a}_u(A, A') = 0$. The terminology is taken from [15, Section 4.3], cf. the following example.

Example 4.16. Let \mathfrak{a} be a linear category and let \mathcal{R} be a transitive relation on $\text{Ob}(\mathfrak{a})$. In [15], the relation \mathcal{R} is called *censoring* if $\mathfrak{a}(B, A) = 0$ for $(B, A) \notin \mathcal{R}$. Clearly, this yields a special case of a censoring subcategory $\mathcal{U}_{\mathcal{R}}$ of the standard grading category \mathcal{U} of \mathfrak{a} . Precisely, we let $\mathcal{U}_{\mathcal{R}}$ be the category with $\text{Ob}(\mathcal{U}_{\mathcal{R}}) = \text{Ob}(\mathfrak{a})$ and

$$\mathcal{U}_{\mathcal{R}}(B, A) = \begin{cases} * & \text{if } (B, A) \in \mathcal{R}, \\ \emptyset & \text{otherwise.} \end{cases} \quad \square$$

A 1-injection $(\varphi, F) : (\mathcal{V}, \mathfrak{a}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ of graded sets is called *censoring* if for all $u : A \longrightarrow A'$ in \mathcal{U}^\sharp which is not in the image of $\mathcal{N}_1(F) : \mathcal{N}_1(\mathfrak{a}) \longrightarrow \mathcal{N}_1(\mathfrak{a})$, we have $\mathfrak{a}_u(A, A') = 0$.

Remark 4.17. (1) A subcategory $\varphi : \mathcal{V} \subseteq \mathcal{U}$ is censoring if and only if the Cartesian map $(\varphi, \delta^{\varphi, \mathfrak{a}}) : (\mathcal{V}, \mathfrak{a}^\varphi) \longrightarrow (\mathcal{U}, \mathfrak{a})$ is a censoring 1-injection.

- (2) A 1-injection $(\varphi, F) : (\mathcal{V}, \mathfrak{a}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ is censoring if and only if $\varphi^\sharp : \mathcal{V}^\sharp \longrightarrow \mathcal{U}^\sharp$ defines a censoring subcategory for the graded category $(\mathcal{U}^\sharp, \mathfrak{a}^\sharp)$. □

Proposition 4.18. Let \mathfrak{a} be a \mathcal{U} -graded category and let M be an \mathfrak{a} -bimodule.

- (1) Let $\varphi : \mathcal{V} \subseteq \mathcal{U}$ be a censoring subcategory. Then the map

$$\varphi^* : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{a}^\varphi, M^\varphi)$$

is an isomorphism of complexes.

- (2) Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be a sub-Cartesian, censoring 1-injection. Then the map

$$F^* : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, F^*M)$$

is an isomorphism of complexes. □

Proof. By the higher remarks, it suffices to prove either one of (1), (2). Let us prove (2). By the assumption, every component

$$\mathrm{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), M_{|\mathcal{U}}(A_0, A_n))$$

in (8) contains some $u_i : A_i \longrightarrow A_{i+1}$ not contained in $\mathrm{Im}(\mathcal{N}_1(\varphi^\sharp))$, which thus has $\mathfrak{a}_{u_i}(A_i, A_{i+1}) = 0$. Consequently, $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}^\sharp}(\mathfrak{a}, M) = 0$. ■

Proposition 4.18 has a useful corollary, which says that a censoring subcategory essentially censors possible uncontrollable parts of bimodules. This was precisely the original intuition behind the terminology in [15].

Corollary 4.19. Let \mathfrak{a} be a \mathcal{U} -graded category and let $\mathcal{V} \subseteq \mathcal{U}$ be a censoring subcategory. Let

$$f : M \longrightarrow N$$

be a morphism of \mathfrak{a} -bimodules. If

$$f^\varphi : M^\varphi \longrightarrow N^\varphi$$

is a quasi-isomorphism, then so is

$$\mathbf{C}_{\mathcal{U}}(\mathfrak{a}, f) : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, N). \quad \square$$

5 Hochschild Cohomology with Support

Let \mathfrak{a} be a \mathcal{U} -graded category, $\varphi : \mathcal{V} \subseteq \mathcal{U}$ be a subcategory, and $(\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be a Cartesian functor. In this section, we investigate some cases where the cohomology of the complex $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a}, M)$ (see Example 4.15) has a nice cohomological interpretation. The main point is that we need some control over the “complement” of \mathcal{V} in \mathcal{U} . Precisely, we assume that this complement (the \mathcal{U} -morphisms not in \mathcal{V}) constitutes an ideal \mathcal{Z} in

\mathcal{U} . In this case, we show in Proposition 5.3 that

$$\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a}, M) \cong \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{Z}}),$$

where $M_{\mathcal{Z}}$ is the natural restriction of M to an \mathfrak{a} -bimodule supported on \mathcal{Z} (i.e., with zero values outside of \mathcal{Z}). Our setup applies in the situation where \mathcal{U} is the category associated to a collection of open subsets of a topological space X ordered by inclusion, \mathcal{V} is the full subcategory of subsets $U \subseteq V$ for a fixed subset V , and \mathcal{Z} contains the inclusions $U' \subseteq U$ with $U \not\subseteq V$.

In Section 5.7, we revisit the arrow category construction from [12] in the map-graded context. For an $(\mathcal{U}, \mathfrak{a})$ - $(\mathcal{V}, \mathfrak{b})$ -bimodule (S, M) , we take the natural inclusion $\mathcal{V} \coprod \mathcal{U} \rightarrow (\mathcal{V} \rightarrow_S \mathcal{U})$ and corresponding Cartesian functor

$$(\mathcal{V} \coprod \mathcal{U}, \mathfrak{b} \coprod \mathfrak{a}) \rightarrow (\mathcal{V} \rightarrow_S \mathcal{U}, \mathfrak{b} \rightarrow_M \mathfrak{a}),$$

as starting point for obtaining map-graded analogs of some of the main results from [12]. Sections 5.9 and 5.10 are entirely modeled upon the treatment in [12], and mainly formulate results from [12] in the map-graded context, making use of the natural Hom and tensor functors from Section 3.2. Further, in Section 5.8, we give an intrinsic characterization of arrow categories based upon the *thin ideals* introduced in Section 5.6.

5.1 Ideals in categories

Let \mathcal{U} be an arbitrary category. Recall from Section 3.1 that a \mathcal{U} -bifunctor S consists of sets $S(V, U)$ for $U, V \in \mathcal{U}$ and actions

$$\mathcal{U}(U, U') \times S(V, U) \times \mathcal{U}(V', V) \rightarrow S(V', U').$$

Denote the category of \mathcal{U} -bimodules by $\text{Bimod}(\mathcal{U})$. A (*two sided*) *ideal* in \mathcal{U} is a subfunctor

$$\mathcal{Z} \subseteq \mathbf{1}_{\mathcal{U}}$$

of the identity bifunctor in the category $\text{Bifun}(\mathcal{U})$ of \mathcal{U} -bifunctors. More precisely, it consists of subsets $\mathcal{Z}(V, U) \subseteq \mathcal{U}(V, U)$ for all $U, V \in \mathcal{U}$ such that the composition of \mathcal{U} restricts to

$$\mathcal{U}(U, U') \times \mathcal{Z}(V, U) \times \mathcal{U}(V', V) \rightarrow \mathcal{Z}(V', U').$$

We put the *morphisms in* \mathcal{Z} equal to $\text{Mor}(\mathcal{Z}) = \coprod_{V, U} \mathcal{Z}(V, U)$.

5.2 Bimodules over ideals

Let \mathfrak{a} be a \mathcal{U} -graded category and let \mathcal{Z} be an ideal in \mathcal{U} . An \mathfrak{a} -bimodule M on \mathcal{Z} consists of:

- (1) k -modules $M_{\mathcal{Z}}(B, A)$ for all $\mathcal{Z}: V \rightarrow U$ in \mathcal{Z} and $A \in \mathfrak{a}_U, B \in \mathfrak{a}_V$;
- (2) actions $\mathfrak{a}_U(A, A') \otimes M_{\mathcal{Z}}(B, A) \otimes \mathfrak{a}_V(B', B) \rightarrow M_{\mathcal{U}\mathcal{Z}V}(B', A')$ for all additional $u: U \rightarrow U', v: V' \rightarrow V$ in \mathcal{U} and $A' \in \mathfrak{a}_{U'}, B' \in \mathfrak{a}_{V'}$;

with the natural axioms. The \mathfrak{a} -bimodules on \mathcal{Z} form an abelian category $\text{Bimod}_{\mathcal{Z}}(\mathfrak{a})$. There are obvious exact functors

$$(-)|^{\mathcal{U}}: \text{Bimod}_{\mathcal{Z}}(\mathfrak{a}) \rightarrow \text{Bimod}_{\mathcal{U}}(\mathfrak{a}): M \mapsto M|^{\mathcal{U}},$$

where $M|^{\mathcal{U}}$ is the extension of M by zero values outside of \mathcal{Z} , and

$$(-)|_{\mathcal{Z}}: \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) \rightarrow \text{Bimod}_{\mathcal{Z}}(\mathfrak{a}): M \mapsto M|_{\mathcal{Z}},$$

where $M|_{\mathcal{Z}}$ is the restriction of M to \mathcal{Z} . Clearly, $(-)|_{\mathcal{Z}}$ is right adjoint to $(-)|^{\mathcal{U}}$,

$$(-)|_{\mathcal{Z}}(-)|^{\mathcal{U}} = \mathbf{1}_{\text{Bimod}_{\mathcal{Z}}(\mathfrak{a})},$$

and the canonical

$$(-)_{\mathcal{Z}} = (-)|^{\mathcal{U}}(-)|_{\mathcal{Z}} \rightarrow \mathbf{1}_{\text{Bimod}_{\mathcal{U}}(\mathfrak{a})}$$

corresponds to inclusions of \mathfrak{a} -modules

$$M_{\mathcal{Z}} \subseteq M,$$

where $M_{\mathcal{Z}}$ is obtained by changing the values of M to zero outside of \mathcal{Z} . We can thus identify $\text{Bimod}_{\mathcal{Z}}(\mathfrak{a})$ with the full subcategory of $\text{Bimod}_{\mathcal{U}}(\mathfrak{a})$ of bimodules *supported on* \mathcal{Z} .

5.3 Ideal-subcategory decomposition

We now turn to our main situation of interest, which is summarized as follows:

- (1) \mathcal{U} is an arbitrary category;
- (2) $\mathcal{V} \subseteq \mathcal{U}$ is a subcategory;
- (3) $\mathcal{Z} \subseteq \mathbf{1}_{\mathcal{U}}$ is an ideal;
- (4) $\text{Mor}(\mathcal{U}) = \text{Mor}(\mathcal{V}) \coprod \text{Mor}(\mathcal{Z})$.

In this case, we call $(\mathcal{Z}, \mathcal{V})$ an *ideal-subcategory decomposition* of \mathcal{U} .

Remark 5.1. Note that the datum of an ideal-subcategory decomposition $(\mathcal{Z}, \mathcal{V})$ of \mathcal{U} is in one-to-one correspondence with the datum of an ideal $\mathcal{Z} \subseteq 1_{\mathcal{U}}$ which is completely prime, that is, $zz' \in \mathcal{Z}$ implies $z \in \mathcal{Z}$ or $z' \in \mathcal{Z}$. Indeed, to recover \mathcal{V} from \mathcal{Z} it suffices to put $\text{Ob}(\mathcal{V}) = \{U \in \mathcal{U} \mid 1_U \notin \mathcal{Z}\}$ and $\text{Mor}(\mathcal{V}) = \text{Mor}(\mathcal{U}) \setminus \text{Mor}(\mathcal{Z})$. \square

Example 5.2. Let X be a topological space, V be an open subset and $Z = X \setminus V$ be its closed complement. Let $\mathcal{U} \subseteq \text{open}(X)$ be a subposet of the open sets of X . Let $\mathcal{V} \subseteq \mathcal{U}$ be the full subcategory with

$$\text{Ob}(\mathcal{V}) = \{U \in \mathcal{U} \mid U \subseteq V\},$$

and $\mathcal{Z} \subseteq 1_{\mathcal{U}}$ the ideal with $u: U \rightarrow U'$ in \mathcal{Z} if and only if $U' \not\subseteq V$. Clearly, $(\mathcal{Z}, \mathcal{V})$ is an ideal-subcategory decomposition of \mathcal{U} . \square

5.4 The localization sequence of bimodule categories

Let \mathfrak{a} be a \mathcal{U} -graded category and $(\mathcal{Z}, \mathcal{V})$ be an ideal-subcategory decomposition of \mathcal{U} . Put $\mathfrak{b} = \mathfrak{a}|_{\mathcal{V}}$. In this section, we take a closer look at the sequence

$$\text{Bimod}_{\mathcal{Z}}(\mathfrak{a}) \xrightarrow{(-)|_{\mathcal{Z}}^{\mathcal{U}}} \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) \xrightarrow{(-)|_{\mathcal{V}}} \text{Bimod}_{\mathcal{V}}(\mathfrak{b}).$$

Clearly, $\text{Bimod}_{\mathcal{Z}}(\mathfrak{a})$ is the kernel of $(-)|_{\mathcal{V}}$, and we already know from Section 5.2 that $(-)|_{\mathcal{Z}}^{\mathcal{U}}$ has a right adjoint $(-)|_{\mathcal{Z}}$. Since $(-)|_{\mathcal{V}}$ is induced by an underlying k -linear functor $\mathfrak{b} \otimes_{\mathcal{V}} \mathfrak{b} \rightarrow \mathfrak{a} \otimes_{\mathcal{U}} \mathfrak{a}$, both adjoints of $(-)|_{\mathcal{V}}$ exist and can be described explicitly. Moreover, since \mathcal{Z} is an ideal, the right adjoint has a particularly easy description. Define the functor

$$(-)|_{\mathcal{V}}^{\mathcal{U}} : \text{Bimod}_{\mathcal{V}}(\mathfrak{b}) \rightarrow \text{Bimod}_{\mathcal{U}}(\mathfrak{a}) : M \mapsto M|^{\mathcal{U}},$$

in the following way. Put

$$(M|^{\mathcal{U}})_u(B, A) = \begin{cases} M_u(B, A) & \text{if } u \in \mathcal{V}, \\ 0 & \text{otherwise.} \end{cases}$$

Then there is a unique way to let \mathfrak{a} act on $M|^{\mathcal{U}}$ so that

$$\mathfrak{a}_u(A, A') \otimes (M|^{\mathcal{U}})_u(B, A) \otimes \mathfrak{a}_{u'}(B', B) \rightarrow (M|^{\mathcal{U}})_{uu'}(B', A')$$

is given by the action of \mathfrak{b} on M if u, u', u'' are in \mathcal{V} , and is zero otherwise. This is a well-defined action thanks to the fact that \mathcal{Z} is an ideal. The functor $(-)|_{\mathcal{V}}^{\mathcal{U}}$ is right adjoint to $(-)|_{\mathcal{V}}$ and we have

$$(-)|_{\mathcal{V}}(-)|_{\mathcal{V}}^{\mathcal{U}} = \mathbf{1}_{\text{Bimod}_{\mathcal{V}}(\mathfrak{b})}.$$

The canonical

$$\mathbf{1}_{\text{Bimod}_{\mathcal{U}}(\mathfrak{a})} \longrightarrow (-)|_{\mathcal{V}}^{\mathcal{U}}(-)|_{\mathcal{V}} = (-)|_{\mathcal{V}}$$

corresponds to quotients of \mathfrak{a} -bimodules

$$M \longrightarrow M_{\mathcal{V}},$$

where $M_{\mathcal{V}}$ is obtained from M by changing the values of M outside \mathcal{V} to zero.

Summarizing, we have an exact sequence of functors

$$0 \longrightarrow (-)|_{\mathcal{Z}} \longrightarrow \mathbf{1}_{\text{Bimod}_{\mathcal{U}}(\mathfrak{a})} \longrightarrow (-)|_{\mathcal{V}} \longrightarrow 0,$$

corresponding to exact sequences of bimodules

$$0 \longrightarrow M_{\mathcal{Z}} \longrightarrow M \longrightarrow M_{\mathcal{V}} \longrightarrow 0. \tag{10}$$

5.5 The localization sequence for Hochschild cohomology

We can now use (10) to obtain the following short exact sequence of Hochschild complexes:

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{Z}}) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{V}}) \longrightarrow 0. \tag{11}$$

Proposition 5.3. The sequences (9) and (11) are canonically isomorphic. □

Proof. This follows from the fact that $(\mathcal{Z}, \mathcal{V})$ is an ideal-subcategory decomposition of \mathcal{U} . Indeed, let us first compare $\mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{V}})$ to $\mathbf{C}_{\mathcal{V}}(\mathfrak{b}, M|_{\mathcal{V}})$. In every product

$$\prod_{(u, A) \in \mathcal{N}(\mathfrak{a})} \text{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), (M_{\mathcal{V}})|_{\mathcal{U}}(A_0, A_n)),$$

turning up in $\mathbf{C}_{\mathcal{U}}^n(\mathfrak{a}, M_{\mathcal{V}})$ we may clearly remove the pieces with $|u| \in \mathcal{Z}$ since then $(M_{\mathcal{V}})|_{\mathcal{U}}(A_0, A_n) = 0$. But since \mathcal{Z} is an ideal and \mathcal{V} a subcategory, the remaining pieces

are precisely the ones with all the $u_i \in \mathcal{V}$, and we recover $\mathbf{C}_{\mathcal{V}}(\mathfrak{b}, M|_{\mathcal{V}})$. Similarly, consider a product

$$\prod_{(u, A) \in \mathcal{N}(\mathfrak{a})} \mathrm{Hom}_k(\mathfrak{a}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), (M_{\mathcal{Z}})_{|u|}(A_0, A_n))$$

in $\mathbf{C}_{\mathcal{U}}^n(\mathfrak{a}, M_{\mathcal{Z}})$. This time, all the pieces with $|u| \in \mathcal{V}$ can be removed. What remains are the pieces in which at least one u_i belongs to \mathcal{Z} , which corresponds precisely to $\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a}, M)$. ■

In fact, the isomorphism $\mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{V}}) \cong \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, M|_{\mathcal{V}})$ is easily understood on the derived level. Indeed, the localization between $\mathrm{Bimod}_{\mathcal{U}}(\mathfrak{a})$ and $\mathrm{Bimod}_{\mathcal{V}}(\mathfrak{b})$ of Section 5.4 yields:

$$\mathrm{RHom}_{\mathrm{Bimod}_{\mathcal{U}}(\mathfrak{a})}(1_{\mathfrak{a}}, (M|_{\mathcal{V}})|^{\mathcal{U}}) \cong \mathrm{RHom}_{\mathrm{Bimod}_{\mathcal{V}}(\mathfrak{b})}(1_{\mathfrak{b}}, M|_{\mathcal{V}}).$$

The more mysterious part in the isomorphic sequences (9) and (11) remains the Hochschild complex with support

$$\mathbf{C}_{\mathcal{U} \setminus \mathcal{V}}(\mathfrak{a}, M) \cong \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{Z}}),$$

but at least it now has an interpretation as an ordinary Hochschild complex of \mathfrak{a} with values in the bimodule $M_{\mathcal{Z}}$ supported on \mathcal{Z} . So far, the most meaningful incarnation of the sequences (9) and (11) is perhaps

$$0 \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M_{\mathcal{Z}}) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, M) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, M|_{\mathcal{V}}) \longrightarrow 0. \tag{12}$$

5.6 Thin ideals

Consider $\varphi : \mathcal{V} \subseteq \mathcal{U}$ with $\mathrm{Ob}(\mathcal{V}) = \mathrm{Ob}(\mathcal{U})$ and let $(\mathcal{Z}, \mathcal{V})$ be an ideal-subcategory decomposition of \mathcal{U} . Let \mathfrak{b} be a \mathcal{V} -graded category. We define a \mathfrak{b} -bimodule on \mathcal{Z} to consist of the following data:

- (1) for all $z : V \longrightarrow U$ in \mathcal{Z} and $B \in \mathfrak{b}_V, A \in \mathfrak{b}_U$, a k -module $M_z(B, A)$;
- (2) actions $\mathfrak{b}_v(A, A') \otimes M_z(B, A) \otimes \mathfrak{b}_{v'}(B', B) \longrightarrow M_{vzv'}(B', A')$ for all additional $v : U \longrightarrow U', v' : V' \longrightarrow V$ in \mathcal{V} and $A' \in \mathfrak{b}_{U'}, B' \in \mathfrak{b}_{V'}$.

We thus obtain the abelian category $\mathrm{Bimod}_{\mathcal{Z}}(\mathfrak{b})$ of \mathfrak{b} -bimodules on \mathcal{Z} .

Now consider a \mathcal{U} -graded category \mathfrak{a} and suppose $\mathfrak{b} = \mathfrak{a}|_{\mathcal{V}}$. There is a functor

$$(-)|^{\mathfrak{a}} : \text{Bimod}_{\mathcal{Z}}(\mathfrak{b}) \longrightarrow \text{Bimod}_{\mathcal{Z}}(\mathfrak{a}) : M \longmapsto M|^{\mathfrak{a}},$$

where the actions $\alpha_{z'}(A, A') \otimes M_z(B, A) \longrightarrow M_{zz'}(B, A')$ and $M_z(B, A) \otimes \alpha_{z'}(B', B) \longrightarrow M_{zz'}(B', A)$ are zero for $z, z',$ and z' in \mathcal{Z} . This yields a well-defined action since \mathcal{Z} is an ideal. Obviously, there is also a functor

$$(-)|_{\mathfrak{b}} : \text{Bimod}_{\mathcal{Z}}(\mathfrak{a}) \longrightarrow \text{Bimod}_{\mathcal{Z}}(\mathfrak{b}) : M \longmapsto M|_{\mathfrak{b}},$$

which restricts the action to \mathfrak{b} . Clearly,

$$(-)|_{\mathfrak{b}}(-)|^{\mathfrak{a}} = 1_{\text{Bimod}_{\mathcal{Z}}(\mathfrak{b})}.$$

Definition 5.4. (1) An ideal $\mathcal{Z} \subseteq 1_{\mathcal{U}}$ is called *thin* if it contains no consecutive morphisms, that is, if we consider $z \in \mathcal{Z}(V, U), u' \in \mathcal{U}(V', V), u \in \mathcal{U}(U, U')$, then it follows that $u' \notin \mathcal{Z}$ and $u \notin \mathcal{Z}$.

(2) An \mathfrak{a} -bimodule M on \mathcal{Z} is called \mathcal{Z} -*thin* if it receives no nontrivial actions from \mathcal{Z} , that is, if we have $z \in \mathcal{Z}(V, U), z' \in \mathcal{Z}(V', V), z'' \in \mathcal{Z}(U, U')$, then it follows that both $\alpha_{z'}(A, A') \otimes M_z(B, A) \longrightarrow M_{z''z'}(B, A')$ and $M_z(B, A) \otimes \alpha_{z''}(B', B) \longrightarrow M_{zz''}(B', A)$ are zero for all A, A', B, B' . □

Example 5.5. Let \mathcal{U} be a category and $\text{Ob}_1 \subseteq \text{Ob}(\mathcal{U})$ and $\text{Ob}_2 \subseteq \text{Ob}(\mathcal{U})$ be two classes of objects with no morphisms going from Ob_2 to Ob_1 . Then the \mathcal{U} -morphisms starting in Ob_1 and landing in Ob_2 form a thin ideal in \mathcal{U} . □

Obviously, if \mathcal{Z} is thin, then every M -bimodule on \mathcal{Z} is \mathcal{Z} -thin. In general, we make the following easy observations.

Proposition 5.6. For an \mathfrak{a} -bimodule M on \mathcal{Z} , the following are equivalent:

- (1) M is \mathcal{Z} -thin;
- (2) $M \cong (M|_{\mathfrak{b}})|^{\mathfrak{a}}$;
- (3) M is in the image of $(-)|^{\mathfrak{a}}$. □

Corollary 5.7. If \mathcal{Z} is thin, then $(-)|^{\mathfrak{a}}$ and $(-)|_{\mathfrak{b}}$ constitute inverse isomorphisms

$$\text{Bimod}_{\mathcal{Z}}(\mathfrak{b}) \cong \text{Bimod}_{\mathcal{Z}}(\mathfrak{a}). \quad \square$$

5.7 Arrow categories

In this section, we introduce the arrow category construction from [12] in the map-graded setting. Let \mathfrak{a} be a \mathcal{U} -graded category, \mathfrak{b} be a \mathcal{V} -graded category, S be a \mathcal{U} - \mathcal{V} -bifunctor, and M be an \mathfrak{a} - S - \mathfrak{b} -bimodule. To these data, we associate the *arrow category* $\mathfrak{b} \rightarrow_M \mathfrak{a}$ which is a $\mathcal{V} \rightarrow_S \mathcal{U}$ graded category. Here, $\mathcal{W} = \mathcal{V} \rightarrow_S \mathcal{U}$ is the underlying arrow category with

$$\mathrm{Ob}(\mathcal{W}) = \mathrm{Ob}(\mathcal{V}) \coprod \mathrm{Ob}(\mathcal{U})$$

and

$$\begin{aligned} \mathcal{W}(V', V) &= \mathcal{V}(V', V), & \mathcal{W}(U, U') &= \mathcal{U}(U, U'), \\ \mathcal{W}(V, U) &= S(V, U), & \mathcal{W}(U, V) &= \emptyset, \end{aligned}$$

for $U, U' \in \mathcal{U}$ and $V, V' \in \mathcal{V}$. Similarly, $\mathfrak{c} = \mathfrak{b} \rightarrow_M \mathfrak{a}$ is the \mathcal{W} -graded category with

$$\mathrm{Ob}(\mathfrak{c}) = \mathrm{Ob}(\mathfrak{a}) \coprod \mathrm{Ob}(\mathfrak{b})$$

and

$$\begin{aligned} \mathfrak{c}_v(B', B) &= \mathfrak{b}_v(B', B), & \mathfrak{c}_u(A, A') &= \mathfrak{a}_u(A, A'), \\ \mathfrak{c}_s(B, A) &= M_s(B, A), \end{aligned}$$

for $v: V' \rightarrow V$ in \mathcal{V} , $B' \in \mathfrak{b}_{V'}$, $B \in \mathfrak{b}_V$, $u: U \rightarrow U'$ in \mathcal{U} , $A \in \mathfrak{a}_U$, $A' \in \mathfrak{a}_{U'}$.

Remark 5.8. Even when \mathfrak{a} and \mathfrak{b} are linear categories with standard grading, and M is an ordinary \mathfrak{a} - \mathfrak{b} -bimodule, the resulting arrow category $\mathfrak{b} \rightarrow_M \mathfrak{a}$ is naturally graded in a nonstandard way since there are no morphisms going from \mathfrak{a} to \mathfrak{b} . \square

Consider the natural inclusions $\varphi_{\mathcal{U}}: \mathcal{U} \rightarrow \mathcal{W}$, $\varphi_{\mathcal{V}}: \mathcal{V} \rightarrow \mathcal{W}$, and $\varphi_{\mathcal{U} \coprod \mathcal{V}}: \mathcal{U} \coprod \mathcal{V} \rightarrow \mathcal{W}$. Clearly, we have $\mathfrak{c}^{\varphi_{\mathcal{U}}} = \mathfrak{a}$, $\mathfrak{c}^{\varphi_{\mathcal{V}}} = \mathfrak{b}$, and $\mathfrak{c}^{\varphi_{\mathcal{U} \coprod \mathcal{V}}} = \mathfrak{a} \coprod \mathfrak{b}$ and we obtain the induced surjections

$$\varphi_{\mathcal{U}}^*: \mathbf{C}_{\mathcal{W}}(\mathfrak{c}) \rightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}),$$

$$\varphi_{\mathcal{V}}^*: \mathbf{C}_{\mathcal{W}}(\mathfrak{c}) \rightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}),$$

and

$$\varphi_{\mathcal{U} \amalg \mathcal{V}}^* \cong \begin{pmatrix} \varphi_{\mathcal{V}}^* \\ \varphi_{\mathcal{U}}^* \end{pmatrix} : \mathbf{C}_{\mathcal{W}}(\mathfrak{c}) \longrightarrow \mathbf{C}_{\mathcal{U} \amalg \mathcal{V}}(\mathfrak{a} \amalg \mathfrak{b}) \cong \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \oplus \mathbf{C}_{\mathcal{U}}(\mathfrak{a}).$$

Further, S defines a thin ideal \mathcal{S} in \mathcal{W} with

$$\mathcal{S}(V, U) = \begin{cases} \mathcal{S}(V, U) & \text{if } U \in \mathcal{U}, V \in \mathcal{V}, \\ \emptyset & \text{else,} \end{cases}$$

and $(\mathcal{S}, \mathcal{U} \amalg \mathcal{V})$ is an ideal-subcategory decomposition in \mathcal{W} . For the category of $\mathfrak{b} \amalg \mathfrak{a}$ -bimodules on \mathcal{S} in the sense of Section 5.6, we clearly have an isomorphism of categories

$$\text{Bimod}_{\mathcal{S}}(\mathfrak{b} \amalg \mathfrak{a}) \cong \text{Bimod}_{\mathcal{S}}(\mathfrak{b}, \mathfrak{a}),$$

and hence, by Corollary 5.7, an isomorphism of categories

$$\text{Bimod}_{\mathcal{S}}(\mathfrak{b}, \mathfrak{a}) \cong \text{Bimod}_{\mathcal{S}}(\mathfrak{c}). \tag{13}$$

5.8 Arrow categories and thin ideals

In this section, we characterize the situation that occurs from the arrow category construction. Let \mathfrak{c} be a \mathcal{W} -graded category and let \mathcal{S} be a thin ideal in \mathcal{W} . We define the full subcategories $\varphi_{\mathcal{U}} : \mathcal{U} \subseteq \mathcal{W}$ and $\varphi_{\mathcal{V}} : \mathcal{V} \subseteq \mathcal{W}$ in the following way. An object $W \in \mathcal{W}$ belongs to \mathcal{V} if there exists a path

$$W \longrightarrow W_1 \longrightarrow \dots \longrightarrow W_n \longrightarrow W_{n+1},$$

for which the last map $W_n \longrightarrow W_{n+1}$ belongs to \mathcal{S} . Similarly, W belongs to \mathcal{U} if there exists a path

$$W_1 \longrightarrow W_2 \longrightarrow \dots \longrightarrow W,$$

for which $W_1 \longrightarrow W_2$ belongs to \mathcal{S} . We put $\mathfrak{a} = \mathfrak{c}^{\varphi_{\mathcal{U}}}$ and $\mathfrak{b} = \mathfrak{c}^{\varphi_{\mathcal{V}}}$. Let S denote the restriction of \mathcal{S} to a \mathcal{U} - \mathcal{V} -bifunctor, and let M be the restriction of $1_{\mathfrak{c}}$ to an \mathfrak{a} - \mathfrak{b} -bimodule.

Proposition 5.9. For $U \in \mathcal{U}$ and $V \in \mathcal{V}$, we have $\mathcal{W}(U, V) = \emptyset$. In particular, the categories \mathcal{U} and \mathcal{V} are disjoint, and there is an injection $\varphi: (\mathcal{V} \rightarrow_S \mathcal{U}) \rightarrow \mathcal{W}$ for which

$$(\mathbf{b} \rightarrow_M \mathbf{a}) = \mathbf{c}^\varphi.$$

The following are equivalent:

- (1) φ is an isomorphism;
- (2) we have $\text{Ob}(\mathcal{W}) = \text{Ob}(\mathcal{V}) \cup \text{Ob}(\mathcal{U})$ and $\mathcal{S}(V, U) = \mathcal{W}(V, U)$ for $V \in \mathcal{V}$ and $U \in \mathcal{U}$.

In this case, we have $(\mathbf{b} \rightarrow_M \mathbf{a}) = \mathbf{c}$. □

Proof. Suppose that there is a morphism $U \rightarrow V$ with $U \in \mathcal{U}$ and $V \in \mathcal{V}$. Then there is a path

$$W_1 \rightarrow W_2 \rightarrow \cdots \rightarrow U \rightarrow V \rightarrow \cdots \rightarrow W_n \rightarrow W_{n+1},$$

with the two morphisms at the ends belonging to \mathcal{S} . But then since \mathcal{S} is an ideal, also $W_2 \rightarrow \cdots \rightarrow W_{n+1}$ belongs to \mathcal{S} . With $W_1 \rightarrow W_2$ in \mathcal{S} , this contradicts the thinness of \mathcal{S} .

The functor φ is clearly injective on objects and on morphisms in \mathcal{U} or \mathcal{V} , and since $\mathcal{S}(V, U) \subseteq \mathcal{W}(V, U)$, the functor is indeed injective. The description of \mathbf{c}^φ clearly follows.

It remains to show the equivalence of (1) and (2). Condition (2) is clearly necessary for φ to be surjective. Conversely, surjectivity readily follows from (2) taking into account that for $U \in \mathcal{U}$ and $V \in \mathcal{V}$, we have $\mathcal{W}(U, V) = \emptyset$. ■

Recall that a delta is a category in which the arrows go only one way, that is, we have the following definition.

Definition 5.10. A *delta* is a category \mathcal{W} such that for all $W \neq W' \in \mathcal{W}$, we have $\mathcal{W}(W, W') = \emptyset$ or $\mathcal{W}(W', W) = \emptyset$. □

Example 5.11. Let \mathcal{W} be a delta with a terminal object $* \in \mathcal{W}$. For $W \in \mathcal{W}$, let $*_W: W \rightarrow *$ denote the unique morphism. Let \mathcal{S} consist of all morphisms $*_W: W \rightarrow *$ for $W \neq *$. Since \mathcal{W} is a delta and $*$ is terminal, for all $W \neq *$ we have $\mathcal{W}(*, W) = \emptyset$ and we have $\mathcal{W}(*, *) = \{1_*\} = \{*_*\}$. Then \mathcal{S} is a thin ideal by Example 5.5. In the above notation, \mathcal{U} consists of $*$ and \mathcal{V} consists of all other objects. Then condition (2) in Proposition 5.9 is fulfilled whence $\mathcal{W} \cong (\mathcal{V} \rightarrow_S \mathcal{U})$. □

5.9 Connecting homomorphism

In the notation of Section 5.7, we now investigate the complex

$$\mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}}),$$

which fits into the following exact sequence from (12):

$$0 \longrightarrow \mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}}) \longrightarrow \mathbf{C}_{\mathcal{W}}(\mathfrak{c}) \xrightarrow{\begin{pmatrix} \varphi_{\mathcal{V}}^* \\ \varphi_{\mathcal{U}}^* \end{pmatrix}} \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \oplus \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow 0$$

Since we are interested in identifying when $\varphi_{\mathcal{V}}^*$ and $\varphi_{\mathcal{U}}^*$ are quasi-isomorphisms, we will describe the components of the connecting homomorphism explicitly. In fact, this morphism is determined by two maps on the chain level which we will describe next:

$$\alpha : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})[1]$$

and

$$\beta : \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \longrightarrow \mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})[1].$$

To do this, we first make $\mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})$ explicit. By definition, $\mathbf{C}_{\mathcal{W}}^n(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})$ is given by

$$\prod_{(u, A) \in \mathcal{N}(\mathfrak{c}) \exists u_i \in \mathcal{S}} \text{Hom}_k(\mathfrak{b}_{u_{n-1}}(A_{n-1}, A_n) \otimes \cdots \otimes M_{u_i}(A_i, A_{i+1}) \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), M_{|u|}(A_0, A_n)).$$

Let $\phi \in \mathbf{C}_{\mathcal{U}}^n(\mathfrak{a})$ be a Hochschild cocycle. Following [12], we define $\alpha(\phi)$ with nonzero components in

$$\text{Hom}_k(M_{u_n}(A_n, A_{n+1}) \cdots \otimes \mathfrak{a}_{u_0}(A_0, A_1), M_{|u|}(A_0, A_{n+1})),$$

given by

$$\alpha(\phi)(x, a_{n-1}, \dots, a_0) = x\phi(a_{n-1}, \dots, a_0),$$

and similarly for β .

Lemma 5.12. The maps α and β are chain maps for which

$$\begin{pmatrix} \beta & \alpha \end{pmatrix} : \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \oplus \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})[1],$$

induces the connecting homomorphisms. □

The triangle

$$\mathbf{C}_{\mathcal{W}}(\mathfrak{c}) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \oplus \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})[1] \longrightarrow \quad (14)$$

$$\begin{pmatrix} \varphi_{\mathcal{V}}^* \\ \varphi_{\mathcal{U}}^* \end{pmatrix} \qquad \qquad \qquad \begin{pmatrix} \beta & \alpha \end{pmatrix}$$

corresponds to a homotopy bi-Cartesian square

$$\begin{array}{ccc} \mathbf{C}_{\mathcal{W}}(\mathfrak{c}) & \xrightarrow{\varphi_{\mathcal{U}}^*} & \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \\ \varphi_{\mathcal{V}}^* \downarrow & & \downarrow \alpha \\ \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) & \xrightarrow{\beta} & \mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})[1] \end{array}$$

We thus have the following proposition.

Proposition 5.13. (1) If α is a quasi-isomorphism, then so is $\varphi_{\mathcal{V}}^*$.

(2) If β is a quasi-isomorphism, then so is $\varphi_{\mathcal{U}}^*$. □

5.10 Derived interpretation

Following [12], we now give a derived interpretation of the morphisms α and β . We start by giving various derived interpretations of $\mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})$.

Let $B(\mathfrak{a}) \rightarrow 1_{\mathfrak{a}}$ and $B(\mathfrak{b}) \rightarrow 1_{\mathfrak{b}}$ be the Bar resolutions in $\mathbf{Bimod}_{\mathcal{U}}(\mathfrak{a})$ and $\mathbf{Bimod}_{\mathcal{V}}(\mathfrak{b})$, respectively, as defined in [14, Section 3.2]. We have

$$\mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \cong \mathbf{Hom}_{\mathbf{Bimod}_{\mathcal{U}}(\mathfrak{a})}(B(\mathfrak{a}), 1_{\mathfrak{a}}) = \mathbf{RHom}_{\mathbf{Bimod}_{\mathcal{U}}(\mathfrak{a})}(1_{\mathfrak{a}}, 1_{\mathfrak{a}})$$

and

$$\mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \cong \mathbf{Hom}_{\mathbf{Bimod}_{\mathcal{V}}(\mathfrak{b})}(B(\mathfrak{b}), 1_{\mathfrak{b}}) = \mathbf{RHom}_{\mathbf{Bimod}_{\mathcal{V}}(\mathfrak{b})}(1_{\mathfrak{b}}, 1_{\mathfrak{b}}).$$

Similarly, we have

$$\mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_{\mathcal{S}})[1] \cong \mathbf{Hom}_{\mathbf{Bimod}_{\mathcal{S}}(\mathfrak{a}, \mathfrak{b})}(B(\mathfrak{a}) \otimes_{\mathfrak{a}} M \otimes_{\mathfrak{b}} B(\mathfrak{b}), M)$$

and

$$\mathbf{Hom}_{\mathbf{Bimod}_{\mathcal{S}}(\mathfrak{a}, \mathfrak{b})}(B(\mathfrak{a}) \otimes_{\mathfrak{a}} M \otimes_{\mathfrak{b}} B(\mathfrak{b}), M) = \mathbf{RHom}_{\mathbf{Bimod}_{\mathcal{S}}(\mathfrak{a}, \mathfrak{b})}(M, M).$$

In particular, from the triangle (14) we obtain a long exact cohomology sequence

$$\cdots \longrightarrow HH_{\mathcal{W}}^i(\mathfrak{c}) \longrightarrow HH_{\mathcal{V}}^i(\mathfrak{b}) \oplus HH_{\mathcal{U}}^i(\mathfrak{a}) \longrightarrow \text{Ext}_{\text{Bimod}_S(\mathfrak{b}, \mathfrak{a})}^i(M, M) \longrightarrow \cdots$$

By Proposition 3.6, we further have

$$\mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_S)[1] \cong \text{Hom}_{\text{Bimod}_{\mathcal{U}}(\mathfrak{a})}(B(\mathfrak{a}), \text{Hom}_{\mathfrak{b}}(M \otimes_{\mathfrak{b}} B(\mathfrak{b}), M)),$$

with $\text{Hom}_{\mathfrak{b}}(M \otimes_{\mathfrak{b}} B(\mathfrak{b}), M) = \text{RHom}_{\mathfrak{b}}(M, M)$. Similarly,

$$\mathbf{C}_{\mathcal{W}}(\mathfrak{c}, (1_{\mathfrak{c}})_S)[1] \cong \text{Hom}_{\text{Bimod}_{\mathcal{V}}(\mathfrak{b})}(B(\mathfrak{b}), \text{Hom}_{\mathfrak{a}^{\text{op}}}(B(\mathfrak{a}) \otimes_{\mathfrak{a}} M, M)),$$

with $\text{Hom}_{\mathfrak{a}^{\text{op}}}(B(\mathfrak{a}) \otimes_{\mathfrak{a}} M, M) = \text{RHom}_{\mathfrak{a}^{\text{op}}}(M, M)$.

Consider the natural morphisms

$$\lambda : 1_{\mathfrak{a}} \longrightarrow \text{Hom}_{\mathfrak{b}}(M, M) \longrightarrow \text{RHom}_{\mathfrak{b}}(M, M)$$

and

$$\omega : 1_{\mathfrak{b}} \longrightarrow \text{Hom}_{\mathfrak{a}^{\text{op}}}(M, M) \longrightarrow \text{RHom}_{\mathfrak{a}^{\text{op}}}(M, M)$$

of bimodules and the induced

$$\mathbf{C}_{\mathcal{U}}(\mathfrak{a}, \lambda) : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, \text{RHom}_{\mathfrak{b}}(M, M))$$

and

$$\mathbf{C}_{\mathcal{V}}(\mathfrak{b}, \omega) : \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, \text{RHom}_{\mathfrak{a}}(M, M)).$$

We have the following map-graded version theorem [15, Theorem 4.1.1] extracted from [12, Section 4].

Theorem 5.14.

- (1) We have $\alpha \cong \mathbf{C}_{\mathcal{U}}(\mathfrak{a}, \lambda)$ and $\beta \cong \mathbf{C}_{\mathcal{V}}(\mathfrak{b}, \omega)$.
- (2) If $\mathbf{C}_{\mathcal{U}}(\mathfrak{a}, \lambda)$ is a quasi-isomorphism (hence, in particular, if λ is a quasi-isomorphism), then so is $\varphi_{\mathcal{V}}^*$.
- (3) If $\mathbf{C}_{\mathcal{V}}(\mathfrak{b}, \omega)$ is a quasi-isomorphism (hence, in particular, if ω is a quasi-isomorphism), then so is $\varphi_{\mathcal{U}}^*$. □

The analog of [12, Section 4.6, Theorem] in the map-graded context can be formulated and proved in a similar fashion. In particular, we mention the following compatibility results explicitly.

Let $(\varphi, F) : (\mathcal{U}, \mathfrak{a}) \longrightarrow (\mathcal{V}, \mathfrak{b})$ be a sub-Cartesian graded functor with associated \mathcal{U} - \mathcal{V} -bifunctor S^φ and \mathfrak{a} - S^φ - \mathfrak{b} -bimodule M^F as in Example 3.2. On the one hand, since (φ, F) is sub-Cartesian, we have the induced morphism of B_∞ -algebras

$$F^* : \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \longrightarrow \mathbf{C}_{\mathcal{U}}(\mathfrak{a}),$$

from Proposition 4.2.

Proposition 5.15. (1) $\lambda : 1_{\mathfrak{a}} \longrightarrow \mathrm{RHom}_{\mathfrak{b}}(M^F, M^F)$ and hence $\varphi_{\mathcal{V}}^*$ is a quasi-isomorphism.

(2) In the homotopy category of B_∞ -algebras, we have $F^* = \varphi_{\mathcal{U}}^* \varphi_{\mathcal{V}}^{*-1}$. \square

Let $(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a})$ be a sub-Cartesian graded functor with associated \mathcal{U} - \mathcal{V} -bifunctor S_φ and \mathfrak{a} - S_φ - \mathfrak{b} -bimodule M_F as in Example 3.2. This time we have the functor

$$F^* : \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \longrightarrow \mathbf{C}_{\mathcal{V}}(\mathfrak{b}).$$

Proposition 5.16. (1) $\omega : 1_{\mathfrak{b}} \longrightarrow \mathrm{RHom}_{\mathfrak{a}^{\mathrm{op}}}(M_F, M_F)$ and hence $\varphi_{\mathcal{U}}^*$ is a quasi-isomorphism.

(2) In the homotopy category of B_∞ -algebras, we have $F^* = \varphi_{\mathcal{V}}^* \varphi_{\mathcal{U}}^{*-1}$. \square

6 Grothendieck Construction

In this section, we present a unified framework for constructing map-graded categories and deconstructing their Hochschild complexes. The classical Grothendieck construction from [1] takes a pseudofunctor $\mathcal{U} \longrightarrow \mathbf{Cat}$ as input and turns it into a category fibered over \mathcal{U} . The main construction from [14] is a k -linearized version of this construction. Now, we go a step further and start from a pseudofunctor

$$(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \longrightarrow \underline{\mathbf{Map}} : \mathcal{C} \longmapsto (\mathcal{U}_{\mathcal{C}}, \mathfrak{a}_{\mathcal{C}}),$$

where \mathcal{C} is a small category and $\underline{\mathbf{Map}}$ is the bicategory of map-graded categories and bimodules described in Section 3.2. Allowing arbitrary bimodules rather than functors between map-graded categories allows us to capture the arrow category with respect to

a bimodule from Section 5.7. In general, we can now deconstruct the Hochschild complex of the Grothendieck construction $(\tilde{\mathcal{U}}, \tilde{\alpha})$ of (\mathcal{U}, α) based upon the internal structure of \mathcal{C} . Here, the strategy is to cover $(\tilde{\mathcal{U}}, \tilde{\alpha})$ by other Grothendieck constructions, using base change for pseudofunctors from Section 6.5. For instance, in Section 6.6, we consider “generalized arrow categories”, and deconstruct them using iterated arrow category constructions. In Section 6.7, we observe how, in the case where \mathcal{C} is a poset, the sheaf property for Hochschild complexes on the one hand, and the arrow category construction on the other hand, can be seen as complementary tools for deconstructing Hochschild cohomology. In Section 6.8, we start from a pseudofunctor $(\mathcal{U}^*, \alpha^*) : \mathcal{C}^* \rightarrow \text{Map}_{\text{sc}}$, which we compare with the natural pseudofunctor of Grothendieck constructions

$$(\mathcal{U}^*, \alpha^*) : \mathcal{C}^* \rightarrow \text{Map}_{\text{sc}} : \mathcal{C} \rightarrow (\tilde{\mathcal{U}}|_{\mathcal{C}}, \tilde{\alpha}|_{\mathcal{C}}),$$

built from the restriction (\mathcal{U}, α) of $(\mathcal{U}^*, \alpha^*)$ to \mathcal{C} . Our main Theorem 6.14 is heavily based upon Keller’s arrow category argument in the case of a fully faithful functor $\mathfrak{b} \rightarrow \mathfrak{a}$, which is in fact a special case of our theorem. As an application, we recover the Mayer–Vietoris triangles for ringed spaces from [15, Section 7.9].

6.1 Diagrams in the bicategory of categories

Let $\underline{\text{Cat}}$ be the bicategory of categories, bifunctors, and natural transformations described in Section 3.1. Let \mathcal{C} be an arbitrary small category and consider a pseudofunctor

$$\mathcal{U} : \mathcal{C} \rightarrow \underline{\text{Cat}} : \mathcal{C} \mapsto \mathcal{U}_{\mathcal{C}}.$$

The pseudofunctor maps a map $c : \mathcal{C} \rightarrow \mathcal{C}'$ to a $\mathcal{U}_{\mathcal{C}'}$ - $\mathcal{U}_{\mathcal{C}}$ -bifunctor $\mathcal{U}(c) = S_c$ and for an additional $c' : \mathcal{C}' \rightarrow \mathcal{C}''$, there is an isomorphism

$$\phi_{c',c} : S_{c'} \circ S_c \rightarrow S_{c'c}.$$

For a third map $c'' : \mathcal{C}'' \rightarrow \mathcal{C}'''$, we have a commutative diagram

$$\begin{array}{ccccc}
 (S_{\mathcal{C}'} \circ S_{\mathcal{C}}) \circ S_{\mathcal{C}} & \xrightarrow{\phi_{c',c} \circ 1} & S_{\mathcal{C}'} \circ S_{\mathcal{C}} & \xrightarrow{\phi_{c'',c'}} & S_{c''c} \\
 \alpha \downarrow & & & & \downarrow 1 \\
 S_{\mathcal{C}'} \circ (S_{\mathcal{C}} \circ S_{\mathcal{C}}) & \xrightarrow{1 \circ \phi_{c',c}} & S_{\mathcal{C}'} \circ S_{\mathcal{C}c} & \xrightarrow{\phi_{c'',c'}} & S_{c''c}
 \end{array} \tag{15}$$

where α is the isomorphism from the bicategory $\underline{\text{Cat}}$. We suppose moreover that $S_{1_C} = 1_{\mathcal{U}_C}$.

We define the *Grothendieck construction* of \mathcal{U} to be the nonlinear \mathcal{C} -graded category \mathcal{U} with \mathcal{U}_C as prescribed and, for $c: C \rightarrow C', U \in \mathcal{U}_C, U' \in \mathcal{U}_{C'}$:

$$\mathcal{U}_c(U, U') = S_c(U, U').$$

Recall that

$$S_{c'} \circ S_c(U, U') = \coprod_{V \in \mathcal{U}_{c'}} S_{c'}(V, U'') \times S_c(U, V) / \sim.$$

The composition on \mathcal{U} is defined as the natural map

$$S_{c'}(U', U'') \times S_c(U, U') \longrightarrow S_{c'} \circ S_c(U, U'') \xrightarrow{\phi_{c',c}} S_{c'c}(U, U''),$$

and we denote the image of (s', s) under composition by $s's$. There is a corresponding category $\tilde{\mathcal{U}}$ over \mathcal{C} with $\text{Ob}(\tilde{\mathcal{U}}) = \coprod_{C \in \mathcal{C}} \mathcal{U}_C$ (see Remark 2.9).

6.2 Diagrams in the bicategory of map-graded categories

The construction from Section 6.1 can be extended in the following way. Let $\underline{\text{Map}}$ be the bicategory of map-graded categories, bimodules, and their morphisms described in Section 3.2. Let \mathcal{C} be an arbitrary small category and consider a pseudofunctor

$$(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \longrightarrow \underline{\text{Map}} : C \longmapsto (\mathcal{U}_C, \mathfrak{a}_C).$$

The pseudofunctor maps $c: C \rightarrow C'$ to an $\mathfrak{a}_{C'}\text{-}\mathfrak{a}_C$ -bimodule $(\mathcal{U}(c) = S_c, \mathfrak{a}(c) = M_c)$ and for an additional $c': C' \rightarrow C''$ there is an isomorphism

$$(\phi_{c',c}, F_{c',c}) : (S_{c'} \circ S_c, M_{c'} \otimes_{\mathfrak{a}_{c'}} M_c) \longrightarrow (S_{c'c}, M_{c'c}).$$

These isomorphisms satisfy the natural coherence axiom similar to (15) and $(S_{1_C}, M_{1_C}) = (1_{\mathcal{U}_C}, 1_{\mathfrak{a}_C})$.

Clearly, there is an underlying pseudofunctor

$$\mathcal{U} : \mathcal{C} \longrightarrow \underline{\text{Cat}} : C \longmapsto \mathcal{U}_C,$$

which determines a nonlinear \mathcal{C} -graded category \mathcal{U} and associated category $\tilde{\mathcal{U}}$. We define the *Grothendieck construction* of $(\mathcal{U}, \mathfrak{a})$ to be the following $\tilde{\mathcal{U}}$ -graded category $\tilde{\mathfrak{a}}$. For

$U \in \mathcal{U}_C$, we define

$$\tilde{\mathfrak{a}}_U = (\mathfrak{a}_C)_U.$$

For a morphism $(c: C \rightarrow C', s \in \mathcal{S}_C(U, U'))$ in $\tilde{\mathcal{U}}$, with $U \in \mathcal{U}_C$, $U' \in \mathcal{U}_{C'}$, and for $A \in (\mathfrak{a}_C)_U$, $A' \in (\mathfrak{a}_{C'})_{U'}$, we put

$$\tilde{\mathfrak{a}}_{(c,s)}(A, A') = (M_C)_s(A, A').$$

Recall that

$$(M_{C'} \otimes_{\mathfrak{a}_{C'}} M_C)_r(A, A'') = \bigoplus_{r=[(s',s)], B} (M_{C'})_{s'}(B, A'') \otimes (M_C)_s(A, B) / \sim.$$

The composition on $\tilde{\mathfrak{a}}$ is defined as the composition of the natural map

$$(M_{C'})_{s'}(A', A'') \otimes (M_C)_s(A, A') \longrightarrow (M_{C'} \otimes_{\mathfrak{a}_{C'}} M_C)_{[(s',s)]}(A, A''),$$

followed by the map

$$(M_{C'} \otimes_{\mathfrak{a}_{C'}} M_C)_{[(s',s)]}(A, A'') \xrightarrow{F_{c',c}} (M_{C'})_{s's}(A, A'').$$

6.3 Diagrams in the bicategory of linear categories

In many applications, we are in a somewhat simplified situation from Section 6.2. Let $\underline{\mathcal{C}\text{at}}(k)$ be the bicategory of k -linear functors and bimodules. There is a natural map $\underline{\mathcal{C}\text{at}}(k) \rightarrow \underline{\text{Map}}: \mathfrak{a} \mapsto \mathfrak{a}_{\text{tr}}$ where \mathfrak{a}_{tr} is the trivially graded category over e from Example 2.11(2). Consider a pseudofunctor

$$\mathfrak{a}: \mathcal{C} \rightarrow \underline{\mathcal{C}\text{at}}(k): C \mapsto \mathfrak{a}_C.$$

If we consider the corresponding $(\mathcal{E}, \mathfrak{a}): \mathcal{C} \rightarrow \underline{\text{Map}}: C \mapsto (e, \mathfrak{a}_C)$, we see that there is a canonical isomorphism $\tilde{\mathcal{E}} \cong \mathcal{C}$ and carrying out the construction from Section 6.2, we thus obtain a \mathcal{C} -graded category $\tilde{\mathfrak{a}}$ with

$$\tilde{\mathfrak{a}}_C = \mathfrak{a}_C.$$

For a morphism $c: C \rightarrow C'$ in \mathcal{C} and for $A \in \mathfrak{a}_C$, $A' \in \mathfrak{a}_{C'}$, we have

$$\tilde{\mathfrak{a}}_c(A, A') = M_C(A, A'),$$

and composition is defined in the obvious way.

Conversely, let \mathfrak{a} be an arbitrary \mathcal{U} -graded category. We define the naturally associated pseudofunctor

$$\mathcal{A} : \mathcal{U} \longrightarrow \underline{\mathbf{Cat}}(k) : U \longrightarrow \mathcal{A}_U,$$

where \mathcal{A}_U is the category with $\text{Ob}(\mathcal{A}_U) = \mathfrak{a}_U$ and $\mathcal{A}(A, A') = \mathfrak{a}_{1_U}(A, A')$ and where for $u : U \longrightarrow U'$ in \mathcal{U} , $\mathcal{A}(u) = M_u$ is the natural $\mathcal{A}_{U'}\text{-}\mathcal{A}_U$ -bimodule with

$$M_u(A, A') = \mathfrak{a}_u(A, A').$$

Then $(\mathcal{U}, \mathfrak{a}) \cong (\mathcal{U}, \tilde{\mathfrak{A}})$ and hence the correspondence between *fibred* \mathcal{U} -graded categories and pseudofunctors $\mathcal{U} \longrightarrow \mathbf{Cat}(k)$ naturally extends to a correspondence between *all* \mathcal{U} -graded categories and pseudofunctors $\mathcal{U} \longrightarrow \underline{\mathbf{Cat}}(k)$.

6.4 Diagrams in the category of map-graded categories

Another specification of the setup from Section 6.2 occurs if we consider a pseudofunctor

$$(\mathcal{U}, \mathfrak{a}) : \mathcal{C}^{\text{op}} \longrightarrow \mathbf{Map} : C \longmapsto (\mathcal{U}_C, \mathfrak{a}_C),$$

mapping a morphism $c : C \longrightarrow C'$ in \mathcal{C} to a map-graded functor

$$(\varphi_c, F_c) : (\mathcal{U}_{C'}, \mathfrak{a}_{C'}) \longrightarrow (\mathcal{U}_C, \mathfrak{a}_C).$$

By Example 3.2(2), this gives rise to a $\mathcal{U}_{C'}\text{-}\mathcal{U}_C$ -bimodule S^{φ_c} and a $\mathfrak{a}_{C'}\text{-}S^{\varphi_c}\text{-}\mathfrak{a}_C$ -bimodule M^{F_c} with

$$S^{\varphi_c}(U, U') = \mathcal{U}_C(U, \varphi_c(U')); \quad M_s^{F_c}(A, A') = \mathfrak{a}_{C_s}(A, F_c(A')).$$

In fact, there are natural pseudofunctors $\mathbf{Cat}^{\text{op}} \longrightarrow \underline{\mathbf{Cat}}$ and $\mathbf{Map}^{\text{op}} \longrightarrow \underline{\mathbf{Map}}$ so that we obtain a composed pseudofunctor

$$(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \longrightarrow \mathbf{Map} \longrightarrow \underline{\mathbf{Map}} : C \longmapsto (\mathcal{U}_C, \mathfrak{a}_C).$$

For $c : C \longrightarrow C'$, $c' : C' \longrightarrow C''$ in \mathcal{C} , we are given natural isomorphisms

$$(\eta, \theta) : (\varphi_c \varphi_{c'}, F_c F_{c'}) = (\varphi_c, F_c)(\varphi_{c'}, F_{c'}) \longrightarrow (\varphi_{c'c}, F_{c'c})$$

satisfying the natural coherence axiom similar to (15). From this, we obtain natural isomorphisms

$$(S^{\varphi_{c'}} \circ S^{\varphi_c}, M^{F_{c'}} \otimes_{\mathfrak{a}_{C'}} M^{F_c}) \longrightarrow (S^{\varphi_{c'c}}, M^{F_{c'c}}),$$

in the following way. On the level of bifunctors, we have

$$\begin{aligned}
 S^{\varphi_{c'}} \circ S^{\varphi_c}(U, U'') &= \coprod_{V \in \mathcal{U}_{c'}} \mathcal{U}_{c'}(V, \varphi_{c'}(U'')) \times \mathcal{U}_C(U, \varphi_c(V)) / \sim \\
 &\xrightarrow{\varphi_c \times 1} \\
 \coprod_{V \in \mathcal{U}_{c'}} \mathcal{U}_C(\varphi_c(V), \varphi_c \varphi_{c'}(U'')) \times \mathcal{U}_C(U, \varphi_c(V)) / \sim \\
 &\xrightarrow{m_{\mathcal{U}_C}} \mathcal{U}_C(U, \varphi_c \varphi_{c'}(U'')) \\
 &\xrightarrow{m_{\mathcal{U}_C}(\eta_{U''}, -)} \mathcal{U}_C(U, \varphi_{c'c}(U'')) = S^{\varphi_{c'c}}(U, U'').
 \end{aligned}$$

Similarly, on the level of bimodules we have

$$\begin{aligned}
 M^{F_{c'}} \otimes_{\mathfrak{a}_{c'}} M_r^{F_c}(A, A'') &= \bigoplus_{r=[(s',s)], B} (\mathfrak{a}_{c'})_{s'}(B, F_{c'}(A'')) \otimes (\mathfrak{a}_C)_s(A, F_C(B)) / \sim \\
 &\xrightarrow{F_c \otimes 1} \\
 \bigoplus_{r=[(s',s)], B} (\mathfrak{a}_C)_{\varphi_c(s')} (F_C(B), F_C F_{c'}(A'')) \otimes (\mathfrak{a}_C)_s(A, F_C(B)) / \sim \\
 &\xrightarrow{m_{\mathfrak{a}_C}} (\mathfrak{a}_C)_{\varphi_c(s')s}(A, F_C F_{c'}(A'')) \\
 &\xrightarrow{m_{\mathfrak{a}_C}(\theta_{A''}, -)} (\mathfrak{a}_C)_{\varphi_c(s')s}(A, F_{c'c}(A'')) = M_{s's}^{F_{c'c}}(A, A'').
 \end{aligned}$$

These maps determine the composition on the Grothendieck construction $(\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$.

Remark 6.1. Note that the nonlinear \mathcal{C} -graded category $\tilde{\mathcal{U}}$ is the original Grothendieck construction of $\mathcal{U} : \mathcal{C} \rightarrow \text{Cat}$ in the sense of Grothendieck. \square

Remark 6.2. For a pseudofunctor $\mathfrak{a} : \mathcal{C} \rightarrow \text{Cat}(k)$ landing in the 2-category of k -linear categories, functors, and natural transformations, composing with $\text{Cat}(k) \rightarrow \underline{\text{Cat}}(k)$

brings us in the situation of Section 6.3. The Grothendieck construction $\tilde{\mathfrak{a}}$ is the one we used in [14] in order to turn a presheaf of k -algebras, or more generally a pseudofunctor $\mathfrak{a} : \mathcal{C} \rightarrow \text{Cat}(k)$, into a \mathcal{C} -graded category and define its structured Hochschild complex $\mathbf{C}_{\mathcal{C}}(\tilde{\mathfrak{a}})$. By [17], this complex computes the natural Hochschild cohomology of a generalized from Gerstenhaber and Schack's Hochschild cohomology of presheaves of algebras [5, 6]. \square

Remark 6.3. Let $(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \rightarrow \text{Map}$ be a pseudofunctor. In a completely similar fashion, the $\mathcal{U}_{\mathcal{C}'}\text{-}\mathcal{U}_{\mathcal{C}}$ -bifunctors S_{φ_c} and $\mathfrak{a}_{\mathcal{C}'}\text{-}S_{\varphi_c}\text{-}\mathfrak{a}_{\mathcal{C}}$ -bimodules M_{F_c} from Example 3.2(1) give rise to a pseudofunctor

$$(\mathcal{U}', \mathfrak{a}') : \mathcal{C} \rightarrow \text{Map} \rightarrow \underline{\text{Map}}. \quad \square$$

6.5 Base change

Let $\mathcal{U} : \mathcal{C} \rightarrow \underline{\text{Cat}}$ be a pseudofunctor as in Section 6.1 and let $\Phi : \mathcal{D} \rightarrow \mathcal{C}$ be an arbitrary functor. There is a resulting composed pseudofunctor

$$\mathcal{U}^{\Phi} : \mathcal{D} \rightarrow \mathcal{C} \rightarrow \underline{\text{Cat}} : D \mapsto \mathcal{U}_{\Phi(D)},$$

with an associated category $\tilde{\mathcal{U}}^{\Phi}$. The natural maps

$$\text{Ob}(\tilde{\mathcal{U}}^{\Phi}) = \coprod_{D \in \mathcal{D}} \mathcal{U}_{\Phi(D)} \rightarrow \coprod_{C \in \mathcal{C}} \mathcal{U}_C = \text{Ob}(\tilde{\mathcal{U}})$$

and

$$\tilde{\mathcal{U}}^{\Phi}((D, U), (D', U')) \rightarrow \tilde{\mathcal{U}}((\Phi(D), U), (\Phi(D'), U')),$$

given by

$$\coprod_{d \in \mathcal{D}(D, D')} S_{\Phi(d)}(U, U') \rightarrow \coprod_{c \in \mathcal{C}(\Phi(D), \Phi(D'))} S_c(U, U'),$$

give rise to a functor

$$\varphi : \tilde{\mathcal{U}}^{\Phi} \rightarrow \tilde{\mathcal{U}}. \quad (16)$$

Proposition 6.4. Let \mathcal{U} and Φ be as above. If $\mathcal{N}_1(\Phi)$ is injective, then so is $\mathcal{N}_1(\varphi)$. \square

Proposition 6.5. Let \mathcal{U} be as above and consider a collection $(\Phi_i : \mathcal{D}_i \rightarrow \mathcal{C})_{i \in I}$ of functors. If, for $n \in \mathbb{N} \cup \{\infty\}$, the collection constitutes an n -cover in Cat , then so does the collection $(\tilde{\Phi}_i : \tilde{\mathcal{U}}^{\Phi_i} \rightarrow \tilde{\mathcal{U}})_{i \in I}$. \square

Now let $(\mathcal{U}, \alpha) : \mathcal{C} \rightarrow \underline{\text{Map}}$ be a pseudofunctor as in Section 6.2 and $\Phi : \mathcal{D} \rightarrow \mathcal{C}$ be a functor. We now obtain a composed pseudofunctor

$$(\mathcal{U}^\Phi, \alpha^\Phi) : \mathcal{D} \rightarrow \mathcal{C} \rightarrow \underline{\text{Map}},$$

with an associated $\tilde{\mathcal{U}}^\Phi$ -graded category $\tilde{\alpha}^\Phi$.

Proposition 6.6. With the notation of Proposition 2.25, we have

$$\tilde{\alpha}^\Phi = (\tilde{\alpha})^\varphi.$$

In particular, there is a natural Cartesian map-graded functor

$$\tilde{\Phi} = (\varphi, \delta^{\varphi, \tilde{\alpha}}) : (\tilde{\mathcal{U}}^\Phi, \tilde{\alpha}^\Phi) \rightarrow (\tilde{\mathcal{U}}, \tilde{\alpha}). \quad \square$$

Next we use base changes to transform a pseudofunctor $(\mathcal{U}, \alpha) : \mathcal{C} \rightarrow \underline{\text{Map}}$ into a functor

$$(\mathcal{U}^*, \alpha^*) : \mathcal{C}^* \rightarrow \text{Map}$$

of Grothendieck constructions. We first define the category \mathcal{C}^* to be the arrow category $\mathcal{C} \rightarrow_{S} e$ for the unique e - \mathcal{C} -bifunctor S with $S(C, *) = \{*\}$. For every $C \in \mathcal{C}$, we obtain a composition

$$(\mathcal{U}|_C, \alpha|_C) : \mathcal{C}/\mathcal{C} \rightarrow \mathcal{C} \rightarrow \underline{\text{Map}},$$

and a Cartesian map-graded functor $(\tilde{\mathcal{U}}|_C, \tilde{\alpha}|_C) \rightarrow (\tilde{\mathcal{U}}, \tilde{\alpha})$. For every $c : C' \rightarrow C$ in \mathcal{C} , we have a natural functor $\mathcal{C}/C' \rightarrow \mathcal{C}/C$ and an induced Cartesian map-graded functor $(\tilde{\mathcal{U}}|_{C'}, \tilde{\alpha}|_{C'}) \rightarrow (\tilde{\mathcal{U}}|_C, \tilde{\alpha}|_C)$ and it is not hard to organize these data into a functor

$$(\mathcal{U}^*, \alpha^*) : \mathcal{C}^* \rightarrow \text{Map}_{\text{sc}} : \mathcal{C} \mapsto (\tilde{\mathcal{U}}|_C, \tilde{\alpha}|_C),$$

where we define $(\tilde{\mathcal{U}}|_*, \tilde{\alpha}|_*) = (\tilde{\mathcal{U}}, \tilde{\alpha})$.

Proposition 6.7. Suppose that \mathcal{C} has finite products. Let $(C_i)_{i \in I}$ be a collection of objects in \mathcal{C} such that for every $C \in \mathcal{C}$ there exists a map $C \rightarrow C_i$ for some i . The composed functor

$$\mathbf{C} : \mathcal{C}^* \rightarrow \text{Map}_{\text{sc}} \rightarrow B_\infty : \mathcal{C} \mapsto \mathbf{C}_{\tilde{\mathcal{U}}|_C}(\tilde{\alpha}|_C)$$

satisfies the sheaf property with respect to the collection of maps $(C_i \rightarrow *)_{i \in I}$ in \mathcal{C}^* . \square

Proof. By Example 2.34(2), the collection $(\mathcal{C}/C_i \rightarrow \mathcal{C})_{i \in I}$ constitutes an ∞ -cover of \mathcal{C} in Cat . By proposition 6.5, the induced restriction maps in Map_{sc} also form an ∞ -cover. Further, it is readily seen that the pullback of $C_i \rightarrow *$ and $C_j \rightarrow *$ in \mathcal{C}^* is given by $C_i \times C_j \rightarrow *$ for the product $C_i \times C_j$ in \mathcal{C} . The pullback of $(\tilde{\mathcal{U}}|_{C_i}, \tilde{\mathfrak{a}}|_{C_i}) \rightarrow (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ and $(\tilde{\mathcal{U}}|_{C_j}, \tilde{\mathfrak{a}}|_{C_j}) \rightarrow (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ is given by $(\tilde{\mathcal{U}}|_{C_i \times C_j}, \tilde{\mathfrak{a}}|_{C_i \times C_j}) \rightarrow (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$. Thus, the result follows from Theorem 4.7. \blacksquare

6.6 Generalized arrow categories

We can cast the arrow category construction from Section 5.7 in the setup from Section 6.2. Consider the path category

$$\mathcal{C}' = \langle 0 \xrightarrow{\leq} 1 \rangle.$$

With the notation of Section 5.7, we obtain a pseudofunctor

$$(\mathcal{W}', c') : \mathcal{C}' \rightarrow \underline{\text{Map}},$$

with

$$\begin{aligned} \mathcal{W}'_0 &= \mathcal{V} \quad \mathcal{W}'_1 = \mathcal{U}, & \mathcal{W}'(0 \leq 1) &= S, \\ c'_0 &= \mathfrak{b}, \quad c'_1 = \mathfrak{a}, & c'(0 \leq 1) &= M. \end{aligned}$$

Then the Grothendieck constructions amount to arrow categories:

$$\tilde{\mathcal{W}}' = (\mathcal{V} \rightarrow_S \mathcal{U}); \quad \tilde{c}' = (\mathfrak{b} \rightarrow_M \mathfrak{a}).$$

We can generalize the arrow category in the following way. Consider the path category

$$\mathcal{C} = \langle 0 \xrightarrow{\leq} 1 \xrightarrow{\leq} \cdots \xrightarrow{\leq} n-1 \xrightarrow{\leq} n \rangle.$$

We will call a category isomorphic to a such a category \mathcal{C} for some n a *chain category*. Then a pseudofunctor

$$(\mathcal{W}, c) : \mathcal{C} \rightarrow \underline{\text{Map}},$$

takes values

$$\begin{aligned} \mathcal{W}_i, \quad \mathcal{W}(i \leq j) &= S_{ij}, \\ \mathfrak{c}_i, \quad \mathfrak{c}(i \leq j) &= M_{ij}, \end{aligned}$$

and is equipped with isomorphisms

$$(\phi_{kji}, F_{kji}) : (S_{jk} \circ S_{ij}, M_{jk} \otimes_{\mathfrak{a}_j} M_{ij}) \longrightarrow (S_{ik}, M_{ik}).$$

Example 6.8. Choosing arbitrary graded categories $(\mathcal{W}_i, \mathfrak{c}_i)$, bifunctors $S_{i,i+1}$, and bimodules $M_{i,i+1}$, it is possible to define the remaining data as generalized compositions and tensor products

$$S_{ij} = S_{j-1,j} \circ \cdots \circ S_{i,i+1}$$

and

$$M_{ij} = M_{j-1,j} \otimes \cdots \otimes M_{i,i+1}. \quad \square$$

Now let $\mathcal{C}^{ij} : \mathcal{C}^{ij} \subseteq \mathcal{C}$ be the subcategory

$$\mathcal{C}^{ij} = \langle \begin{array}{ccccccc} i & \xrightarrow{\quad} & i+1 & \xrightarrow{\quad} & \cdots & \xrightarrow{\quad} & j-1 & \xrightarrow{\quad} & j \\ \leq & & \leq & & & & \leq & & \leq \end{array} \rangle,$$

and put $(\mathcal{W}^{ij}, \mathfrak{c}^{ij}) = (\mathcal{W}^{\Phi^{ij}}, \mathfrak{c}^{\Phi^{ij}})$. We thus obtain natural injective Cartesian map-graded functors $\tilde{\Phi}^{ij} : (\tilde{\mathcal{W}}^{ij}, \tilde{\mathfrak{c}}^{ij}) \longrightarrow (\tilde{\mathcal{W}}, \tilde{\mathfrak{c}})$ as in Section 6.5.

On \mathcal{C}^{01} , we can define a new pseudofunctor

$$(\mathcal{W}', \mathfrak{c}') : \mathcal{C}^{01} \longrightarrow \underline{\text{Map}},$$

with

$$\begin{aligned} \mathcal{W}'_0 &= \mathcal{W}_0, \quad \mathcal{W}'_1 = \mathcal{W}^{1n}, \quad \mathcal{W}(0 \leq 1) = S, \\ \mathfrak{c}'_0 &= \mathfrak{c}_0, \quad \mathfrak{c}'_1 = \mathfrak{c}^{1n}, \quad \mathfrak{c}'(0 \leq 1) = M, \end{aligned}$$

where S is the natural $\tilde{\mathcal{W}}^{1n}\text{-}\mathcal{W}_0$ -bimodule and M the natural $\tilde{\mathfrak{c}}^{ij}\text{-}S\text{-}\mathfrak{c}_0$ bimodule determined by $\tilde{\Phi}^{1n}$, $\tilde{\Phi}^{00}$ and the identity bimodule on $(\tilde{\mathcal{W}}, \tilde{\mathfrak{c}})$ (we have $(\tilde{\mathcal{W}}^{00}, \tilde{\mathfrak{c}}^{00}) \cong (\mathcal{W}_0, \mathfrak{c}_0)$). Thus, we

have

$$S(0, i) = S_{0i}, \quad M(0, i) = M_{0i}.$$

It is readily seen, for instance, using the analysis from Section 5.8, that

$$(\tilde{\mathcal{W}}, \tilde{\tau}') \cong (\tilde{\mathcal{W}}', \tilde{\tau}').$$

By induction on n , we can thus apply the results of Section 5.7.

Alternatively, we can take a global approach. Let $\Phi : \text{ob}\mathcal{C} \subseteq \mathcal{C}$ be the subcategory consisting of the objects and identity morphisms of \mathcal{C} . There is a corresponding injective Cartesian morphism $\tilde{\Phi} : (\tilde{\mathcal{W}}^\Phi, \tilde{\tau}^\Phi) \longrightarrow (\tilde{\mathcal{W}}, \tilde{\tau})$. For every individual object $i \in \mathcal{C}$, we have a further subcategory $\Phi^i : i \subseteq \text{ob}\mathcal{C}$ corresponding to $\tilde{\Phi}^i : (\mathcal{W}_i, c_i) \longrightarrow (\tilde{\mathcal{W}}^\Phi, \tilde{\tau}^\Phi)$. Clearly, the complement of $\tilde{\mathcal{W}}^\Phi \subseteq \tilde{\mathcal{W}}$ constitutes an ideal which we denote by \mathcal{S} . From (12), we thus obtain an exact sequence of Hochschild complexes:

$$0 \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}}(\tilde{\tau}, (1_{\tilde{\tau}})_{\mathcal{S}}) \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}}(\tilde{\tau}) \longrightarrow \bigoplus_{i=0}^n \mathbf{C}_{\mathcal{W}_i}(c_i) \longrightarrow 0.$$

Now consider a general pseudofunctor $(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \longrightarrow \underline{\text{Map}}$. We can use the internal structure of \mathcal{C} in order to describe $(\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ as an arrow category. Let \mathcal{Z} be a thin ideal in \mathcal{C} with associated subcategories $\mathcal{C}_0 \subseteq \mathcal{C}$ of “objects below \mathcal{Z} ” and \mathcal{C}_1 of “objects above \mathcal{Z} ” as described in Section 5.8 and suppose $\text{Ob}(\mathcal{C}) = \text{Ob}(\mathcal{C}_0) \cup \text{Ob}(\mathcal{C}_1)$ such that $\mathcal{C} \cong (\mathcal{C}_0 \rightarrow_{\mathcal{Z}} \mathcal{C}_1)$ for the natural restriction of $1_{\mathcal{C}}$ to a \mathcal{C}_1 - \mathcal{C}_0 -bifunctor \mathcal{Z} . The inclusions $\mathcal{C}_0 \subseteq \mathcal{C}$ and $\mathcal{C}_1 \subseteq \mathcal{C}$ give rise to injective Cartesian graded functors

$$(\tilde{\mathcal{U}}_0, \tilde{\mathfrak{a}}_0) \longrightarrow (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}}), \quad (\tilde{\mathcal{U}}_1, \tilde{\mathfrak{a}}_1) \longrightarrow (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}}).$$

Consider the natural restriction of $1_{\tilde{\mathcal{U}}}$ to a $\tilde{\mathcal{U}}_1$ - $\tilde{\mathcal{U}}_0$ -bifunctor T and the natural restriction of $1_{\tilde{\mathfrak{a}}}$ to an $\tilde{\mathfrak{a}}_1$ - T - $\tilde{\mathfrak{a}}_0$ -bimodule N . Put $\tilde{\mathcal{Z}} = \coprod S_{\mathcal{Z}}(U', U)$ running over $z : \mathcal{C}' \longrightarrow \mathcal{C}$ in \mathcal{Z} , $U \in \mathcal{U}_{\mathcal{C}}$, $U' \in \mathcal{U}_{\mathcal{C}'}$.

Proposition 6.9. The thin ideal $\tilde{\mathcal{Z}}$ in $\tilde{\mathcal{U}}$ gives rise to isomorphisms

$$\tilde{\mathcal{U}} \cong (\tilde{\mathcal{U}}_0 \rightarrow_T \tilde{\mathcal{U}}_1), \quad \tilde{\mathfrak{a}} \cong (\tilde{\mathfrak{a}}_0 \rightarrow_N \tilde{\mathfrak{a}}_1). \quad \square$$

Remark 6.10. Note that a generalized arrow category over

$$\mathcal{C} = \langle 0 \xrightarrow{\leq} 1 \xrightarrow{\leq} 2 \rangle$$

is used in the proof of [12, Section 4.6, Theorem (d)]. □

6.7 Covers by arrow categories

The example from Section 6.6 suggests how we can view some map-graded categories as being assembled from the primary arrow category construction, which can be seen as a certain way of glueing categories by means of a bimodule. Another way to glue map-graded categories is along covers of underlying grading categories as described in Section 2.6. More generally, we obtained a sheaf of Hochschild complexes on Map_{sc} in Section 4.3 and in particular Mayer–Vietoris sequences in Section 4.5. In this section, we explain how these different tools can be combined.

Let (\mathcal{C}, \leq) be (the category associated to) a finite poset. For elements $a, b \in \mathcal{C}$, we denote $a \sqsubset b$ if $a \leq b$ and if $a \leq c \leq b$ for $c \in \mathcal{C}$ implies $a = c$ or $c = b$. Let t_0, \dots, t_n be the maximal elements of \mathcal{C} and s_0, \dots, s_m be the minimal elements. For every composition chain

$$s_i = c_0 \sqsubset c_1 \sqsubset \dots \sqsubset c_k \sqsubset c_{k+1} \sqsubset \dots \sqsubset c_p = t_j,$$

we obtain a generated subcategory which is a chain category in the sense of Section 6.6. Every chain of elements

$$a_0 \leq a_1 \leq a_2 \leq \dots \leq a_l$$

can be refined and fitted into a composition chain

$$s_i = c_0 \sqsubset \dots \sqsubset c_n = a_0 \sqsubset \dots \sqsubset c_m = a_l \sqsubset c_{m+1} \sqsubset \dots \sqsubset c_p = t_j,$$

containing all the elements a_k . Thus, the collection of subcategories $\mathcal{D} \subseteq \mathcal{C}$ generated by composition chains of \mathcal{C} constitutes an ∞ -cover of \mathcal{C} . Clearly, the intersection of two such categories is itself a chain category, of strictly smaller length.

Let $(\Phi_i : \mathcal{D}_i \subseteq \mathcal{C})_{i \in I}$ be the collection of chain categories generated by composition chains of \mathcal{C} . Put $\mathcal{D}_{ij} = \mathcal{D}_i \cap \mathcal{D}_j$ and $\Phi_{ij} : \mathcal{D}_{ij} \rightarrow \mathcal{C}$, $\Phi_{ij}^i : \mathcal{D}_{ij} \rightarrow \mathcal{D}_i$.

Now consider a pseudofunctor

$$(\mathcal{W}, \mathbf{c}) : \mathcal{C} \longrightarrow \underline{\text{Map}}.$$

Put $(\mathcal{W}_i, \mathbf{c}_i) = (\mathcal{W}^{\phi_i}, \mathbf{c}^{\phi_i})$ and $(\mathcal{W}_{ij}, \mathbf{c}_{ij}) = (\mathcal{W}^{\phi_{ij}}, \mathbf{c}^{\phi_{ij}})$. We obtain a pullback diagram of injective Cartesian map-graded functors

$$\begin{array}{ccc} (\tilde{\mathcal{W}}_i, \tilde{\mathbf{c}}_i) & \xrightarrow{\tilde{\phi}_i} & (\tilde{\mathcal{W}}, \tilde{\mathbf{c}}) \\ \tilde{\phi}_{ij}^i \uparrow & & \uparrow \tilde{\phi}_j \\ (\tilde{\mathcal{W}}_{ij}, \tilde{\mathbf{c}}_{ij}) & \xrightarrow{\tilde{\phi}_{ij}^j} & (\tilde{\mathcal{W}}_j, \tilde{\mathbf{c}}_j) \end{array} \quad (17)$$

By Proposition 6.5, the $\tilde{\phi}_i$ constitute an ∞ -cover in $\underline{\text{Map}}$ and thus we can use Theorem 4.7 to relate the different Hochschild complexes. By the sheaf property, we obtain an exact sequence

$$0 \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}}(\tilde{\mathbf{c}}) \longrightarrow \bigoplus_{i \in I} \mathbf{C}_{\tilde{\mathcal{W}}_i}(\tilde{\mathbf{c}}_i) \longrightarrow \bigoplus_{\{i, j\}} \mathbf{C}_{\tilde{\mathcal{W}}_{ij}}(\tilde{\mathbf{c}}_{ij}).$$

Alternatively, we can proceed inductively by first isolating one composition chain category \mathcal{D}_0 and defining \mathcal{D}_1 to be the subcategory generated by all the other composition chains. Both \mathcal{D}_1 and $\mathcal{D}_{01} = \mathcal{D}_0 \cap \mathcal{D}_1$ are covered by strictly fewer composition chain categories than \mathcal{D} , and the occurring chains do not increase in length. Proceeding in a similar fashion as before, we then obtain a Mayer–Vietoris sequence as in Section 4.5:

$$0 \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}}(\tilde{\mathbf{c}}) \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}_0}(\tilde{\mathbf{c}}_0) \oplus \mathbf{C}_{\tilde{\mathcal{W}}_1}(\tilde{\mathbf{c}}_1) \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}_{01}}(\tilde{\mathbf{c}}_{01}) \longrightarrow 0.$$

Remark 6.11. The approach discussed in this section can be extended from a poset to a *delta*, that is, a category \mathcal{C} in which the arrows go only one way: for $C, C' \in \mathcal{C}$, we have $\mathcal{C}(C, C') = \emptyset$ or $\mathcal{C}(C', C) = \emptyset$. \square

Example 6.12. Put $\mathcal{C} = \{s, t_0, t_1\}$ with $s \leq t_0$ and $s \leq t_1$ and consider $(\mathcal{W}, \mathbf{c}) : \mathcal{C} \longrightarrow \underline{\text{Map}}$ with

$$\mathcal{W}(s) = \mathcal{W}_s, \quad \mathcal{W}(t_0) = \mathcal{W}_0, \quad \mathcal{W}(t_1) = \mathcal{W}_1, \quad \mathcal{W}(s \leq t_0) = \mathcal{S}_0, \quad \mathcal{W}(s \leq t_1) = \mathcal{S}_1,$$

$$\mathbf{c}(s) = \mathbf{c}_s, \quad \mathbf{c}(t_0) = \mathbf{c}_0, \quad \mathbf{c}(t_1) = \mathbf{c}_1, \quad \mathbf{c}(s \leq t_0) = \mathbf{M}_0, \quad \mathbf{c}(s \leq t_1) = \mathbf{M}_1.$$

The category \mathcal{C} is covered by the two chain categories

$$\mathcal{D}_0 = \langle s \leq t_0 \rangle, \quad \mathcal{D}_1 = \langle s \leq t_1 \rangle,$$

and we have $\mathcal{D}_{01} = \mathcal{D}_0 \cap \mathcal{D}_1 = \langle s \rangle$. We denote the inclusions by $\Phi_i, \Phi_{ij}, \Phi_{ij}^i$ as before. The diagram (17) is given by

$$\begin{array}{ccc} (\mathcal{W}_s \rightarrow_{S_0} \mathcal{W}_0, \mathfrak{c}_s \rightarrow_{M_0} \mathfrak{c}_0) & \xrightarrow{\tilde{\Phi}_0} & (\tilde{\mathcal{W}}, \tilde{\mathfrak{c}}) \\ \tilde{\Phi}_{01}^0 \uparrow & & \uparrow \tilde{\Phi}_1 \\ (\tilde{\mathcal{W}}_s, \tilde{\mathfrak{c}}_s) & \xrightarrow{\tilde{\Phi}_{01}^1} & (\mathcal{W}_s \rightarrow_{S_1} \mathcal{W}_1, \mathfrak{c}_s \rightarrow_{M_1} \mathfrak{c}_1) \end{array} \tag{18}$$

and we obtain a Mayer-Vietoris sequence

$$0 \longrightarrow \mathbf{C}_{\tilde{\mathcal{W}}}(\tilde{\mathfrak{c}}) \longrightarrow \bigoplus_{i=0}^1 \mathbf{C}_{\mathcal{W}_s \rightarrow_{S_i} \mathcal{W}_i}(\mathfrak{c}_s \rightarrow_{M_i} \mathfrak{c}_i) \longrightarrow \mathbf{C}_{\mathcal{W}_s}(\mathfrak{c}_s) \longrightarrow 0.$$

Note that this example also fits into the setup from Proposition 6.7, with $C_0 = t_0, C_1 = t_1$, and $C_0 \times C_1 = s$. □

6.8 Comparison of pseudofunctors

Let \mathcal{C} be a delta (see Definition 5.10). Now let us start from a pseudofunctor

$$(\mathcal{U}, \mathfrak{a}) : \mathcal{C} \longrightarrow \text{Map}_{\text{sc}}.$$

For every $c : C' \longrightarrow C$, we have an associated sub-Cartesian graded functor

$$(\varphi_c, F_c) : (\mathcal{U}_{C'}, \mathfrak{a}_{C'}) \longrightarrow (\mathcal{U}_C, \mathfrak{a}_C),$$

with associated \mathcal{U}_C - $\mathcal{U}_{C'}$ -bifunctor $S_c = S_{\varphi_c}$ and \mathfrak{a}_C - S_{φ_c} - $\mathfrak{a}_{C'}$ -bimodule $M_c = M_{\varphi_c}$ from Example 3.2. Let \mathcal{C}^* be the category \mathcal{C} with terminal object $*$ and morphisms $*_C : C \longrightarrow *$ attached as described in Section 6.5 and let $(\mathcal{U}^*, \mathfrak{a}^*) : \mathcal{C}^* \longrightarrow \text{Map}_{\text{sc}}$ be any pseudofunctor such that the restriction to \mathcal{C} equals $(\mathcal{U}, \mathfrak{a})$.

Example 6.13. We can consider $(\mathcal{U}^*, \mathfrak{a}^*) : \mathcal{C}^* \rightarrow \text{Map}_{\text{sc}}$ with $(\mathcal{U}_C^*, : \mathcal{C}) \mapsto (\mathcal{U}_C, \mathfrak{a}_C)$ with $(\mathcal{U}_*^*, \mathfrak{a}_*^*) = (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}})$ and the graded functors $(\mathcal{U}_C, \mathfrak{a}_C) \rightarrow (\tilde{\mathcal{U}}|_C, \tilde{\mathfrak{a}}|_C)$ induced by the natural functors $e \rightarrow \mathcal{C}^* : * \rightarrow \mathcal{C}$. \square

As described in Section 6.5, we also obtain an associated pseudofunctor

$$(\mathcal{U}^*, \mathfrak{a}^*) : \mathcal{C}^* \rightarrow \text{Map}_{\text{sc}},$$

with sub-Cartesian graded functors

$$(\tilde{\varphi}_C, \tilde{F}_C) : (\tilde{\mathcal{U}}|_{C'}, \tilde{\mathfrak{a}}|_{C'}) \rightarrow (\tilde{\mathcal{U}}|_C, \tilde{\mathfrak{a}}|_C)$$

and

$$(\tilde{\varphi}_{*C}, \tilde{F}_{*C}) : (\tilde{\mathcal{U}}|_C, \tilde{\mathfrak{a}}|_C) \rightarrow (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}}).$$

In this section, we compare the Hochschild complexes associated to $(\mathcal{U}^*, \mathfrak{a}^*)$ and $(\mathcal{U}^*, \mathfrak{a}^*)$. For every $C \in \mathcal{C}$, the functor $e \rightarrow \mathcal{C}/\mathcal{C} : * \mapsto (1_C : \mathcal{C} \rightarrow \mathcal{C})$ gives rise to a natural graded functor

$$(\varphi_C, F_C) : (\mathcal{U}_C, \mathfrak{a}_C) \rightarrow (\tilde{\mathcal{U}}|_C, \tilde{\mathfrak{a}}|_C).$$

Note that $(\tilde{\varphi}_C, \tilde{F}_C)(\varphi_{C'}, F_{C'}) \neq (\varphi_C, F_C)(\varphi_C, F_C)$. Let $(\theta, H) : (\tilde{\mathcal{U}}, \tilde{\mathfrak{a}}) \rightarrow (\mathcal{U}_*^*, \mathfrak{a}_*^*)$ be the natural sub-Cartesian functor with

$$(\theta, H)(\tilde{\varphi}_{*C}, \tilde{F}_{*C})(\varphi_C, F_C) = (\varphi_{*C}^*, F_{*C}^*).$$

By Example 5.11, the Grothendieck construction of $(\mathcal{U}^*, \mathfrak{a}^*)$ can be described as an arrow category, yielding a commutative diagram

$$\begin{array}{ccc}
 \mathbf{C}_{\tilde{\mathcal{U}}^*}(\tilde{\mathfrak{a}}^*) & \xrightarrow{\cong} & \mathbf{C}_{\mathcal{U}_*^*}(\mathfrak{a}_*^*) \\
 \varphi_{\tilde{\mathcal{U}}}^* \downarrow & \swarrow H^* & \downarrow (F_{*C}^*)^* \\
 \mathbf{C}_{\tilde{\mathcal{U}}}(\tilde{\mathfrak{a}}) & & \mathbf{C}_{\mathcal{U}_C}(\mathfrak{a}_C) \\
 (\tilde{F}_{*C})^* \downarrow & \nearrow F_C^* & \\
 \mathbf{C}_{\tilde{\mathcal{U}}|_C}(\tilde{\mathfrak{a}}|_C) & &
 \end{array} \tag{19}$$

We will prove the following theorem.

Theorem 6.14. (1) For every $c: C' \rightarrow C$ in \mathcal{C} , the natural diagram

$$\begin{array}{ccc}
 \mathbf{G}_{\tilde{\mathcal{U}}|_C}(\tilde{\mathfrak{a}}|_C) & \xrightarrow{F_c^*} & \mathbf{G}_{\mathcal{U}_C}(\mathfrak{a}_C) \\
 \tilde{F}_c^* \downarrow & & \downarrow F_c^* \\
 \mathbf{G}_{\tilde{\mathcal{U}}|_{C'}}(\tilde{\mathfrak{a}}|_{C'}) & \xrightarrow{F_{c'}^*} & \mathbf{G}_{\mathcal{U}_{C'}}(\mathfrak{a}_{C'})
 \end{array} \tag{20}$$

is commutative in the homotopy category of B_∞ -algebras, and the horizontal arrows are quasi-isomorphism.

- (2) The horizontal arrows naturally give rise to a morphism between pseudo-functors $\mathcal{C} \rightarrow B_\infty$:

$$\mathbf{G}_{\mathcal{U}^*|_C}(\mathfrak{a}^*|_C) \rightarrow \mathbf{G}_{\mathcal{U}}(\mathfrak{a}).$$

- (3) If the canonical morphism

$$\omega: 1_{\mathfrak{a}^*} \rightarrow \mathrm{RHom}_{\mathfrak{a}^{\mathrm{op}}}(M_H, M_H)$$

induces a quasi-isomorphism $\mathbf{G}_{\mathcal{U}^*}(\mathfrak{a}^*, \omega)$, then H^* is a quasi-isomorphism and we obtain a morphism of pseudofunctors $\mathcal{C}^* \rightarrow \mathrm{ho}(B_\infty)$:

$$\mathbf{G}_{\mathcal{U}^*}(\mathfrak{a}^*) \rightarrow \mathbf{G}_{\mathcal{U}^*}(\mathfrak{a}^*),$$

in which the top horizontal arrow is given by $(H^*)^{-1}$. This holds in particular is we choose $(\mathcal{U}^*, \mathfrak{a}^*)$ as in Example 6.13. □

As a consequence of the theorem, suppose that the conditions of Proposition 6.7 are fulfilled and $\mathbf{G}_{\mathcal{U}^*}(\mathfrak{a}^*): \mathcal{C}^* \rightarrow B_\infty$ is a sheaf of Hochschild complexes. Then exact sequences of Hochschild complexes following from the sheaf property for $\mathbf{G}_{\mathcal{U}^*}^\bullet(\mathfrak{a}^*)$ naturally translate to exact triangles in terms of $\mathbf{G}_{\mathcal{U}^*}^\bullet(\mathfrak{a}^*)$ in which we were originally interested. In particular, we naturally obtain Mayer–Vietoris exact triangles and their induced long exact cohomology sequences. Before proving the theorem, we start with a key example.

Example 6.15. The first example is in the setup for the original arrow category, as described in the beginning of Section 6.6. So $\mathcal{C} = \{0, 1\}$ with $0 \leq 1$. We consider $(\mathcal{W}, \mathcal{C})$:

$\mathcal{C} \longrightarrow \text{Map}_{\text{sc}}$ determined by $(\mathcal{W}_0, c_0) = (\mathcal{V}, \mathfrak{b})$, $(\mathcal{W}_1, c_1) = (\mathcal{U}, \mathfrak{a})$ and a single sub-Cartesian graded functor

$$(\varphi, F) : (\mathcal{V}, \mathfrak{b}) \longrightarrow (\mathcal{U}, \mathfrak{a}).$$

Using the bifunctor S_φ and the bimodule M_F , we obtain the arrow category

$$(\tilde{\mathcal{W}}, \tilde{c}) = (\mathcal{V} \rightarrow_\varphi \mathcal{U}, \mathfrak{b} \rightarrow_F \mathfrak{a}).$$

Obviously, $\mathcal{C}/1 \cong \mathcal{C}$, $\mathcal{C}/0 \cong e$, and the diagram (20) reduces to

$$\begin{array}{ccc} \mathbf{C}_{\mathcal{V} \rightarrow_\varphi \mathcal{U}}(\mathfrak{b} \rightarrow_F \mathfrak{a}) & \xrightarrow{\varphi_{\mathcal{U}}^*} & \mathbf{C}_{\mathcal{U}}(\mathfrak{a}) \\ \varphi_{\mathcal{V}}^* \downarrow & & \downarrow F^* \\ \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) & \xrightarrow{1} & \mathbf{C}_{\mathcal{V}}(\mathfrak{b}) \end{array}$$

The fact that $\varphi_{\mathcal{U}}^*$ is a quasi-isomorphism and the diagram commutes in the homotopy category of B_∞ -algebras is the content of Proposition 5.16. □

Proof. We will concentrate on the proof of (1). The proof of (3) then follows from diagram (19) in which $\varphi_{\mathcal{U}}^*$ becomes a quasi-isomorphism by Theorem 5.14.

Let $C \in \mathcal{C}$ be fixed. The category \mathcal{C}/C is a delta with terminal object $1_C : C \longrightarrow C$. Let $e_C \subseteq \mathcal{C}/C$ be the category with single object 1_C and single morphism 1_C . Let $\mathcal{D}_C \subseteq \mathcal{C}/C$ be the full subcategory consisting of all objects except 1_C . For every $c : C' \longrightarrow C$, there is a unique morphism $c : c \longrightarrow 1_C$ in \mathcal{C}/C . According to Example 5.11, we obtain a thin ideal $\mathcal{Z} = \{c : c \longrightarrow 1_C \mid 1_C \neq c : C' \longrightarrow C\}$ and we have

$$\mathcal{C}/C \cong \mathcal{D}_C \rightarrow_{\mathcal{Z}} e_C,$$

with $Z(c, 1_C) = \{c\}$. The restriction of $(\mathcal{U}|_C, \mathfrak{a}|_C)$ along $e_C \subseteq \mathcal{C}/C$ corresponds to the constant category $(\mathcal{U}_C, \mathfrak{a}_C)$. Denote the restriction of $(\mathcal{U}|_C, \mathfrak{a}|_C)$ along $\mathcal{D}_C \subseteq \mathcal{C}/C$ by $(\mathcal{V}_C, \mathfrak{b}_C)$. Now consider $1_C \neq c : C' \longrightarrow C$. Since \mathcal{C} is a delta, there is a natural factorization $\mathcal{C}/C' \longrightarrow \mathcal{D}_C \longrightarrow \mathcal{C}/C$ from which we obtain a natural factorization of $(\tilde{\varphi}_c, \tilde{F}_c)$:

$$(\tilde{\mathcal{U}}|_{C'}, \tilde{\mathfrak{a}}|_{C'}) \xrightarrow{(\tilde{\varphi}_c^0, \tilde{F}_c^0)} (\tilde{\mathcal{V}}_C, \tilde{\mathfrak{b}}_C) \xrightarrow{(\varphi_{\mathcal{V}}, F_{\mathfrak{b}})} (\tilde{\mathcal{U}}|_C, \tilde{\mathfrak{a}}|_C).$$

We further obtain a natural sub-Cartesian graded functor (ψ, G) fitting into commutative diagrams

$$\begin{array}{ccc}
 (\tilde{\mathcal{V}}_C, \tilde{\mathfrak{b}}_C) & \xrightarrow{(\psi, G)} & (\mathcal{U}_C, \mathfrak{a}_C) \\
 \uparrow (\tilde{\varphi}_c^0, \tilde{F}_c^0)_{(\varphi_{C'}, F_{C'})} & \nearrow (\varphi_c, F_c) & \\
 (\mathcal{U}_{C'}, \mathfrak{a}_{C'}) & &
 \end{array} \tag{21}$$

for $1_C \neq c: C' \rightarrow C$. By Proposition 6.9, we have

$$\tilde{\mathcal{U}}|_C \cong (\tilde{\mathcal{V}}_C \rightarrow_{\psi} \mathcal{U}_C), \quad \tilde{\mathfrak{a}}|_C \cong (\tilde{\mathfrak{b}}_C \rightarrow_G \mathfrak{a}_C),$$

where we use the bimodules S_{ψ} and M_G . By Proposition 5.16 (see also Example 6.15), we have a commutative diagram

$$\begin{array}{ccc}
 \mathbf{C}_{\tilde{\mathcal{U}}|_C}(\tilde{\mathfrak{a}}|_C) & \xrightarrow{F_C^*} & \mathbf{C}_{\mathcal{U}_C}(\mathfrak{a}_C) \\
 \downarrow F_b^* & \swarrow G^* & \\
 \mathbf{C}_{\tilde{\mathcal{V}}_C}(\tilde{\mathfrak{b}}_C) & &
 \end{array}$$

in which F_C^* is a quasi-isomorphism. Composing this diagram with

$$\mathbf{C}_{\tilde{\mathcal{V}}_C}(\tilde{\mathfrak{b}}_C) \xrightarrow{(\tilde{F}_c^0)^*} \mathbf{C}_{\tilde{\mathcal{U}}|_{C'}}(\tilde{\mathfrak{a}}|_{C'}) \xrightarrow{F_{C'}^*} \mathbf{C}_{\mathcal{U}_{C'}}(\mathfrak{a}_{C'}),$$

and making use of (21), we obtain the commutative diagram (20) as desired. ■

Example 6.16. As an application, we look at the setup from [15, Section 7.6]. We consider the poset $\tilde{\Delta} = \{I \mid I \subseteq \{1, 2, \dots, n\}\}$ ordered by reversed inclusion and the subposet $\Delta \subseteq \tilde{\Delta}$ of all $J \neq \emptyset$. We consider a stack $\tilde{\mathcal{S}}$ of Grothendieck categories on $\tilde{\Delta}$ with exact restriction functors and fully faithful right adjoints, and denote \mathcal{S} for its restriction to Δ . We further assume that the conditions (C1) and (C2) listed in [15, Section 7.6] are satisfied. Let $\text{Mod}(\mathcal{S})$ denote the category of presheaf objects in \mathcal{S} . The examples to have in mind are the stack of categories of sheaves of modules on a cover of a ringed space, and the stack of quasi-coherent sheaf categories on a finite affine cover of a separated scheme.

From $\tilde{\mathcal{S}}$, we obtain a pseudofunctor of categories of injectives

$$\epsilon : \tilde{\Delta} \mapsto \text{Cat}(k) : I \mapsto \epsilon_I = \text{Inj}(\tilde{\mathcal{S}}(I)).$$

For $J \subseteq I$, we have a corresponding fully faithful functor

$$F_{IJ} : \epsilon_I \longrightarrow \epsilon_J.$$

Since these functors are fully faithful, there is a corresponding $(e, \epsilon) : \tilde{\Delta} \longrightarrow \text{Map}_{\text{sc}}$ and the results from this section apply. The collection of objects $\{1\}, \{2\}, \dots, \{n\}$ in Δ satisfies the conditions in Proposition 6.7, where further $\Delta^* \cong \tilde{\Delta}$ and the product of subsets I and J is given by $I \cap J$. We denote the pseudofunctor

$$\epsilon^* = (\epsilon|_{\Delta})^* : \Delta^* \longrightarrow \text{Cat}(k) : I \mapsto \tilde{\epsilon}|_I,$$

where $\tilde{\epsilon}|_I$ is the Grothendieck construction of the restriction of ϵ to $\Delta/I = \{J \in \Delta \mid I \subseteq J\}$ and $\tilde{\epsilon}|_* = \tilde{\epsilon}|_{\Delta}$ is the Grothendieck construction of $\epsilon|_{\Delta}$. Thus, we conclude that $\mathbf{C}(\epsilon^*)$ satisfies the sheaf property with respect to the collection of maps $\{i\} \longrightarrow *$ for $i \in \{1, \dots, n\}$. The first part of the proof of [15, Theorem 7.7.1] amounts to the verification of the condition in Theorem 6.14(3). We thus obtain from Theorem 6.14 a morphism of pseudofunctors $\Delta^* \longrightarrow \text{ho}(B_{\infty})$:

$$\mathbf{C}(\epsilon^*) \longrightarrow \mathbf{C}(\epsilon),$$

in which all component maps are quasi-isomorphisms. The Mayer–Vietoris exact triangle for ringed spaces proved in [15, Theorem 7.9.1] becomes an immediate corollary of our theorem, and the proof given in [15] is in fact a special case of the proof of Theorem 6.14. \square

Remark 6.17. The sheaves of Hochschild complexes we obtain in this paper naturally give rise to hypercohomology spectral sequences. However, if we start for instance from a ringed space (X, \mathcal{O}_X) , the site on which an associated sheaf of Hochschild complexes of Grothendieck constructions of categories of injectives lives is fundamentally different from the standard site associated to X . In Example 6.16, this difference is “brigged” by a bimodule between the categories associated to the different suprema associated to a cover of a ringed space (on the one hand, the ringed space and on the other hand, the downset of the cover). A global approach along these lines (possibly combined with techniques from [13]) should lead to new Hochschild cohomology spectral sequences.

The construction of a Hochschild cohomology local-to-global spectral sequence for general ringed spaces based upon map-graded Hochschild cohomology and hypercoverings remains work in progress [18]. \square

Acknowledgements

The author is very grateful to Michel Van den Bergh for the original idea of introducing Hochschild complexes of map-graded categories, as a tool for obtaining a Hochschild cohomology local-to-global spectral sequence for arbitrary ringed spaces, dating back to one of her research stays at the Mittag-Leffler institute in 2004. A write up of the envisaged approach, which involves the theory of hypercoverings, remains a joint work in progress [18]. She also thanks him for drawing her attention to Remark 5.1.

Funding

The author acknowledges the support of the European Union for the ERC grant No 257004-HHNcdMir and the support of the Research Foundation Flanders (FWO) under Grant No G.0112.13N.

References

- [1] *Revêtements étales et groupe fondamental (SGA 1)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)] 3, Société Mathématique de France, Paris, 2003, Séminaire de géométrie algébrique du Bois Marie 1960–61. [Geometric Algebra Seminar of Bois Marie 1960–61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Mathematics 224, Springer, Berlin].
- [2] Bendiffalah, B. and D. Guin. “Cohomologie de l’algèbre triangulaire et applications.” *Journal of Algebra* 282, no. 2 (2004): 513–37.
- [3] Cibils, C., E. Marcos, M. J. Redondo, and A. Solotar. “Cohomology of split algebras and of trivial extensions.” *Glasgow Mathematical Journal* 45, no. 1 (2003): 21–40.
- [4] Gerstenhaber, M. and S. D. Schack. “On the deformation of algebra morphisms and diagrams.” *Transactions of the American Mathematical Society* 279, no. 1 (1983): 1–50.
- [5] Gerstenhaber, M. and S. D. Schack. “Algebraic Cohomology and Deformation Theory.” *Deformation Theory of Algebras and Structures and Applications (Il Ciocco, 1986)*, 11–264. NATO Advanced Science Institutes Series C: Mathematical and Physical Sciences 247. Dordrecht: Kluwer Academic Publishers, 1988.
- [6] Gerstenhaber, M. and S. D. Schack. “The cohomology of presheaves of algebras. I. Presheaves over a partially ordered set.” *Transactions of the American Mathematical Society* 310, no. 1 (1988): 135–65.
- [7] Green, E. L., E. N. Marcos, and N. Snashall. “The Hochschild cohomology ring of a one point extension.” *Communications in Algebra* 31, no. 1 (2003): 357–79.

- [8] Green, E. L. and Ø. Solberg. "Hochschild Cohomology Rings and Triangular Rings." *Representations of Algebra*, vol. I, II, 192–200. Beijing: Beijing Normal University Press, 2002.
- [9] Guccione, J. A. and J. J. Guccione. "Hochschild cohomology of triangular matrix algebras." (2001): preprint arXiv:math/0104068.
- [10] Guil-Asensio, F. and M. Saorín. "On automorphism groups induced by bimodules." *Archiv der Mathematik (Basel)* 76, no. 1 (2001): 12–9.
- [11] Happel, D. "Hochschild Cohomology of Finite-Dimensional Algebras." *Séminaire d'Algèbre Paul Dubreil et Marie-Paul Malliavin, 39ème Année (Paris, 1987/1988)*, 108–26. Lecture Notes in Mathematics 1404. Berlin: Springer, 1989.
- [12] Keller, B. "Derived invariance of higher structures on the Hochschild complex." (2003): preprint, <http://www.math.jussieu.fr/~keller/publ/dih.dvi>.
- [13] Lowen, W. "A sheaf of Hochschild complexes on quasi compact opens." *Proceedings of the American Mathematical Society* 136, no. 9 (2008): 3045–50, preprint arXiv:0707.2605.
- [14] Lowen, W. "Hochschild cohomology of presheaves as map-graded categories." *International Mathematics Research Notices* (2008), Art. ID rnn118, 32.
- [15] Lowen, W. and M. Van den Bergh. "Hochschild cohomology of abelian categories and ringed spaces." *Advances in Mathematics* 198, no. 1 (2005): 172–221.
- [16] Lowen, W. and M. Van den Bergh. "Deformation theory of abelian categories." *Transactions of the American Mathematical Society* 358, no. 12 (2006): 5441–83.
- [17] Lowen, W. and M. Van den Bergh. "A Hochschild cohomology comparison theorem for prestacks." *Transactions of the American Mathematical Society* 363, no. 2 (2011): 969–86.
- [18] Lowen, W. and M. Van den Bergh. "A local-to-global spectral sequence for Hochschild cohomology." in preparation.
- [19] Michelena, S. and M. I. Platzeck. "Hochschild cohomology of triangular matrix algebras." *Journal of Algebra* 233, no. 2 (2000): 502–25.
- [20] Mitchell, B. "Rings with several objects." *Advances in Mathematics* 8, no. 1 (1972): 1–161.
- [21] Swan, R. G. "Hochschild cohomology of projective schemes." *Journal of Pure and Applied Algebra* 110, no. 1 (1996): 57–80.
- [22] Vistoli, A. "The deformation theory of local complete intersections." (1997): preprint alg-geom/9703008.